
# **Change of Variables**

# **Background**

Area of parallelogram = base x perpendicular height = |b||a|sin heta = |a imes b|

#### **Jacobian**



$$J=rac{\partial(x,y)}{\partial(u,v)}=egin{pmatrix}rac{\partial x}{\partial u} & rac{\partial x}{\partial v}\ rac{\partial y}{\partial u} & rac{\partial y}{\partial v} \end{pmatrix}$$

For x = g(u,v) and y = h(u,v)

$$\iint_R f(x,y) dx dy = \iint_G f(g(u,v),h(u,v)) J(u,v) du dv$$

u(x,y) and v(x,y)

$$\Delta u pprox u_x \Delta x + u_y \Delta y$$

$$\Delta v pprox v_x \Delta x + v_y \Delta y$$

$$egin{pmatrix} \Delta u \ \Delta v \end{pmatrix} pprox egin{pmatrix} u_x & u_y \ v_x & v_y \end{pmatrix} egin{pmatrix} \Delta x \ \Delta y \end{pmatrix}$$

The scaling factor for the area is the determinant of the matrix.

$$<\Delta x,0> \to <\Delta u, \Delta v> pprox < U_x \Delta x, V_x \Delta x>$$

$$<0,\Delta y> o <\Delta u,\Delta v> pprox < U_y\Delta y,V_y\Delta y>$$


area' = det () dxdy

#### **Example**

Given region defined by 2x-y=0 , 2x+y=0 and the y axis and x axis. Compute  $\iint_R (4x^2-y^2)^4 dx dy$ 

$$u = 2x - y$$

$$v=2x + y$$



$$u-v = -2y$$

$$\iint_{R} (4x^{2}-y^{2})^{4} dx dy = \int_{0}^{2} \int_{u=-v}^{u=0} ((uv)^{4}) rac{1}{4} du dv$$
 )

### **Example**

Region defined by y = x-1, y=x-2 and y=1, y=2

$$u = x - y$$

v=u


$$\iint_R f(x,y) dx dy = \int_1^2 \int_1^2 f(g(u,v),h(u,v)) (1) du dv$$

## **Example**

Find the scaling factor (dxdy vs dudv)

$$u = 3x - 2y$$

$$v = x + y$$



dA=dxdy dA'=dudv

$$\iint dx dy = \iint \frac{1}{5} du dv$$

### **Example**

Find the area of a circle

$$\iint_R dx dy = \int_{-r}^r \int_{ heta=0}^{ heta=2\pi} r dr d heta = \pi r^2$$

### **Example**

Find the area of an ellipse with semi-axes a,b  $(\x^2 \cot a^2 + \{y^2 \cot b^2 = 1\})$  Change the Variables

Extra close brace or missing open brace

Extra close brace or missing open brace

$$\iint_R dx dy = rac{1}{2} \iint_{u^2+v^2<1} du dv = ab \iint_{u^2+v^2<1} du dv$$

ab (area of unit circle  $=ab\pi$ 

# **Example**

Find the volume of the region under the curve  $z=9-x^2-y^2$   $\int_{x=-3}^{x=3}\int_{y=-\sqrt{9-x^2}}^{\sqrt{9-x^2}}9-x^2-y^2dxdy$   $\int_{r=0}^{r=3}\int_{\theta=0}^{\theta=2\pi}(9-r^2)rdrd\theta=\frac{81\pi}{2}$ 

# **Example**

Compute  $\int_0^1 \int_0^1 x^2 y dx dy$  using a change of Variables u=x v=xy  $\int_0^1 \int_0^1 x^2 y dx dy = \setminus (\int_0^1 \int_v^1 v du dv \setminus)$