

# Pset 7

Darwin Do

April 26, 2022

- (a) Darwin Do
- (b) 919941748
- (c) Collaborators: Graham Stodolski, Raffael Davila
- (d) I have followed the academic integrity and collaboration policy
- (e) Hours: 5

# 1 Strongly Independent Set

To prove that SIS is NP-hard, we show that  $\text{IS} \leq_p \text{SIS}$  where IS is the Independent-Set problem as discussed in lecture.

*Proof.* Let  $G = (V, E)$  be an undirected graph that is a valid input to the IS problem. We will form  $G' = (V', E')$  by replacing every edge  $(u, v) \in E$  with two edges,  $(u, w)$  and  $(w, v)$  where  $w$  is a new node. Then, we connect all the  $w$  nodes together.

If  $G$  is **yes** on the IS problem, then  $G'$  will also be **yes** on the SIS problem with the same  $k$ . This is because any nodes in the independent set  $S$  of size  $\geq k$  will have a distance between each other of at least 2 by definition. Due to the adding of the  $(u, w)$  and  $(w, v)$  edges, this distance will become at least 4. So the nodes in  $S$  are a valid strong independent set in  $G'$  with size  $\geq k$ . Otherwise, if the candidate set  $S$  is **no** for the IS problem on  $G$ ,  $G'$  does not have an independent set of size  $\geq k$ . There will be at least 2 vertices  $u, v \in S$  that are connected and have a distance of 1 between each other. When we create the  $(u, w)$   $(w, v)$  edge pairs in  $G'$ , the distance between  $u$  and  $v$  in  $G'$  will be 2 and break the SIS principle.

If  $G'$  is **yes** on the SIS problem, then  $G$  will also be **yes** on the IS problem with the same  $k$ . Let  $S$  be the strong independent set of size  $k$  in  $G'$ . We know that there can be no  $w$  nodes in  $S$  as all the  $w$  nodes are connected to each other so you can reach any node from a  $w$  node within 2 edges. That means we can remove the  $w$  nodes from  $G'$  and re-connect the  $(u, v)$  nodes to get  $G$ . The nodes in the strong independent set  $S$  are the same nodes in the independent set for  $G$  since none of the vertices in  $S$  have a distance of 1 between them by definition of SIS. Otherwise, if the candidate set  $S$  is **no** for the SIS problem on  $G'$ ,  $G$  does not have an independent set of size  $\geq k$ . In this case, is at least one pair of nodes  $u, v \in S$  whose distance is  $\leq 2$ . If it is 1, then we have already violated the independent set principle. If the distance is 2, then we know the path from  $u$  to  $v$  is  $(u, w) \rightarrow (w, v)$  by construction. Once we remove the  $w$  node and reconnect the  $(u, v)$ , they will be neighbors and violate the independent set principle.

We have shown  $\text{IS} \leq_p \text{SIS}$ . Since IS is NP-hard, SIS is NP-hard as well.  $\square$

Now we show that  $\text{SIS} \in \text{NP}$ .

*Proof.* Let  $S$  be a candidate strongly independent set for graph  $G$ . For all  $v \in S$ , we perform a BFS from  $v$  that only goes 2 levels from  $v$ . If we encounter any other nodes in  $S$  from our 2-level BFS, we know that this candidate is not a strongly independent set. Otherwise if complete this process for all  $v \in S$  and no elements of  $S$  are in the 2-level BFS, we know this is an strongly independent set.

This modified version of BFS is faster than BFS as it only goes 2 levels. The runtime of normal BFS is  $O(|V| + |E|)$ . We are running this modified version of BFS on every node in  $S$ , so the upper bound runtime is  $O((|V| + |E|) \cdot |V|)$

which is polynomial to the graph. Verifying a candidate takes a polynomial amount of time so **SIS** is in NP.  $\square$

**SIS** is in NP and is NP-hard so it is NP-complete.

## 2 Dominating Set

We first show that DS is NP-hard by showing that  $VC \leq_p DS$  where VC is the **Vertex-Cover** problem as discussed in lecture.

*Proof.* Let  $G = (V, E)$  be an undirected graph that is a valid input to the VC problem. We will form  $G' = (V', E')$  by replacing every edge  $(u, v) \in E$  with two edges,  $(u, w)$  and  $(w, v)$  where  $w$  is a new node. Then we connect all the vertices  $v \in V$  together with edges.

If  $G$  is **yes** on the VC problem then there is a vertex set  $S$  with size  $\leq k$  where every edge has at least one vertex in  $S$ . This set  $S$  will also be a valid dominating set in  $G'$ . We know that every edge in  $G$  is touching at least one node in  $S$ . By construction, that means every  $w$  node will be touching a node in  $S$ . Then since we connect all the nodes  $v \in V$  together, we know all  $v \in V$  is either a node in  $S$  or touching a  $S$  node. Every  $v \in V'$  is either in  $S$  or touching a node in  $S$  so  $G'$  is a dominating set of size  $\leq k$ . Otherwise, if set  $S$  is **no** for VC on  $G$ , there will be at least one edge  $(u, v) \in E$  where  $u$  and  $v$  are not in  $S$ . In  $G'$ , the  $w$  node connecting to the  $u, v$  nodes will be connected to 2 nodes not in  $S$  breaking the dominating set principle.

If  $G'$  is **yes** on the DS problem, then  $G$  is also **yes** on the VC problem. Let  $S$  be the dominating set in  $G'$  of size  $\leq k$ . By definition, we know that every  $w$  node is a neighbor of at least 1 node in  $S$ .

When we remove the  $w$  node and the  $(v, w)$ ,  $(w, u)$  edges, we connect  $(v, u)$  again to form the edge in  $E$ . Since we know that at least  $u$  or  $v$  is in  $S$ , the  $(v, u)$  edge follows the vertex-cover principle. We do this step for all edges in  $G'$  that are connected to a  $w$  node. For edges that are not connected to a  $w$  node, (instead connected to each  $v \in V$ ), we simply remove these from  $G'$  to get  $G$  so we don't need to worry about them. Therefore the  $S$  set is a valid vertex cover of size  $\leq k$ . Otherwise, if set  $S$  is **no** for DS on  $G'$ , there will be at least one node  $v^*$  who has 0 neighbors in  $S$ . We know  $v^* \notin V$  as every  $v \in V$  is connected to each other so all  $v \in V$  is either in  $S$  or a neighbor of  $S$  since  $|S| > 0$ . That means  $v^*$  is a  $w$  node with edges  $(u, w)$  and  $(w, v)$  where  $v$  and  $u$  are not in  $S$ . When deleting the  $w$  nodes to form  $G$ , the  $(u, v)$  edge will violate the vertex-cover principle and  $S$  will not be a valid vertex-cover for  $G$ . □

Now we show  $DS \in NP$

*Proof.* Let  $S$  be a candidate dominating set for graph  $G$  with size  $\leq k$ . For all  $v \in V$ , we check if either  $v$  is in  $S$  or if one of the neighbors of  $v$  is in  $S$ . If this condition fails, we know that  $S$  is not a valid dominating set. Otherwise if none of the nodes fail this check, we know  $S$  is a valid dominating set of size  $\leq k$  on  $G$ .

The worst-case runtime for this is  $O(|V|^2)$  as we iterate through all the nodes and look at each node's neighbors. This is polynomial time so DS is in NP. □

DS is in NP and is NP-hard so it is NP-complete.