


aRchiteutis preliminary report

Dropped samples

No samples were removed by the read-count floor or the rarefaction-curve check.

composition

Stacked composition. Samples ordered by `fpc`. Within-sample abundances rescaled to 0-1.



donut


Mean composition across samples, drawn as a donut. Relative abundance.



Apilactobacillus Buchnera Candidatus Carsonella Gilliamella Mycobacterium Pseudomonas Staphylococcus Strepto
bacillus Burkholderia Escherichia Klebsiella Plantactinospora Serratia Streptococcus


barplot

Per-taxon distribution across samples. Geometry: `box`.




alpha

Shannon and Simpson on counts. Split by target. Geometry: `box`.




rarefaction

Observed, Shannon and Simpson rarefaction. Thin line per sample, loess by target.




beta

Sample distance heatmap. Method: `bray`. Samples clustered with ward.D2.



heattree


Rank tree coloured by mean relative abundance. `metacoder::heat_tree` when that package and a taxonomy table are available; otherwise a formula tree.



difftree


Log2 fold change between two target levels on a taxonomy tree. ggtree layout when that package is installed; otherwise a rectangular cladogram. A bar at each tip shows the log2 fold change.

log2(pupa / larvae)



upset

Taxon presence intersections across target levels.



This is a preliminary report only, not a final analysis.