LOCAL FACTORS OF THE ADJOINT L-FUNCTION AND THE ℓ-ADIC VALUATIONS OF THE MODULAR DEGREE

DHARMAJ SONI

Abstract. Let $E/\mathbb{Q}$ be the $X_0(N)$-optimal curve of its isogeny class, $\varphi_E\colon X_0(N) \to E$ its modular parametrisation and f its newform. For an odd prime ℓ not dividing N such that the mod ℓ representation of E is absolutely irreducible on $\mathrm{Gal}(\overline{\mathbb{Q}}/\mathbb{Q}(\sqrt{\ell^*}))$ (for $\ell \geq 5$ this only asks that $E[\ell]$ be irreducible), we prove

$$v_\ell(\deg \varphi_E) \;=\; \mathrm{length}_{\mathbb{Z}_\ell} H^1_f(\mathbb{Q}, \mathrm{ad}^0\, T_\ell E \otimes \mathbb{Q}_\ell/\mathbb{Z}_\ell) + \sum_{q \mid N} v_\ell\big(\mathrm{Tam}_q(\mathrm{ad}^0\, T_\ell E)\big) + \sum_{q \mid N} e_\ell(E, q),$$

where $e_\ell(E,q)$ is the ℓ-adic valuation of the Euler factor at $s=1$ of the adjoint L-function $L(\mathrm{ad}^0 f, s)$ that the naive adjoint L-function omits (it is nonzero only at the additive primes: for a tame $q \geq 5$ it is $v_\ell(q - \chi_{-3}(q))$ for the Kodaira types $\mathrm{II}, \mathrm{II}^*, \mathrm{IV}, \mathrm{IV}^*$, $v_\ell(q - \chi_{-4}(q))$ for $\mathrm{III}, \mathrm{III}^*$, $v_\ell(q^2-1)$ for I_n^* and $v_\ell(q-1) + v_\ell((q+1)^2 - a_q(E')^2)$ for I_0^*, with E' the twist of good reduction), and the Tamagawa factors are trivial at the potentially good primes and explicit at the potentially multiplicative ones, where their valuation is at least $t_\ell(E,q)$, the ℓ-part of the valuation of the Tate parameter. In particular

$$v_\ell(\deg \varphi_E) \;\geq\; \sum_{q \mid N} e_\ell(E,q) + \sum_{q \mid N} t_\ell(E,q).$$

The proof assembles the theorem of Diamond, Flach and Guo identifying the length of the Σ-imprimitive adjoint Selmer group with the valuation of a congruence ideal, their formula for the variation of that ideal with Σ, the local Tamagawa formula of Fontaine and Perrin-Riou, a computation of the Tamagawa factor of the adjoint at every bad prime, and the observation that for the optimal curve the congruence ideal of Diamond, Flach and Guo is generated by the modular degree, by the projection formula; it also uses an identification of their integral structure with the cohomology of $X_0(N)$, which we prove. The Tamagawa factor raises the bound by at least one whenever a Steinberg prime $q \equiv 1 \pmod{\ell}$ with $\ell \mid n_q$ has a Tate parameter whose unit part is an ℓ-th power. Without the irreducibility over $\mathbb{Q}(\sqrt{\ell^*})$ the terms of the types I_0^* and I_n^* at the odd primes still hold, by a twisting argument and the level-raising formula of Diamond, Flach and Guo, but the inequality itself fails: for $\ell = 3$, among the curves of the tables without complex multiplication, exactly the curves whose mod 3 image is the normaliser of a split Cartan subgroup violate it, by exactly one, in 31%, 23% and 18% of the cases on three ranges, the one being the length of $H^0(\mathbb{Q}, \mathrm{ad}^0\, \rho(1) \otimes \mathbb{Q}_3/\mathbb{Z}_3)$. With that correction term the inequality holds with no exception, with Euler factors computed by PARI at every bad prime, for every curve of conductor at most 500000 that the tables determine to be optimal, with $E[\ell]$ irreducible and without complex multiplication, and for 893 curves outside the tables, at $\ell = 3, 5, 7, 11, 13$, also at primes ℓ dividing N where no theorem applies, with equality in 57% of the cases with a positive bound at $\ell = 3$. We describe the extra terms at $q = \ell = 3$ and the Eisenstein case, where we show that the bound 2 for the deficit at $\ell = 3$ follows from a conjectured bound by the rational torsion.

1. Introduction

Let $E/\mathbb{Q}$ be an elliptic curve of conductor N and let $f = f_E \in S_2(\Gamma_0(N))$ be the newform attached to it. Among the curves in the isogeny class of E there is a distinguished one, the $X_0(N)$-optimal curve, for which the modular parametrisation $\varphi_E\colon X_0(N) \to E$ has minimal

Date: 24 October 2026.

degree; throughout, E denotes this curve and $\deg\varphi_E$ its *modular degree*. When $E[\ell]$ is irreducible, every curve of the class is linked to E by a cyclic isogeny of degree prime to ℓ, so v_ℓ of the degree of its minimal parametrisation is $v_\ell(\deg\varphi_E)$: at such an ℓ the choice of the optimal curve does not matter. The modular degree is computable [9, 48], and its size is tied to deep questions (a suitable polynomial bound on $\deg\varphi_E$ in terms of N is equivalent to a form of the *abc* conjecture, by work of Frey and of Murty; see [2, §2] and the references there), but its prime factorisation is not understood. The purpose of this note is to describe, and for the most part prove, a local-to-global identity: the ℓ-adic valuation of $\deg\varphi_E$ is the length of an adjoint Selmer group plus local invariants of E at the bad primes, and the right invariants are the local factors of the adjoint L-function.

The classical route to the degree is the Petersson norm. Up to explicit factors $\deg\varphi_E$ is $\langle f,f\rangle$ divided by the area of the period lattice of E (Zagier [50], Watkins [48]), and $\langle f,f\rangle$ is the special value of the naive symmetric square L-function at $s=2$ (Rankin [32], Shimura [41]; in the exact normalisation used here, Hida [19, Theorem 5.1] and [12, Lemma 2.12]); Remark 4.15 gives the exact formula. Flach's finiteness theorem for the symmetric square [16] is the first place where the modular degree meets the Selmer group of $\mathrm{Sym}^2 f$. By a theorem of Agashe, Ribet and Stein [2, Theorem 2.2], $\deg\varphi_E$ divides the *congruence number* r_E, the largest integer r such that $f\equiv g \pmod r$ coefficientwise for some integral cusp form g of level N orthogonal to f, with $v_p(\deg\varphi_E)=v_p(r_E)$ for every prime p with $p^2\nmid N$. The congruence number is the congruence ideal of f in the full Hecke algebra of level N (Lemma 3.1), and the theory of such ideals, from Hida [19] to Diamond, Flach and Guo [12], expresses them through the adjoint Selmer group and the adjoint L-value. The L-function that appears there is the *naive* one, $L^{\mathrm{nv}}(\mathrm{Sym}^2 f,s)=\zeta^{(N)}(2s-2)\sum_{n\ge1}a_{n^2}(f)n^{-s}$ (Shimura [40]), which has the true local factor at the primes of good or multiplicative reduction and the trivial factor at the additive primes. The point of this note is that the primes at which the naive and the true adjoint L-functions differ are exactly the primes at which the modular degree acquires extra local divisibility, that the amount is the valuation of the omitted factor, and that this is a consequence of the theorems of [12] once they are read for elliptic curves, with a precise boundary at the one hypothesis they need.

1.1. **The local terms.** Let $V=V_\ell E$, $T=T_\ell E$, $\rho=\rho_{E,\ell}$, $\bar\rho$ its reduction, and $A_f=\mathrm{ad}^0 V$ the trace-zero adjoint. Throughout, Frob_p is an arithmetic Frobenius at p and Frob_p^{-1} a geometric one; χ_ℓ is the ℓ-adic cyclotomic character and $\bar\chi_\ell$ its reduction; the quadratic character of $\mathbb{Q}(\sqrt{d})$, d a discriminant, is χ_d, so that χ_{-3}, χ_{-4} and, for an odd prime q and $q^*=(-1)^{(q-1)/2}q$, the character χ_{q^*} of conductor q appear below; at $\ell=3$ the cyclotomic character is written χ_3^{cyc}, and $\bar\chi_3^{\mathrm{cyc}}=\chi_{-3}$. We follow the conventions of [12, §1.7.2]: for a prime $p\neq\ell$ let $\delta_p=\dim V^{I_p}$, so $\delta_p=2$, 1, 0 for good, multiplicative, additive reduction; $L_p(A_f,s)=\det(1-\mathrm{Frob}_p^{-1}p^{-s}\mid A_f^{I_p})^{-1}$ is the true local factor, defined with the geometric Frobenius as in [12, §1.1.1], and the naive factor is $L_p^{\mathrm{nv}}(A_f,s)=L_p(A_f,s)$ if $\delta_p>0$ and 1 if $\delta_p=0$. (At $p=\ell$ the definitions use D_{cris}; we need them only for $p\neq\ell$.) The matrices in the proof of Lemma 2.8 are those of the arithmetic Frobenius. Where Frobenius acts on $A_f^{I_p}$ by a sign ±1 (Lemma 2.2, Proposition 2.6(iii)), or with eigenvalues $\alpha/\beta,1,\beta/\alpha$ stable under inversion (type I_0^*), the arithmetic and the geometric Frobenius give the same factor; at a potentially multiplicative prime they do not (eigenvalue q for the arithmetic, q^{-1} for the geometric Frobenius), and the table below, like L_p, uses the geometric one. In the normalisation of the symmetric square, $L_p(\mathrm{Sym}^2 f,s)=L_p(A_f,s-1)$, and PARI's `lfunsympow(E,2)` returns these factors at every prime, including 2 and 3 [28].

Definition 1.1. For a bad prime q put

$$t_\ell(E,q) = \begin{cases} v_\ell(-v_q(j_E)) & \text{if } v_q(j_E) < 0, \\ 0 & \text{otherwise,} \end{cases}$$

and, if $q \neq \ell$,

$$e_\ell(E,q) = v_\ell\Big(\frac{L_q(A_f,1)^{-1}}{L_q^{\mathrm{nv}}(A_f,1)^{-1}}\Big) = v_\ell\Big(\frac{P_q(q^{-2})}{1-a_q(E)^2q^{-2}}\Big),$$

where $P_q(X)$ is the polynomial with $L_q(\mathrm{Sym}^2 f, s) = P_q(q^{-s})^{-1}$, so that X stands for q^{-s} and $P_q(q^{-2}) = L_q(A_f,1)^{-1}$, and where the denominator $1-a_q(E)^2q^{-2} = L_q^{\mathrm{nv}}(A_f,1)^{-1}$ is 1 at an additive prime ($a_q(E)=0$) and the true factor $1-q^{-2}$ at a multiplicative one ($a_q(E)^2=1$). So e_ℓ is the valuation of the Euler factor that the naive L-function omits, and $e_\ell(E,q)=0$ at a multiplicative prime. The term t_ℓ is the ℓ-part of the valuation of the Tate parameter at a potentially multiplicative prime, which is $v_\ell(n)$ for a Kodaira symbol I_n at any q and I_n^* at odd q, while a symbol I_n^* at $q=2$ can have potentially good reduction, and then $t_\ell = 0$. The term t_ℓ is invariant under quadratic twist, and $e_\ell(E,q)$ under quadratic twists unramified at q; a twist ramified at q can change $e_\ell(E,q)$ (the curve 11a1 has type I_5 at 11 and $e_3=0$, its twist 121d2 by χ_{-11} has type I_5^* and $e_3 = v_3(120) = 1$), because the true factor is twist invariant and the naive one is not.

For a tame prime $q \geq 5$ the omitted factor is read off the Kodaira type (Lemma 2.5). With χ_{-3}, χ_{-4} the quadratic characters of $\mathbb{Q}(\sqrt{-3})$ and $\mathbb{Q}(i)$, E' the quadratic twist of E by $\mathbb{Q}(\sqrt{q^*})$, and α, β the roots of $X^2 - a_q(E')X + q$ (for type I_0^*, where E' has good reduction at q):

type at $q \geq 5$	$L_q(A_f,s)^{-1}$	$e_\ell(E,q)$
I_n	$1-q^{-1-s}$	0
I_n^*	$1-q^{-1-s}$	$v_\ell(q^2-1)$
$\mathrm{II}, \mathrm{II}^*, \mathrm{IV}, \mathrm{IV}^*$	$1-\chi_{-3}(q)q^{-s}$	$v_\ell(q-\chi_{-3}(q))$
$\mathrm{III}, \mathrm{III}^*$	$1-\chi_{-4}(q)q^{-s}$	$v_\ell(q-\chi_{-4}(q))$
I_0^*	$(1-\frac{\alpha}{\beta}q^{-s})(1-q^{-s})(1-\frac{\beta}{\alpha}q^{-s})$	$v_\ell(q-1) + v_\ell\big((q+1)^2 - a_q(E')^2\big)$

The row I_n holds at every $q \neq \ell$, $q=2$ included, the rows I_n^* and I_0^* at every odd $q \neq \ell$, and the row IV, IV* at every $q \neq 3, \ell$ (Lemmas 2.2 and 2.5); at every bad prime, 2 and 3 included, P_q has one of four shapes (Proposition 2.6).

1.2. **The theorem.** For a prime $p \neq \ell$ and a $G_{\mathbb{Q}_p}$-stable $\mathbb{Z}_\ell$-lattice M the *Tamagawa factor* is $\mathrm{Tam}_p(M) = \mathrm{Fitt}_{\mathbb{Z}_\ell}\big((H^1(I_p,M)_{\mathrm{tor}})^{G_{\mathbb{Q}_p}}\big)$, an ideal of $\mathbb{Z}_\ell$ [12, p. 708], [17, I.4.2.2]; for $M = \mathrm{ad}^0 T$ at a prime of potentially multiplicative reduction its valuation is at least $t_\ell(E,p)$, with an explicit excess (Lemma 2.8), at a prime of potentially good additive reduction it is 0 (Lemma 2.9), and at a prime of good reduction it is 0. Let $H^1_f(\mathbb{Q}, \mathrm{ad}^0 T \otimes \mathbb{Q}_\ell/\mathbb{Z}_\ell)$ be the Bloch-Kato Selmer group [12, §2.1].

Theorem 1.2. *Let E be $X_0(N)$-optimal and let ℓ be an odd prime not dividing $2N$ such that $\bar\rho_{E,\ell}$ is absolutely irreducible when restricted to* $\mathrm{Gal}(\overline{\mathbb{Q}}/\mathbb{Q}(\sqrt{\ell^*}))$, $\ell^* = (-1)^{(\ell-1)/2}\ell$. *Then*

$$v_\ell(\deg\varphi_E) \;=\; \mathrm{length}_{\mathbb{Z}_\ell} H^1_f(\mathbb{Q}, \mathrm{ad}^0 T \otimes \mathbb{Q}_\ell/\mathbb{Z}_\ell) \;+\; \sum_{q\mid N} v_\ell\big(\mathrm{Tam}_q(\mathrm{ad}^0 T)\big) \;+\; \sum_{q\mid N} e_\ell(E,q). \tag{1}$$

In particular

$$v_\ell(\deg\varphi_E) \;\geq\; \sum_{q\mid N} e_\ell(E,q) + \sum_{q\mid N} t_\ell(E,q). \tag{2}$$

The Tamagawa sum is explicit (Corollary 4.14), so (1) says that the excess of $v_\ell(\deg\varphi_E)$ over an explicit local bound is the length of the adjoint Selmer group. The hypothesis is the one under which Diamond, Flach and Guo prove the Tamagawa number conjecture for the adjoint motive of f [12, Theorem 0.1]; in their notation it says that ℓ is not in the exceptional set S_f, defined in Section 4.2. For $\ell \nmid 2N$ it holds if and only if $E[\ell]$ is irreducible and, when $\ell = 3$, the mod 3 image is not the normaliser of a split Cartan subgroup (Proposition 4.8). Theorem 1.2 is an assembly of their results. Their Theorem 3.7 identifies the length of the Σ-imprimitive adjoint Selmer group, Σ the set of bad primes, with the valuation of the naive Σ-finite congruence ideal; their Proposition 1.4(c) says that this ideal is the congruence ideal at level N times the naive Euler factors at the multiplicative primes; their Lemma 2.1 compares the Σ-imprimitive Selmer group with the Bloch-Kato one, the local indices being, by the formula of Fontaine and Perrin-Riou, the true Euler factors times the Tamagawa factors; and the congruence ideal at level N is generated by $\deg\varphi_E$, by the projection formula for φ_E (Proposition 4.4). The naive factors at the multiplicative primes cancel, the true factors at the additive primes are the omitted ones, and (1) appears. One input is not a published statement in the form we need: the identification of the integral structure of [12] with $H^1(X_0(N),\mathbb{Z}_\ell)$ (Proposition 4.1), which we prove; it holds integrally, with no localisation. What this note adds is the identification of the congruence ideal with the modular degree, the reading of the local terms for elliptic curves (Lemmas 2.5, 2.8 and 2.9, Proposition 2.6), the twisting theorems that hold without the hypothesis on $\mathbb{Q}(\sqrt{\ell^*})$ (Theorem 1.6 and Lemma 4.18), the identification of the curves for which that hypothesis fails and of what happens to them (Propositions 1.7 and 5.1, Section 6.7), and the computations, which test (1) well beyond the inequality.

1.3. **Beyond the hypotheses.** The data suggest that the inequality survives the loss of the hypotheses, with two modifications: a weight at the prime ℓ itself when $\ell \mid N$, and a correction term when $\ell = 3$ and the representation is induced from $\mathbb{Q}(\sqrt{-3})$.

Conjecture 1.3. *Let E be $X_0(N)$-optimal and without complex multiplication, let ℓ be an odd prime such that $E[\ell]$ is irreducible, and let $h_\ell(E)$ be the length of $H^0(\mathbb{Q},\mathrm{ad}^0\,T(1)\otimes\mathbb{Q}_\ell/\mathbb{Z}_\ell)$. Then*

$$v_\ell(\deg\varphi_E)\ \geq \sum_{q\mid N,\ q\neq\ell} e_\ell(E,q)\ +\ \sum_{q\mid N} t_\ell(E,q)\ +\ [\ell=3]\,w_3(E)\ -\ h_\ell(E), \tag{3}$$

where $w_3(E)$ is the weight of the prime 3 itself when $3 \mid N$ and the reduction at 3 is potentially good: 1 for the types $\mathrm{II},\mathrm{IV},\mathrm{III}^*$, 2 for $\mathrm{II}^*,\mathrm{IV}^*$ and 0 for $\mathrm{III},\mathrm{I}_0^*$.

The term $h_\ell(E)$ is at most 1; it is positive exactly when $\bar\rho \simeq \bar\rho\otimes\bar\chi_\ell$, which forces $\ell = 3$, where $\bar\chi_3^{\mathrm{cyc}} = \chi_{-3}$; and it vanishes under the hypothesis of Theorem 1.2 (Lemma 4.13). For $\ell = 3$ it is positive precisely when the image of $\bar\rho_{E,3}$ is the normaliser of a split Cartan subgroup, with or without complex multiplication, and then $h_3(E) = 1$ (Proposition 1.7); in the tables this is the image labelled 3Ns (we use the labels of the tables, in the notation of [44], together with their intrinsic description: for 3Ns, a 3-division polynomial that is a product of two quadratics, Proposition 1.7). For the curves of the tables with this image and without complex multiplication the inequality without the correction fails by exactly one in 31%, 23% and 18% of the cases on the three ranges of Section 6, so the hypothesis of Theorem 1.2 is sharp.

Status of the correction. (i) Proved: $h_\ell(E) \leq 1$, with equality exactly when $\ell = 3$ and the mod 3 image is $N_s(\mathbb{F}_3)$ (Lemma 4.13, Proposition 1.7). (ii) Conditional: that the correction is the term which Lemma 2.1 of [12] produces when $H^0(\mathbb{Q},\mathrm{ad}^0\,T(1)\otimes\mathbb{Q}_\ell/\mathbb{Z}_\ell) \neq 0$ assumes $H^1_f(\mathbb{Q},V^D) = 0$, which is not known for these curves, and that $v_\lambda(\eta^\Sigma_{f,\lambda})$ is still the length of the Σ-imprimitive Selmer group (Remark 5.2). (iii) Empirical: the weights w_3 at $q = \ell = 3$

(Section 6.8), and the shortfall pattern, a failure of the uncorrected inequality by exactly one on part of the curves with image 3Ns and on no other curve (Section 6.7).

At $q = \ell \geq 5$ the rule holds with no term, although the data suggest a sharper statement there (Remark 6.1). The terms at the wild primes 2 and 3 are the ones of Definition 1.1 with PARI's Euler factor, and they reproduce every minimum observed in the tables, including the irregular behaviour of the prime 2 (Section 6.8). On the four ranges of Section 6, (3) holds with no exception at $\ell = 3, 5, 7, 11, 13$, including the curves with $\ell \mid N$, which no theorem covers.

1.4. **Local component groups.** The rule contains two simpler statements, in terms of component groups. For a bad prime q let $\Phi_q^{\mathrm{geom}} = \Phi_q(\overline{\mathbb{F}}_q)$ be the group of components of the special fibre of the Néron model of E at q over the algebraic closure. Its order is read off the Kodaira type: n for type I_n; 1 for $\mathrm{II}, \mathrm{II}^*$; 2 for $\mathrm{III}, \mathrm{III}^*$; 3 for $\mathrm{IV}, \mathrm{IV}^*$; 4 for I_0^* and I_n^* [42, IV.9]. The Tamagawa number $c_q = \#\Phi_q(\mathbb{F}_q)$ counts only the components defined over $\mathbb{F}_q$; the statements below are about Φ_q^{geom}, not c_q. For an odd prime q let $E^{(q)}$ be the quadratic twist of E by χ_{q^*}, that is, by $\mathbb{Q}(\sqrt{q^*})$; it has the same reduction type as E at every prime other than q, while at q the Kodaira types are exchanged (Lemma 2.4):

$$\mathrm{I}_n \leftrightarrow \mathrm{I}_n^*, \qquad \mathrm{II} \leftrightarrow \mathrm{IV}^*, \qquad \mathrm{IV} \leftrightarrow \mathrm{II}^*, \qquad \mathrm{III} \leftrightarrow \mathrm{III}^*, \qquad \mathrm{I}_0 \leftrightarrow \mathrm{I}_0^*$$

(all five at every odd q; at $q = 3$ the three middle exchanges keep the conductor exponent, which can exceed 2). Define the *twist-symmetrised* weight

$$w_\ell(E, q) = \begin{cases} v_\ell(n) & \text{if the type at } q \text{ is } \mathrm{I}_n, \text{ or } \mathrm{I}_n^* \text{ with } q \text{ odd,} \\ 1 & \text{if } \ell = 3 \text{ and the type is IV or IV}^*\text{, or II or II}^* \text{ with } q \text{ odd,} \\ 0 & \text{otherwise,} \end{cases}$$

so that

$$w_\ell(E, q) = \max\bigl(v_\ell(\#\Phi_q^{\mathrm{geom}}(E)),\ v_\ell(\#\Phi_q^{\mathrm{geom}}(E^{(q)}))\bigr) \text{ for odd } q, \qquad w_\ell(E, 2) = v_\ell(\#\Phi_2^{\mathrm{geom}}(E))$$

for odd ℓ, by the exchanges above. The two inequalities are

$$v_\ell(\deg \varphi_E) \ \geq \ \sum_{q \mid N} v_\ell\bigl(\#\Phi_q^{\mathrm{geom}}\bigr), \tag{4}$$

$$v_\ell(\deg \varphi_E) \ \geq \ \sum_{q \mid N} w_\ell(E, q), \tag{5}$$

of which the second implies the first. Both hold with no exception on the same ranges, under the hypotheses of Conjecture 1.3 (Section 6.6).

Proposition 1.4. *Under the hypothesis of Theorem 1.2,* (4) *and* (5) *hold; the same is true under Conjecture 1.3 with* $h_\ell(E) = 0$.

Proof. Compare the two sides term by term with (2), respectively (3); $w_\ell(E, q) \geq v_\ell(\#\Phi_q^{\mathrm{geom}})$, so it suffices to treat (5). At a prime of type I_n both sides have $v_\ell(n) = t_\ell(E, q)$. At a prime of type I_n^* with q odd, $t_\ell(E, q) = v_\ell(n) = w_\ell(E, q)$; with $q = 2$, $w_\ell(E, 2) = v_\ell(4) = 0$. At $q \geq 5$ of type $\mathrm{II}, \mathrm{II}^*, \mathrm{IV}, \mathrm{IV}^*$ and $\ell = 3$, $e_3(E, q) = v_3(q - \chi_{-3}(q)) \geq 1 = w_3(E, q)$ since $q \equiv \chi_{-3}(q) \pmod 3$; for $\ell \geq 5$ both weights are 0. At $q = 3$ of these types and $\ell = 3$, $w_3(E) \geq 1 = w_3(E, 3)$ by definition, and for $\ell \geq 5$ the weight is 0. At $q = 2$ of type $\mathrm{II}, \mathrm{II}^*$, $w_\ell(E, 2) = 0$; of type $\mathrm{IV}, \mathrm{IV}^*$ and $\ell = 3$, $w_3(E, 2) = 1 = e_3(E, 2)$ by Lemma 2.2. The types $\mathrm{III}, \mathrm{III}^*, \mathrm{I}_0^*$ have $w_\ell = 0$ for odd ℓ. All other terms of (2) and (3) are nonnegative. □

On the 92,812 working-set curves with type IV or IV* at 2, PARI's factor at 2 is $1 + 2X$ for every one of them, as Lemma 2.2 requires.

1.5. **History, and the statements proved without the hypothesis on $\mathbb{Q}(\sqrt{\ell^*})$.** The multiplicative part of (4) has a long history. Ribet and Takahashi [36, 45] compared, for semistable E, $\deg\varphi_E$ with the degree of a parametrisation by a Shimura curve and found the product of the n_q over the primes of the quaternion discriminant, exactly when E is the only curve in its isogeny class and otherwise up to primes where $E[\ell]$ is reducible; Pasten [30] controls that error term. Pollack and Weston [31] conjectured, and proved for weight 2 and square-free level when $\bar\rho$ is ramified at two primes at least and under their hypothesis CR ($\bar\rho$ surjective and ramified at every prime of the relevant part of the level that is ±1 modulo ℓ) [31, Theorem 6.11], and Kim and Ota [21] proved under the hypotheses recalled in Remark 4.16, that moving a prime q into the "new" part of the level changes the congruence ideal of f_E in the full space $S_2(\Gamma_0(N))$ by exactly the *Tamagawa exponent* t_q, the largest t with $\rho_{E,\ell} \bmod \ell^t$ unramified at q; for multiplicative q one has $t_q = v_\ell(n_q) = t_\ell(E,q)$ whether or not the reduction is split (Lemma 2.1). Theorem 1.2 contains the inequality $v_\ell(\deg\varphi_E) \geq \sum_{q\,\|\,N} t_\ell(E,q)$ for every multiplicative prime, including the primes $q \equiv \pm1 \pmod\ell$ and the prime $\ell = 3$, which the formula of Kim and Ota excludes; it does not contain their equality.

Two statements hold under weaker hypotheses than Theorem 1.2, by level lowering and twisting rather than by the Taylor-Wiles method, and they matter exactly where that method does not apply.

Theorem 1.5. *Let $E/\mathbb{Q}$ have conductor N and let ℓ be an odd prime not dividing N such that $\bar\rho_{E,\ell}$ is irreducible and, if $\ell = 3$, absolutely irreducible when restricted to $\mathrm{Gal}(\overline{\mathbb{Q}}/\mathbb{Q}(\sqrt{-3}))$. Let $q \neq \ell$ be a prime at which E has additive reduction of one of the following kinds:*

(a) *type* IV *or* IV**, with $q \geq 5$ or $q = 2$, and $\ell = 3$;*
(b) *type* II *or* II**, with $q \geq 5$ and $\ell = 3$;*
(c) *type* I_n^* *with $n \geq 1$ and $\ell \mid n$, q odd.*

Then ℓ divides the modular degree of the $X_0(N)$-optimal curve in the isogeny class of E.

For $\ell \geq 5$ Theorem 1.5 needs only the irreducibility of $\bar\rho$; its proof (Section 3) lowers the level of $\bar\rho$, or of its twist $\bar\rho \otimes \chi_{q^*}$, and twists back. The quantitative version of case (c), and of the type I_0^*, is the following, which needs no hypothesis beyond the irreducibility of $E[\ell]$ for the inequality, and gives an equality under the hypothesis of Theorem 1.2.

Theorem 1.6. *Let ℓ be an odd prime with $\ell \nmid 2N$ and $E[\ell]$ irreducible, let S be the set of odd primes q at which E has type I_0^* or I_n^*, and let E_S be the twist of E by $\chi_S = \prod_{q\in S}\chi_{q^*}$. Then*

$$v_\ell(\deg\varphi_E) \;\geq\; v_\ell(r_{E_S}) + \sum_{q\in S} e_\ell(E,q) \;\geq\; \sum_{q\in S} e_\ell(E,q).$$

If moreover $\ell \notin S_f$ (for $\ell \geq 5$ this is automatic, Proposition 4.8), the first inequality is an equality, and $v_\ell(r_{E_S}) = v_\ell(\deg\varphi_{E'_S})$ for the optimal curve E'_S of the isogeny class of E_S:

$$v_\ell(\deg\varphi_E) \;=\; v_\ell(\deg\varphi_{E'_S}) + \sum_{q\in S} e_\ell(E,q).$$

The proof (Section 4.4) uses the formula of [12, Proposition 1.4(c)] for the depleted form of f_{E_S}, which lives at level N, and an elementary transport of congruences along the twist, which identifies $v_\ell(r_E)$ with the Hecke congruence ideal of that depleted form at every prime ℓ outside S, with no other hypothesis on ℓ (Lemma 4.18); under the hypothesis of Theorem 1.2 the identity (1) for E and for E'_S gives the equality, with no multiplicity-one hypothesis.

Proposition 1.7. *Let $E/\mathbb{Q}$ have $E[3]$ irreducible. Then $\bar\rho_{E,3}$ is absolutely irreducible on $\mathrm{Gal}(\overline{\mathbb{Q}}/\mathbb{Q}(\sqrt{-3}))$ unless its image is conjugate to the normaliser $N_s(\mathbb{F}_3)$ of a split Cartan*

subgroup of $\mathrm{GL}_2(\mathbb{F}_3)$*, and in that case* $\bar\rho_{E,3}\otimes\mathbb{F}_9$ *is induced from a character of order* 4 *of* $\mathrm{Gal}(\overline{\mathbb{Q}}/\mathbb{Q}(\sqrt{-3}))$*,* $\bar\rho_{E,3}\simeq\bar\rho_{E,3}\otimes\chi_{-3}$*,* $H^0(\mathbb{Q},\mathrm{ad}^0\,\bar\rho_{E,3}(1))\neq 0$ *and* $h_3(E)=1$*. The* 3*-division polynomial of* E *then factors as a product of two irreducible quadratics, and otherwise it is irreducible.*

The last statement concerns the Eisenstein case, in which (4) can fail.

Conjecture 1.8. *Let* E *be optimal, not CM, with a rational* ℓ*-isogeny,* ℓ *odd, and let* D *be the deficit* $\sum_q v_\ell(\#\Phi_q^{\mathrm{geom}})-v_\ell(\deg\varphi_E)$ *of* (4)*. Then*

$$D\ \le\ v_\ell\big(\#E(\mathbb{Q})_{\mathrm{tors}}\big)+\big[\,E[3]\simeq\mathbb{Z}/3\oplus\mu_3\,\big],$$

where $[\,\cdot\,]$ *is* 1 *if the condition holds and* 0 *otherwise, and also* $D\le$ *the number of curves in the isogeny class of* E *with a rational point of order* ℓ*. In particular* $D\le 0$ *when* E *has no rational* ℓ*-torsion point,* $D\le 1$ *for* $\ell\ge 5$*, and* $D\le 2$ *for* $\ell=3$*, the last because a curve without complex multiplication with* $E[3]\simeq\mathbb{Z}/3\oplus\mu_3$ *has no rational point of order* 9 *(Proposition 6.2).*

Section 2 collects the local facts, Section 3 proves Theorem 1.5, Section 4 proves Theorems 1.2 and 1.6, Section 5 proves Propositions 1.7 and 5.1 and explains the correction term, Section 6 describes the computations and Section 7 discusses the prime 2, where the analogous statements overlap with the literature on Watkins's conjecture.

2. Local invariants

Let q be a prime of bad reduction. We use Tate's algorithm [46], [42, IV.9] for the Kodaira type and c_q, and the following facts.

Lemma 2.1. *Let* q *be a prime of multiplicative reduction,* $n_q=v_q(\Delta_E)$*, and* $\ell\neq q$ *a prime. Then* $\rho_{E,\ell}\bmod\ell^t$ *is unramified at* q *if and only if* $\ell^t\mid n_q$*. This holds for split and for nonsplit reduction.*

Proof. Over the unramified quadratic extension of $\mathbb{Q}_q$ (which does not change the inertia group) E is a Tate curve with parameter q_E, $v_q(q_E)=n_q$. The ℓ^t-torsion is generated by μ_{ℓ^t} and an ℓ^t-th root of q_E. By Kummer theory over $\mathbb{Q}_q^{\mathrm{ur}}$, which contains μ_{ℓ^t} as $q\neq\ell$, inertia acts trivially on $E[\ell^t]$ if and only if q_E is an ℓ^t-th power in $\mathbb{Q}_q^{\mathrm{ur}}$; the units of its valuation ring are ℓ^t-th powers by Hensel's lemma, so this holds if and only if $\ell^t\mid n_q$. □

At an additive prime $q\neq\ell$ the inertia invariants of the torsion are those of the component group: $E[\ell^k]^{I_q}\simeq\Phi_q^{\mathrm{geom}}[\ell^k]$ for every k. Indeed $E[\ell^k]^{I_q}=E(\mathbb{Q}_q^{\mathrm{ur}})[\ell^k]$; the group $E(\mathbb{Q}_q^{\mathrm{ur}})$ is an extension of Φ_q^{geom} by the group $E_0(\mathbb{Q}_q^{\mathrm{ur}})$ of points reducing to the identity component [42, IV.9], and $E_0(\mathbb{Q}_q^{\mathrm{ur}})$ is an extension of the additive group of $\overline{\mathbb{F}}_q$ by a pro-q group, so it is uniquely ℓ-divisible; the snake lemma for multiplication by ℓ^k gives the isomorphism [39, §1].

Lemma 2.2. *Let* $q\neq 3$ *be a prime at which* E *has reduction of type* IV *or* IV**. Then the reduction at* q *is potentially good, the image of the inertia group* I_q *in* $\mathrm{Aut}(T_\ell E)$ *is cyclic of order* 3 *for every prime* $\ell\neq q$*, and its image in* $\mathrm{Aut}(E[3])$ *is a nontrivial unipotent subgroup. Consequently* $E[3]^{I_q}$ *is a line,* $\bar\rho_{E,3}$ *is tamely ramified at* q*, the Serre conductor of* $\bar\rho_{E,3}$ *has exponent* 1 *at* q *and the conductor of* E *has exponent* 2 *at* q*. Moreover, for* $\ell\neq q$*,* $(\mathrm{ad}^0\,V_\ell E)^{I_q}$ *is a line on which* Frob_q *acts by* $\chi_{-3}(q)$*, so that* $P_q(X)=1-\chi_{-3}(q)qX$ *and* $e_\ell(E,q)=v_\ell(q-\chi_{-3}(q))$*; in particular* $P_2(X)=1+2X$ *and* $e_3(E,2)=1$ *when* $q=2$*.*

Proof. By the remark above, $E[3]^{I_q}\simeq\Phi_q^{\mathrm{geom}}[3]\simeq\mathbb{Z}/3$ is a line. Inertia acts nontrivially on $E[3]$, since otherwise $\Phi_q^{\mathrm{geom}}[3]$ would have order 9. If the reduction were potentially multiplicative, E would be over $\mathbb{Q}_q$ the twist of a Tate curve by a quadratic character χ, ramified since the

reduction is additive [42, V.5]; then $\rho_{E,3}(\sigma) = \chi(\sigma)\left(\begin{smallmatrix}1 & *\\ 0 & 1\end{smallmatrix}\right)$ on I_q in a suitable basis, the corner being n times the 3-adic tame character; the subgroup of I_q on which the corner is 0 modulo 3 has index dividing 3, so it is not contained in the kernel of χ, and $\bar\rho_{E,3}(I_q)$ would contain -1, which fixes no line. So the reduction is potentially good, and $\rho_{E,3}(I_q)$ is finite [39, §2]. It lies in $\mathrm{SL}_2(\mathbb{Z}_3)$, since χ_3^{cyc} is unramified at $q \neq 3$, and reduction modulo 3 is injective on it, the kernel of $\mathrm{GL}_2(\mathbb{Z}_3) \to \mathrm{GL}_2(\mathbb{F}_3)$ being torsion-free. The elements of $\mathrm{SL}_2(\mathbb{F}_3)$ fixing a nonzero vector form a group of order 3 of unipotent elements; so $\bar\rho_{E,3}(I_q)$ is that group and $\rho_{E,3}(I_q)$ is cyclic of order 3. The kernel of $\rho_{E,\ell}|_{I_q}$ does not depend on $\ell \neq q$ [39, §2], so the same holds on $T_\ell E$. The order 3 is prime to q, so the image is a quotient of the tame inertia group and the ramification is tame. The Serre conductor exponent of $\bar\rho_{E,3}$ at q is $\dim E[3] - \dim E[3]^{I_q} = 1$, and the conductor exponent of E is $\dim V - \dim V^{I_q} = 2$, there being no wild part. Let σ generate $\rho(I_q)$; its eigenvalues are the two primitive cube roots of unity, so $\sigma^2+\sigma+1 = 0$, and the trace-zero endomorphisms commuting with σ form the line spanned by $\sigma - \sigma^{-1} = 2\sigma + 1$. The arithmetic Frobenius acts on the tame quotient of I_q by $x \mapsto x^q$, so $\rho(\mathrm{Frob}_q)\sigma\rho(\mathrm{Frob}_q)^{-1} = \sigma^q = \sigma^{\chi_{-3}(q)}$, and Frob_q, and therefore also Frob_q^{-1}, acts on the line by $\chi_{-3}(q)$. Hence $L_q(A_f, s)^{-1} = 1 - \chi_{-3}(q)q^{-s}$, $P_q(X) = 1 - \chi_{-3}(q)qX$, and, the naive factor being 1, $e_\ell(E, q) = v_\ell(1 - \chi_{-3}(q)q^{-1}) = v_\ell(q - \chi_{-3}(q))$. For $q = 2$: $\chi_{-3}(2) = -1$ and $e_3(E, 2) = v_3(3/2) = 1$. □

At every prime $q \neq 3$ at which $\rho_{E,3}(I_q)$ is cyclic of order 3, $q = 2$ included, a generator σ satisfies $\sigma^2 + \sigma + 1 = 0$ on T_3E, so T_3E is a torsion-free module of $\mathbb{Z}_3$-rank 2 over the discrete valuation ring $\mathbb{Z}_3[X]/(X^2 + X + 1) \simeq \mathbb{Z}_3[\zeta_3]$, with X acting as σ, hence free of rank one over it; Lemma 2.9 uses this, also at $q = 2$. For $q \geq 5$ the order 3 is also the semistability defect $12/\gcd(v_q(\Delta_E), 12)$ of Kraus [23]; at $q = 2$ the conductor exponent 2 can also be read off [13]. At $q = 3$ the types IV, IV* are wildly ramified, and the lemma does not apply.

Remark 2.3. For the other additive types at $q \geq 5$ the image of inertia mod 3 is not unipotent: it is $\{\pm 1\}$ (type I_0^*), cyclic of order 4 (types III, III*), minus a unipotent, of order 6 (types II, II*), or a ramified quadratic character times a unipotent (type I_n^*). For all these types $E[3]^{I_q} = 0$ and the Serre conductor of $\bar\rho_{E,3}$ has exponent 2 at q. The twist by χ_{q^*} lowers that exponent for I_0^*, I_n^*, II, II*; it lowers the conductor of E only for I_0^* and I_n^*.

Lemma 2.4. *Let q be an odd prime, $\chi = \chi_{q^*}$ the quadratic character of conductor q and $E^{(q)}$ the twist of E by χ, that is, by $\mathbb{Q}(\sqrt{q^*})$.*

(i) *At every prime $p \neq q$ the curves E and $E^{(q)}$ have the same Kodaira type, the same geometric component group and the same conductor exponent.*

(ii) *If E has type I_n at q ($n \geq 0$) then $E^{(q)}$ has type I_n^* at q, and conversely, with the same n. If E has type* II, III, IV *at q then $E^{(q)}$ has type* IV*, III*, II* *respectively, and conversely. For the exchanges $\mathrm{I}_n \leftrightarrow \mathrm{I}_n^*$ ($n \geq 0$) at every odd q, and for all the exchanges at $q \geq 5$, the conductor exponent of $E^{(q)}$ at q is 2 when its reduction is additive and 1 when it is of type I_n, $n \geq 1$. At $q = 3$ the exchanges* II $\leftrightarrow$ IV*, III $\leftrightarrow$ III*, IV $\leftrightarrow$ II* *leave the conductor exponent unchanged; it can exceed 2.*

(iii) *(Atkin-Li [4]) Let g be a newform of weight 2 on $\Gamma_0(M)$ with $q^2 \nmid M$. Then the twist $g \otimes \chi$, with coefficients $a_n(g \otimes \chi) = \chi(n)a_n(g)$, is a newform of weight 2 on $\Gamma_0(M')$, $M' = \mathrm{lcm}(M, q^2)$; that is, $M' = Mq^2$ if $q \nmid M$ and $M' = Mq$ if $q \parallel M$.*

Proof. (i) The character χ is unramified at p, so E and $E^{(q)}$ become isomorphic over an unramified extension of $\mathbb{Q}_p$, and the Kodaira type, the geometric component group and the conductor exponent are invariant under unramified base change. (ii) If $(a_1, a_2, a_3, a_4, a_6)$ is a minimal model of E at q with invariants c_4, c_6, Δ, the twist has a model with invariants $(q^*)^2c_4$, $(q^*)^3c_6$, $(q^*)^6\Delta$ and the same j-invariant; only their q-adic valuations are used below. For potentially multiplicative reduction, $v_q(j) < 0$, the type is I_n or I_n^* with $n = -v_q(j)$ according as the reduction is

multiplicative or additive, and a ramified quadratic twist exchanges the two. For $n = 0$: a model $y^2 = x^3+a_2x^2+a_4x+a_6$ with good reduction at q has twist $y^2 = x^3+q^*a_2x^2+(q^*)^2a_4x+(q^*)^3a_6$, for which the cubic of Tate's algorithm, after removing the powers of q, is $T^3+\epsilon a_2T^2+a_4T+\epsilon a_6$ with $\epsilon = q^*/q = \pm1$; its roots are ϵ times those of $x^3 + a_2x^2 + a_4x + a_6$, so they are distinct modulo q, the criterion for type I_0^*; conversely a curve of type I_0^* at an odd q has a model of the latter shape, and its twist has good reduction. For $q \geq 5$ and potentially good reduction the type is determined by $v_q(\Delta)$ of a minimal model, which lies in $\{0, 2, 3, 4, 6, 8, 9, 10\}$ for the types $\mathrm{I}_0, \mathrm{II}, \mathrm{III}, \mathrm{IV}, \mathrm{I}_0^*, \mathrm{IV}^*, \mathrm{III}^*, \mathrm{II}^*$; the twisted model has $v_q(\Delta)$ increased by 6, and is minimal exactly when the result is less than 12, otherwise 12 is subtracted. For the exchanges $\mathrm{I}_n \leftrightarrow \mathrm{I}_n^*$ at every odd q and for all exchanges at $q \geq 5$ the ramification is tame: $V_\ell E^{(q)} = V_\ell E \otimes \chi$ with χ tame and $V_\ell E$ multiplicative, good, or (for $q \geq 5$) tame; so the conductor exponents are the tame ones, $\dim V - \dim V^{I_q}$. At $q = 3$, for the types $\mathrm{II}, \mathrm{III}, \mathrm{IV}$ and their duals, the ramification can be wild, and we use the formula of Ogg and Saito [26, 37], [42, IV.11]: $v_3(\Delta_{\min}) = f_3 + m - 1$, where f_3 is the conductor exponent and m the number of irreducible components of the special fibre of the minimal regular model, which is $1, 2, 3, 5, 7, 8, 9$ for the types $\mathrm{II}, \mathrm{III}, \mathrm{IV}, \mathrm{I}_0^*, \mathrm{IV}^*, \mathrm{III}^*, \mathrm{II}^*$. The reduction of $E^{(3)}$ is potentially good, as j is unchanged, and not of type I_0 or I_0^*, since by the case $n = 0$ above that would make E of type I_0^* or I_0. The character χ_{-3} is tame, so it does not change the Swan conductor; and neither $V_\ell E$ nor $V_\ell E^{(3)}$ has nonzero inertia invariants, since the image of I_3 on each is not contained in $\{\pm1\}$ (otherwise a quadratic twist would have good reduction and the type would be I_0 or I_0^*), so it contains an element of order 3, 4 or 6, which has no eigenvalue ±1. Hence $f_3(E^{(3)}) = f_3(E)$. The twisted model has discriminant $3^6\Delta$, so $v_3(\Delta_{\min})$ changes by 6 modulo 12; with f_3 fixed, Ogg's formula gives $m(E^{(3)})-1 \equiv m(E)-1+6 \pmod{12}$, and as $0 \leq m-1 \leq 8$ on both sides, $m \mapsto m \pm 6$: $1 \leftrightarrow 7$, $2 \leftrightarrow 8$, $3 \leftrightarrow 9$, which are the three exchanges (the types with $m = 7, 8, 9$ and potentially good reduction being IV^*, III^*, II^*, as I_n^* with $n \geq 1$ is potentially multiplicative at odd q). For the classification at 3 compare [22, 27]. (iii) is [4, Theorem 3.1 and p. 228]; in the case $q \parallel M$, the local component of g at q is an unramified twist of the Steinberg representation, whose twist by a ramified quadratic character has conductor exponent 2. □

Lemma 2.5 (the omitted factors at tame primes)**.** *Let $q \geq 5$ be a bad prime, $q \neq \ell$. Then $L_q(A_f, s)^{-1}$ and $e_\ell(E, q)$ are as in the table of Section 1.1. The rows I_n^* and I_0^* hold for every odd prime $q \neq \ell$, the row I_n for every $q \neq \ell$.*

Proof. The inertia group acts on V through a finite cyclic quotient when the reduction is potentially good, and through a quadratic character times a unipotent when it is potentially multiplicative; $L_q(A_f, s)^{-1} = \prod(1 - \lambda q^{-s})$ over the eigenvalues λ of the geometric Frobenius Frob_q^{-1} on $(\mathrm{ad}^0 V)^{I_q}$, in the normalisation in which a good prime has the eigenvalues $\alpha/\beta, 1, \beta/\alpha$ on $\mathrm{ad}^0 V$.

Type I_n. The local component of f is an unramified twist of the Steinberg representation; $(\mathrm{ad}^0 V)^{I_q}$ is a line on which the geometric Frobenius has eigenvalue q^{-1}, so $L_q(A_f, s)^{-1} = 1 - q^{-1-s}$, which is also the naive factor since $a_q^2 = 1$: $e_\ell = 0$. This holds at every $q \neq \ell$. *Type* I_n^**, q odd.* $V = V' \otimes \chi_{q^*}$ with V' Steinberg at q (Lemma 2.4), $\mathrm{ad}^0 V = \mathrm{ad}^0 V'$ and $a_q = 0$, so the naive factor is 1 and $e_\ell = v_\ell(1 - q^{-2}) = v_\ell(q^2 - 1)$. *Types* $\mathrm{II}, \mathrm{II}^*, \mathrm{IV}, \mathrm{IV}^*$. The image of I_q in $\mathrm{Aut}\, V$ is cyclic of order 3 or 6 [23], generated by σ or $-\sigma$ with σ of order 3 and eigenvalues $\zeta_3^{\pm1}$. On $\mathrm{ad}^0 V$ the sign disappears and the invariants form the line spanned by $2\sigma + 1$, whose square is -3, inside $\mathrm{End}_{I_q}(T) = \mathbb{Z}_\ell[\sigma] \simeq \mathbb{Z}_\ell \otimes \mathbb{Z}[\zeta_3]$ (not a field when $\ell \equiv 1 \pmod 3$); Frobenius centralises σ if $q \equiv 1 \pmod 3$ and inverts it if $q \equiv 2$, for either choice of Frobenius, so it acts on the line by $\chi_{-3}(q)$, and $e_\ell = v_\ell(1 - \chi_{-3}(q)q^{-1}) = v_\ell(q - \chi_{-3}(q))$. *Types* $\mathrm{III}, \mathrm{III}^*$. The image has order 4, generated by τ with eigenvalues $\pm i$; the invariant line of $\mathrm{ad}^0 V$ is spanned by τ itself,

which has trace 0 (it is diagonalisable only over $\mathbb{Q}_\ell(i)$), and Frobenius acts on it by $\chi_{-4}(q)$, so $e_\ell = v_\ell(q-\chi_{-4}(q))$. *Type* I_0^**, q odd.* $V = V' \otimes \chi_{q^*}$ with V' unramified with arithmetic Frobenius eigenvalues α, β; $\mathrm{ad}^0 V$ is unramified with eigenvalues $\alpha/\beta, 1, \beta/\alpha$ for either Frobenius, and

$$L_q(A_f,1)^{-1} = \Big(1-\frac{\alpha}{\beta q}\Big)\Big(1-\frac{1}{q}\Big)\Big(1-\frac{\beta}{\alpha q}\Big) = \frac{q-1}{q}\cdot\frac{(q+1)^2-a_q(E')^2}{q^2},$$

using $\alpha\beta = q$ and $\alpha+\beta = a_q(E')$. Only Lemma 2.4 for the exchanges $\mathrm{I}_n \leftrightarrow \mathrm{I}_n^*$ and $\mathrm{I}_0 \leftrightarrow \mathrm{I}_0^*$ was used in the last two rows, and it holds at every odd q; at $q=3$ the formulas read $L_3(A_f,1)^{-1} = \frac{8}{9}$ for I_n^* and $\frac{2}{3}\cdot\frac{16-a_3(E')^2}{9}$ for I_0^*. In each case the values were confirmed against PARI's `lfunsympow` on every curve of the tables. □

The shape of the factor at every bad prime, 2 and 3 included, is determined by the image of inertia.

Proposition 2.6. *Let q be a bad prime and $\ell \neq q$ an odd prime. Then $P_q(X)$ has one of the following four shapes.*

(i) $P_q(X) = 1 - X$*, if $v_q(j_E) < 0$; then $e_\ell(E,q) = 0$ if the reduction is multiplicative and $e_\ell(E,q) = v_\ell(q^2-1)$ if it is additive.*

(ii) $P_q(X) = (1-qX)(1-(a^2-2q)X+q^2X^2)$*, if the reduction is potentially good and $\rho(I_q) = \{\pm 1\}$; here $a = a_q(E')$ for a quadratic twist E' of E with good reduction at q, and $e_\ell(E,q) = v_\ell(q-1) + v_\ell((q+1)^2-a^2)$.*

(iii) $P_q(X) = 1 - cqX$ *with $c = \pm 1$, if the reduction is potentially good and $\rho(I_q)$ is cyclic of order 3, 4 or 6; here c is the eigenvalue of Frobenius on the line $(\mathrm{ad}^0 V)^{I_q}$, and $e_\ell(E,q) = v_\ell(q-c)$.*

(iv) $P_q(X) = 1$*, if the reduction is potentially good and $\rho(I_q)$ is not abelian, which happens only for $q = 2, 3$; then $e_\ell(E,q) = 0$.*

Proof. (i) Over $\mathbb{Q}_q$, $\rho \simeq \rho' \otimes \chi$ with ρ' the representation of a Tate curve, possibly twisted by an unramified character, and χ quadratic [42, V.5]; so $\mathrm{ad}^0 V \simeq \mathrm{ad}^0 V'$, inertia acts on V' through a nontrivial unipotent, and as in the proof of Lemma 2.8 the invariants of $\mathrm{ad}^0 V$ are the line spanned by e, on which the geometric Frobenius acts by q^{-1}. So $L_q(A_f,s)^{-1} = 1-q^{-1-s}$ and $P_q(X) = L_q(A_f, s-1)^{-1} = 1-X$ with $X = q^{-s}$; the naive factor is the true one if $\delta_q = 1$ and 1 if $\delta_q = 0$. In the other cases $\rho(I_q)$ is a finite subgroup of $\mathrm{SL}_2(\mathbb{Z}_\ell)$ whose elements have characteristic polynomials in $\mathbb{Z}[X]$ [39, §2], so their orders are $1, 2, 3, 4, 6$; the reduction is additive, $\delta_q = 0$, and the naive factor is 1. (ii) The quadratic character $\psi = \rho|_{I_q}$ of I_q is invariant under conjugation by $G_{\mathbb{Q}_q}$, and it extends to a character χ of $G_{\mathbb{Q}_q} = I_q \rtimes \langle\phi\rangle$, ϕ a Frobenius lift generating a complement isomorphic to $\widehat{\mathbb{Z}}$, by $\chi(i\phi^k) = \psi(i)$. Every quadratic character of $G_{\mathbb{Q}_q}$ is the restriction of a quadratic Dirichlet character χ' (for q odd a product of χ_{q^*} and a character unramified at q; for $q=2$ a product of χ_{-4}, χ_8 and χ_5), and $E' = E \otimes \chi'$ has good reduction at q. Then $\mathrm{ad}^0 V \simeq \mathrm{ad}^0 V_\ell E'$ is unramified with eigenvalues $\alpha/\beta, 1, \beta/\alpha$, $\alpha+\beta = a$, $\alpha\beta = q$, so $P_q(X) = (1-\alpha^2X)(1-qX)(1-\beta^2X)$, which is the stated polynomial; a is determined up to sign, and a^2 is well defined. The value of e_ℓ is the computation of Lemma 2.5. (iii) A finite abelian subgroup of SL_2 in characteristic 0 is cyclic; let g generate $\rho(I_q)$, of order $m \in \{3,4,6\}$, with eigenvalues $\zeta^{\pm 1}$, $\zeta \neq \zeta^{-1}$. The trace-zero endomorphisms commuting with g form the line spanned by $g - g^{-1}$. Frobenius normalises $\rho(I_q)$, so it conjugates g to some g^k with the same eigenvalues, that is, to $g^{\pm 1}$, and it acts on the line by $c = \pm 1$, the same sign for the arithmetic and the geometric Frobenius. So $L_q(A_f,s)^{-1} = 1-cq^{-s}$, $P_q(X) = 1-cqX$ and $e_\ell = v_\ell(1-c/q) = v_\ell(q-c)$. (iv) A faithful two-dimensional representation of a non-abelian finite group in characteristic 0 is absolutely irreducible, so the commutant of $\rho(I_q)$ consists of scalars and $(\mathrm{ad}^0 V)^{I_q} = 0$. For $q \geq 5$ the image is tame, hence cyclic. □

Remark 2.7. At $q = 2, 3$ the Kodaira type does not determine P_q, but PARI does, and Definition 1.1 applies verbatim. For instance a type IV or IV* at 2 always has $P_2(X) = 1 + 2X$ and $e_3 = 1$ (Lemma 2.2); a type II or II* at 2 usually has $P_2 = 1$, but 336f1 (type II at 2) has $P_2 = 1 + 2X$, $e_3(E, 2) = 1$ and $\deg \varphi_E = 24$, and 1232l1 (type II* at 2), the twist of 77a1 by χ_{-4} with $a_2(77\text{a}1) = 0$, has $P_2 = (1 - 2X)(1 + 2X)^2 = 1 + 2X - 4X^2 - 8X^3$, $e_3(E, 2) = 2$ and $\deg \varphi_E = 288$, so that the bound of Theorem 1.2 at $\ell = 3$ is attained by the prime 2 alone. A type I_n^* at 2 has potentially good reduction in a little over half of the cases (below conductor 10,000: 58%, 2,340 of 4,037, among the optimal curves, and 61%, 5,935 of 9,667, among all curves), usually with $P_2 = 1$, and then $t_\ell = 0$. Among all curves of conductor below 10,000, every cubic factor at 2 (1,143) and at 3 (2,283) equals the formula of (ii) for a twist of good reduction found independently (534 and 1,308 among the optimal non-CM curves); that a cubic factor has shape (ii) is automatic, (ii) being the only cubic shape. The prime 2 can contribute at $\ell \geq 5$: 592c1, the twist of 37a1 by χ_{-4} with $a_2(37\text{a}1) = -2$, has type II* at 2, $P_2 = (1 - 2X)(1 + 4X^2)$, $e_5(E, 2) = 1$ and $\deg \varphi_E = 80 = 2^4 \cdot 5$. At $q = \ell$ the Euler factor is not the relevant quantity, and the weights w_3 of Conjecture 1.3 are empirical.

Lemma 2.8 (the Tamagawa factor at a multiplicative prime)**.** *Let q be a prime of multiplicative reduction, $n = n_q$, ℓ an odd prime different from q, $T = T_\ell E$ and $M = \mathrm{ad}^0 T$. Let $q_E \in \mathbb{Q}_q^\times$ be the Tate parameter, $u = q_E q^{-n} \in \mathbb{Z}_q^\times$ its unit part, and $c(u)$ the class of the reduction of u in $\mathbb{F}_q^\times/(\mathbb{F}_q^\times)^{\ell^v}$, $v = v_\ell(q-1)$, identified with an element of $\mathbb{Z}/\ell^v$ by the choice of a generator; $v_\ell(c(u))$ does not depend on that choice, $v_\ell(0)$ is read as v, and $v_\ell(c(u)) \geq k$ if and only if the reduction of u is an ℓ^k-th power in $\mathbb{F}_q^\times$, for $k \leq v$. Then $H^1(I_q, M)_{\mathrm{tor}} \simeq (\mathbb{Z}/\ell^{v_\ell(n)})^2$ and*

$$v_\ell\big(\mathrm{Tam}_q(\mathrm{ad}^0 T)\big) = v_\ell(n) + \min\big(v_\ell(n), v_\ell(q-1), v_\ell(c(u))\big).$$

In particular $v_\ell(\mathrm{Tam}_q(\mathrm{ad}^0 T)) \geq t_\ell(E, q)$, with equality unless $q \equiv 1 \pmod{\ell}$, $\ell \mid n$ and the reduction of u is an ℓ-th power in $\mathbb{F}_q^\times$. The same formula holds at a prime $q \neq \ell$ of additive potentially multiplicative reduction, $q = 2$ included, with $n = -v_q(j_E)$ and q_E the Tate parameter, determined by $j(q_E) = j_E$.

Proof. Over the unramified quadratic extension of $\mathbb{Q}_q$ the curve is a Tate curve, so T has a basis e_1, e_2 in which $G = G_{\mathbb{Q}_q}$ acts through matrices $\left(\begin{smallmatrix} \chi_\ell \varepsilon & \varepsilon\kappa \\ 0 & \varepsilon \end{smallmatrix}\right)$, with ε unramified of order at most 2 and κ the Kummer cocycle of q_E in $H^1(G, \mathbb{Z}_\ell(1)) = \varprojlim \mathbb{Q}_q^\times/(\mathbb{Q}_q^\times)^{\ell^k}$; the nonsplit curve is the Tate curve twisted by ε, and $\varepsilon\kappa$, not κ, satisfies the cocycle relation for the twisted matrices. The inertia group $I = I_q$ acts through its maximal pro-ℓ tame quotient, generated by an element t with $te_1 = e_1$, $te_2 = e_2 + n'e_1$, $v_\ell(n') = v_\ell(n)$, since the restriction of κ to I is n times the tame character and ε is trivial on I; the kernel of $I \to \langle t \rangle$ has trivial pro-ℓ quotient, so by inflation-restriction $H^1(I, M') = H^1(\langle t \rangle, M') = M'/(t-1)M'$ for every $\mathbb{Z}_\ell$-module M' of finite type with this action. In the basis $e = e_1 \otimes e_2^\vee$, $h = e_1 \otimes e_1^\vee - e_2 \otimes e_2^\vee$, $f = e_2 \otimes e_1^\vee$ of $M = \mathrm{ad}^0 T$ one has $(t-1)e = 0$, $(t-1)h = -2n'e$, $(t-1)f = n'h - n'^2 e$, and as 2 is a unit, $(t-1)M = n'(\mathbb{Z}_\ell e \oplus \mathbb{Z}_\ell h)$; so $H^1(I, M) = (\mathbb{Z}_\ell/n')\bar{e} \oplus (\mathbb{Z}_\ell/n')\bar{h} \oplus \mathbb{Z}_\ell \bar{f}$ and its torsion is $(\mathbb{Z}/\ell^{v_\ell(n)})^2$. The group G/I acts on $H^1(I, M)$ by $(\sigma \cdot z)(i) = \sigma z(\sigma^{-1} i \sigma)$ for a cocycle z; for the arithmetic Frobenius Frob, $\mathrm{Frob}^{-1}\, t\, \mathrm{Frob} = t^{q^{-1}}$ and $z(t^k) \equiv k\, z(t)$ modulo $(t-1)M$, so Frob acts on $M/(t-1)M$ as $q^{-1}\,\mathrm{Ad}(\rho(\mathrm{Frob}))$, that is, by conjugation by $\rho(\mathrm{Frob})$ followed by multiplication by q^{-1}. Writing $\rho(\mathrm{Frob}) = \left(\begin{smallmatrix} \varepsilon q & c' \\ 0 & \varepsilon \end{smallmatrix}\right)$ with $\varepsilon = \varepsilon(\mathrm{Frob})$, the adjoint action sends e to qe and h to $h - 2c'\varepsilon e$, so on the torsion $\bar{e} \mapsto \bar{e}$ and $\bar{h} \mapsto q^{-1}(\bar{h} - 2c'\varepsilon\bar{e})$. The fixed points are the $x\bar{e} + y\bar{h}$ with $(q-1)y \equiv 0$ and $c'y \equiv 0 \pmod{\ell^{v_\ell(n)}}$: $\ell^{v_\ell(n)}$ choices of x and $\ell^{\min(v_\ell(n), v_\ell(q-1), v_\ell(c'))}$ of y. Finally, q being a uniformiser of $\mathbb{Q}_q$, $H^1(\mathbb{Q}_q, \mathbb{Z}_\ell(1))$ splits as $\mathbb{Z}_\ell\kappa_q \oplus H^1_{\mathrm{ur}}$, and $\kappa = n\kappa_q + \kappa_u$, where κ_u is unramified, inflated from the class of u in $H^1(G/I, \mathbb{Z}_\ell(1)) = \mathbb{Z}_\ell(1)/(q-1) \simeq \mathbb{F}_q^\times \otimes \mathbb{Z}_\ell$. The value $c' = \varepsilon\kappa(\mathrm{Frob})$ is a cocycle value: it is ε times the class of u for the Frobenius lift fixing a compatible system

of ℓ-power roots of q, and in general it is well defined only modulo $\ell^{\min(v_\ell(n),v_\ell(q-1))}$ (changing the lift changes it by a multiple of n', changing the cocycle by a multiple of $q-1$). Neither indeterminacy affects $\min(v_\ell(n), v_\ell(q-1), v_\ell(c'))$, and $v_\ell(c')$ may be replaced by $v_\ell(c(u))$ there. The identification of Tam_q with the Fitting ideal of the G-invariants of the torsion of $H^1(I,M)$ is [17, I.4.2.2], as recalled in [12, p. 708]. As a check, the same computation for $M=T$ gives $(\mathbb{Z}/\ell^{v_\ell(n)})\bar{e}_1$ with Frob acting by ε, that is, the ℓ-part of the Tamagawa number $c_q(E)$, as it should. At an additive potentially multiplicative prime, E is over $\mathbb{Q}_q$ the twist of the Tate curve E_{q_E} by a ramified quadratic character [42, V.5]; the $G_{\mathbb{Q}_q}$-module $\mathrm{ad}^0 T$ is unchanged by the twist, and the computation above, which only uses that module, applies with $n = v_q(q_E) = -v_q(j_E)$. At $q=2$, $v_\ell(q-1)=0$ and the minimum vanishes. □

Lemma 2.9 (the Tamagawa factor at a potentially good additive prime)**.** *Let ℓ be odd and let $q \neq \ell$ be a prime at which E has additive, potentially good reduction. Then $H^1(I_q, \mathrm{ad}^0 T)_{\mathrm{tor}} = 0$; in particular $v_\ell(\mathrm{Tam}_q(\mathrm{ad}^0 T)) = 0$.*

Proof. Let $M = \mathrm{ad}^0 T$ and let K be the kernel of the action of $I = I_q$ on M; the quotient $\Delta = I/K$ is the image of the finite group $\rho(I)$ modulo $\rho(I) \cap \{\pm 1\}$. Inflation-restriction gives the exact sequence $0 \to H^1(\Delta, M) \to H^1(I,M) \to \mathrm{Hom}(K,M)^\Delta$. The group K is the inertia group of a finite extension of $\mathbb{Q}_q^{\mathrm{ur}}$, and a continuous homomorphism from K to the pro-ℓ group M factors through its maximal pro-ℓ abelian quotient, which is $\mathbb{Z}_\ell(1)$ as $q \neq \ell$; so $\mathrm{Hom}(K,M) \simeq M(-1)$ is torsion-free, and the torsion of $H^1(I,M)$ lies in $H^1(\Delta,M)$, which is killed by $\#\Delta$. The elements of $\rho(I)$ have orders $1,2,3,4,6$ (proof of Proposition 2.6), so $\#\Delta$ is divisible only by 2 and 3, and $H^1(\Delta,M)=0$ for $\ell \geq 5$. Let $\ell = 3$. The 3-primary part of $H^1(\Delta,M)$ injects into $H^1(P,M)$ for a Sylow 3-subgroup P of Δ, by restriction and corestriction. A finite 3-subgroup of SL_2 in characteristic 0 is cyclic, and it has exponent 3, so P is trivial or generated by the image of an element $\sigma \in \rho(I)$ of order 3. Then T is free of rank one over $\mathbb{Z}_3[X]/(X^2+X+1)$, with X acting as σ, so in a suitable basis $\sigma = \left(\begin{smallmatrix} 0 & -1 \\ 1 & -1 \end{smallmatrix}\right)$; with $e = \left(\begin{smallmatrix} 0 & 1 \\ 0 & 0 \end{smallmatrix}\right)$, the elements e, $\sigma e \sigma^{-1}$, $\sigma^2 e \sigma^{-2}$ have coordinates $(1,0,0)$, $(0,0,-1)$, $(1,-1,-1)$ in the basis e,h,f of $\mathrm{ad}^0 T$, a matrix of determinant -1. So $\mathrm{ad}^0 T \simeq \mathbb{Z}_3[P]$ is free over $\mathbb{Z}_3[P]$ and $H^1(P, \mathrm{ad}^0 T) = 0$. Hence $H^1(I,M)_{\mathrm{tor}} = 0$, and so is its group of $G_{\mathbb{Q}_q}$-invariants. □

By contrast $\mathrm{Tam}_q(T)$ can be divisible by 3 at a prime of type IV or IV*: then $H^1(\langle\sigma\rangle, T) = T/(1-\zeta_3)T \simeq \mathbb{F}_3$.

3. Proof of Theorem 1.5

Throughout this section ℓ is odd and $\ell \nmid N$. Let $f = f_E \in S_2(\Gamma_0(N))$ be the newform of the isogeny class of E, let E_0 be the optimal curve in the class, and let $r = r_{E_0}$ be its congruence number: the largest integer r such that $f \equiv g \pmod r$ coefficientwise for some $g \in S_2(\Gamma_0(N))$ with integer coefficients orthogonal to f [2, §1]. By [2, Theorem 2.2], $\deg \varphi_{E_0}$ divides r and $v_p(\deg \varphi_{E_0}) = v_p(r)$ for every prime p with $p^2 \nmid N$; as $\ell \nmid N$ it suffices to prove $\ell \mid r$. Let $\mathbb{T}$ be the Hecke algebra of level N, the subring of $\mathrm{End}\, S_2(\Gamma_0(N))$ generated over $\mathbb{Z}$ by all T_n, $n \geq 1$, including the operators U_p for $p \mid N$, and write $\bar{\rho} = \bar{\rho}_{E,\ell}$, irreducible by hypothesis, hence absolutely irreducible, complex conjugation having the distinct eigenvalues 1 and -1. The two lemmas below reduce the theorem to the production of a congruent newform of lower level.

Lemma 3.1 (congruence criterion)**.** *Let $\phi_f \colon \mathbb{T} \to \mathbb{Z}$, $T_n \mapsto a_n(f)$, let $I = \ker \phi_f$ and let J be the annihilator in $\mathbb{T}$ of the orthogonal complement $f^\perp$ of f. Then $\mathbb{T}/(I+J)$ is cyclic of order r; equivalently, r generates the congruence ideal $\phi_f(J) = \phi_f(\mathrm{Ann}_{\mathbb{T}} I)$. Consequently, if there is a normalised eigenform $g \in S_2(\Gamma_0(N)) \otimes \overline{\mathbb{Q}}$ for all of $\mathbb{T}$, orthogonal to f, with coefficients in the*

ring of integers $\mathcal{O}$ of a number field, and a prime $\lambda \mid \ell$ of $\mathcal{O}$ such that $a_n(g) \equiv a_n(f) \pmod{\lambda}$ for every $n \geq 1$, then $\ell \mid r$. The same statements hold with $\mathbb{Z}$ replaced by $\mathbb{Z}_\ell$ and r by its ℓ-part.

Proof. Writing $\mathbb{T} \otimes \mathbb{Q} = \mathbb{Q} \times \mathbb{T}'$ with $\mathbb{T}'$ acting faithfully on $f^\perp$ (the f-line is $\mathbb{T}$-stable, and so is $f^\perp$, because f is an eigenvector of the adjoints of the T_n as well), one has $I = \mathbb{T} \cap (0 \times \mathbb{T}')$ and $J = \mathbb{T} \cap (\mathbb{Q} \times 0)$, so $I \cap J = 0$ and $\mathbb{T}/(I+J) \simeq \mathbb{Z}/\phi_f(J)$ is cyclic, of order r' say. The pairing $S_2(\Gamma_0(N);\mathbb{Z}) \times \mathbb{T} \to \mathbb{Z}$, $(g,t) \mapsto a_1(tg)$, is perfect [34, Theorem 2.2], [1, Lemma 2.1] (see [2, §4.1]), and under it the forms killed by I are $\mathbb{Z}f$ and the forms killed by J are $L = f^\perp \cap S_2(\Gamma_0(N);\mathbb{Z})$. Dualising the inclusion $\mathbb{T} \subset \mathbb{T}/I \times \mathbb{T}/J$, whose cokernel is $\mathbb{T}/(I+J)$, shows that $\mathbb{Z}f \oplus L$ has index r' in $S_2(\Gamma_0(N);\mathbb{Z})$. The quotient $S_2(\Gamma_0(N);\mathbb{Z})/(\mathbb{Z}f \oplus L)$ embeds in $\mathbb{Q}f/\mathbb{Z}f$ by projection onto $\mathbb{Q}f$ along $f^\perp$, so it is cyclic, generated by the class of a form h with projection f/r'; then $r'h = f + g_0$ with $g_0 \in L$, so $f \equiv -g_0 \pmod{r'}$ and $r' \leq r$; conversely a congruence $f \equiv g_1 \pmod{r}$ with $g_1 \in L$ gives the integral form $(f-g_1)/r$, whose class has order r, so $r \mid r'$. Hence $r = r' = \#\mathbb{T}/(I+J)$. Now let $\phi_g \colon \mathbb{T} \to \mathcal{O}$ be the eigencharacter of g. It kills J because $g \in f^\perp$, and $\phi_g \equiv \phi_f \pmod{\lambda}$ on all of $\mathbb{T}$ by hypothesis, so $\phi_g(I) \subset \lambda$. Thus $I+J$ lies in the kernel of $\mathbb{T} \to \mathcal{O}/\lambda$, a maximal ideal of residue characteristic ℓ, and ℓ divides $\#\mathbb{T}/(I+J) = r$. Every step holds verbatim over $\mathbb{Z}_\ell$, the pairing being perfect after tensoring with $\mathbb{Z}_\ell$. □

Lemma 3.2 (oldforms)**.** *Let h be a newform of weight 2 on $\Gamma_0(M)$ with $M \mid N$ and $h \neq f$, with coefficients in $\mathcal{O}$, and let $\lambda \mid \ell$ be a prime of $\mathcal{O}$ such that $a_p(h) \equiv a_p(f) \pmod{\lambda}$ for all primes $p \nmid \ell N$. Assume that $p \nmid M$ for every prime $p \parallel N$ at which $\bar\rho$ is unramified. Then, after enlarging $\mathcal{O}$ and λ, the span of the forms $h(dz)$, $d \mid N/M$, contains an eigenform g as in Lemma 3.1. Consequently $\ell \mid r$.*

Proof. The forms $h(dz)$, $d \mid N/M$, span a $\mathbb{T}$-stable subspace V of $S_2(\Gamma_0(N))$ orthogonal to the newform f (whether $M < N$, or $M = N$ and $h \neq f$). On V the operators T_n with $(n,N) = 1$ act by the scalars $a_n(h)$, and U_p for $p \mid M$ with $v_p(M) = v_p(N)$ acts by $a_p(h)$. For $p \mid N/M$ put $e_p = v_p(N/M)$; on the span of $h(p^j z)$, $0 \leq j \leq e_p$, one has $U_p h(p^j z) = h(p^{j-1}z)$ for $j \geq 1$, and $U_p h = a_p(h)h - p\,h(pz)$ if $p \nmid M$, $U_p h = a_p(h)h$ if $p \mid M$. The eigenvalues of U_p there are therefore the roots α_p, β_p of $X^2 - a_p(h)X + p$, together with 0 if $e_p \geq 2$, when $p \nmid M$; and $a_p(h)$, together with 0 if $e_p \geq 1$, when $p \mid M$. The operator U_p is not semisimple on this span when 0 is a repeated eigenvalue, but each listed eigenvalue has an eigenvector: the p-stabilisations $h - \beta_p h(pz)$, $h - \alpha_p h(pz)$ for α_p, β_p, the form h for $a_p(h)$, and the depleted forms $h - a_p(h)h(pz) + p\,h(p^2 z)$ (if $p \nmid M$), $h - a_p(h)h(pz)$ (if $p \parallel M$) or h itself (if $p^2 \mid M$, where $a_p(h) = 0$) for the eigenvalue 0. Since V is the tensor product of these local spaces and $\mathbb{T}$ is commutative, for every choice of one such eigenvalue μ_p at each $p \mid N/M$ there is a common eigenform $g \in V \otimes \overline{\mathbb{Q}}$ of all of $\mathbb{T}$ with $U_p g = \mu_p g$ for $p \mid N/M$, $U_p g = a_p(h)g$ for the other $p \mid N$, and $T_n g = a_n(h)g$ for $(n,N) = 1$; normalise it by $a_1(g) = 1$. We choose the μ_p and check $a_p(g) \equiv a_p(f) \pmod{\lambda}$ prime by prime; the congruence for all n then follows from the multiplicativity of the coefficients, from $a_{p^k} = a_p^k$ for $p \mid N$, and from the recursion $a_{p^{k+1}} = a_p a_{p^k} - p\,a_{p^{k-1}}$ for $p \nmid N$, which f and g both satisfy.

$p \nmid \ell N$. $a_p(g) = a_p(h) \equiv a_p(f)$ by hypothesis. Consequently $\bar\rho_h \simeq \bar\rho$ over a common finite extension of $\mathbb{F}_\ell$, by Chebotarev and Brauer-Nesbitt, $\bar\rho$ being irreducible, hence absolutely irreducible.

$p = \ell$. Both f and h have weight 2 and level prime to ℓ. By the theorems of Deligne and Fontaine describing $\bar\rho_f|_{D_\ell}$ and $\bar\rho_h|_{D_\ell}$ [15, §2], the residue class of a_ℓ is determined by the restriction to D_ℓ: it is 0 when that restriction is irreducible, and otherwise the eigenvalue of Frobenius on the unramified quotient. That quotient is well defined: the two characters of the semisimplification restrict to I_ℓ as $\bar\chi_\ell$ and 1, and $\bar\chi_\ell|_{I_\ell} \neq 1$ since ℓ is odd, so the unramified quotient is the module of I_ℓ-coinvariants. Hence $a_\ell(h) \equiv a_\ell(f)$.

$p^2 \mid N$. Here E has additive reduction at p and $a_p(f) = 0$. If $v_p(M) = v_p(N)$ then $a_p(h) = 0$, because a newform on $\Gamma_0(M)$ with $p^2 \mid M$ has $a_p = 0$ [3, Theorem 3]. Otherwise $e_p \geq 1$, and $e_p \geq 2$ if $p \nmid M$, so 0 is an available eigenvalue: take $\mu_p = 0$.

$p \,\|\, N$, $\bar{\rho}$ *unramified at* p. By hypothesis $p \nmid M$, so $e_p = 1$ and $\mu_p \in \{\alpha_p, \beta_p\}$. Here $\rho_{E,\ell}|_{D_p}$ is an extension of an unramified character ε by $\varepsilon\chi_\ell$ with $\varepsilon(\mathrm{Frob}_p) = a_p(f) = \pm 1$ [10, Theorem 3.1(e)], so the eigenvalues of $\bar{\rho}(\mathrm{Frob}_p)$ are $a_p(f)p$ and $a_p(f)$. As $\bar{\rho}_h \simeq \bar{\rho}$ is unramified at p and $p \nmid M$, the characteristic polynomial of $\bar{\rho}_h(\mathrm{Frob}_p)$ is $X^2 - a_p(h)X + p \equiv (X - a_p(f)p)(X - a_p(f)) \pmod{\lambda}$, so one of α_p, β_p is congruent to $a_p(f)$ modulo a prime above λ in $\mathcal{O}[\alpha_p]$; take it as μ_p.

$p \,\|\, N$, $\bar{\rho}$ *ramified at* p. Then $p \mid M$, since otherwise $\bar{\rho}_h$ would be unramified at p; so $v_p(M) = 1 = v_p(N)$ and $a_p(g) = a_p(h) = \pm 1$. Both $\bar{\rho}|_{D_p}$ and $\bar{\rho}_h|_{D_p}$ are extensions of an unramified character by its twist by $\bar{\chi}_\ell$, with Frobenius eigenvalue $a_p(f)$, respectively $a_p(h)$, on the unramified quotient [10, Theorem 3.1(e)], and both are non-split because $\bar{\rho}$ is ramified at p. A non-split extension of two characters has a unique stable line, so the isomorphism $\bar{\rho}|_{D_p} \simeq \bar{\rho}_h|_{D_p}$ matches the sublines and hence the quotients, and $a_p(h) \equiv a_p(f)$.

In all cases $a_p(g) \equiv a_p(f) \pmod{\lambda}$, and Lemma 3.1 applies. □

We use the optimal-level theorem of Ribet, Carayol and Diamond [33, 7, 11] in the form of [10, Theorem 3.15]: *let ℓ be odd and let $\bar{\sigma}\colon \mathrm{Gal}(\overline{\mathbb{Q}}/\mathbb{Q}) \to \mathrm{GL}_2(\overline{\mathbb{F}}_\ell)$ be irreducible and modular, absolutely irreducible on $\mathrm{Gal}(\overline{\mathbb{Q}}/\mathbb{Q}(\sqrt{-3}))$ if $\ell = 3$; then $\bar{\sigma}$ arises from a newform of weight 2 and level $N(\bar{\sigma})\ell^{\delta(\bar{\sigma})}$ with nebentypus of order prime to ℓ, where $N(\bar{\sigma})$ is the Serre conductor and $\delta(\bar{\sigma}) = 0$ when $\bar{\sigma}|_{D_\ell}$ is good.* For $\bar{\sigma} = \bar{\rho}_{E',\ell}$ with E' an elliptic curve of conductor prime to ℓ, the restriction to D_ℓ comes from the finite flat group scheme $E'[\ell]$ and is good, and $\det \bar{\sigma} = \bar{\chi}_\ell$ forces the nebentypus, which is trivial modulo λ and of order prime to ℓ, to be trivial. So in that case $\bar{\sigma}$ arises from a newform of weight 2 on $\Gamma_0(N(\bar{\sigma}))$. Recall also that $N(\bar{\rho}_{E',\ell})$ divides the conductor of E' [7], and that $N(\bar{\sigma})$ is prime to every prime at which $\bar{\sigma}$ is unramified.

Case (a). Here $q^2 \,\|\, N$ and $\ell = 3$ (Lemma 2.2, which gives conductor exponent 2 at q, also for $q = 2$). By Lemma 2.2 the Serre conductor $M = N(\bar{\rho})$ has exponent exactly 1 at q, so $M \mid N/q$. The optimal-level theorem gives a newform g of weight 2 on $\Gamma_0(M)$ with $\bar{\rho}_g \simeq \bar{\rho}$, that is, $a_p(g) \equiv a_p(f) \pmod{\lambda}$ for all $p \nmid 3N$. Take $h = g$: it has level $M < N$, and $p \nmid M$ for every prime p at which $\bar{\rho}$ is unramified. Lemma 3.2 gives $3 \mid r$; at $p = q$ it uses the eigenvalue 0 of U_q, available since $v_q(M) = 1 < 2 = v_q(N)$.

Case (b). Here $q^2 \,\|\, N$, $\ell = 3$, and E has type II or II* at $q \geq 5$. Let $\chi = \chi_{q^*}$ and $E' = E^{(q)}$. By Lemma 2.4, E' has type IV* or IV at q, the same types as E elsewhere, and conductor N; its newform is $f' = f \otimes \chi$, a newform of weight 2 on $\Gamma_0(N)$ (both have $a_q = 0$), and $\bar{\rho}_{E',3} = \bar{\rho} \otimes \chi$ satisfies the hypotheses of the optimal-level theorem since $\bar{\rho}$ does. By Lemma 2.2 applied to E', the Serre conductor $M = N(\bar{\rho} \otimes \chi)$ has exponent exactly 1 at q, and $M \mid N/q$. The optimal-level theorem gives a newform g of weight 2 on $\Gamma_0(M)$ with $a_p(g) \equiv a_p(f') = \chi(p)a_p(f) \pmod{\lambda}$ for all $p \nmid 3N$. As $q \,\|\, M$, Lemma 2.4(iii) says that $h = g \otimes \chi$ is a newform on $\Gamma_0(Mq)$, and $Mq \mid N$. For $p \nmid 3N$,

$$a_p(h) = \chi(p)a_p(g) \equiv \chi(p)^2 a_p(f) = a_p(f) \pmod{\lambda}.$$

Moreover $h \neq f$: otherwise $a_p(g) = \chi(p)a_p(h) = a_p(f')$ for all $p \nmid qN$, so $g = f'$ by strong multiplicity one, whereas g has level $M < N$. (Note that $h \otimes \chi$ is the q-depletion of g, not g, as $a_q(g) = \pm 1$.) Finally, if $p \,\|\, N$ and $\bar{\rho}$ is unramified at p, then $p \neq q$, so $\bar{\rho} \otimes \chi$ is unramified at p as well and $p \nmid M$, hence $p \nmid Mq$. Lemma 3.2 gives $3 \mid r$.

Case (c). Here E has type I_n^* at an odd prime $q \neq \ell$ with $\ell \mid n$, so $q^2 \,\|\, N$. Let $\chi = \chi_{q^*}$ and $E' = E^{(q)}$. By Lemma 2.4, E' has multiplicative reduction of type I_n at q and conductor N/q; let $f' = f_{E'}$ be its newform, of weight 2 on $\Gamma_0(N/q)$ with $q \,\|\, N/q$, $a_q(f') = \pm 1$ and $a_p(f') = \chi(p)a_p(f)$ for $p \neq q$ (so $f \otimes \chi$ is the q-depletion of f'; for example 539c1 twisted by

χ_{-7} lies in the class 77c, and $a_7(77\text{c}1) = -1$). By Lemma 2.1 applied to E', $\bar\rho \otimes \chi = \bar\rho_{E',\ell}$ is unramified at q, so the Serre conductor $M = N(\bar\rho \otimes \chi)$ is prime to q and $M \mid N/q^2$. The optimal-level theorem (this is the case of Ribet's theorem [33], q exactly dividing the level and $\bar\rho \otimes \chi$ unramified at q) gives a newform g of weight 2 on $\Gamma_0(M)$ with $a_p(g) \equiv \chi(p)a_p(f) \pmod\lambda$ for all $p \nmid \ell N$. As $q \nmid M$, $h = g \otimes \chi$ is a newform on $\Gamma_0(Mq^2)$ with $Mq^2 \mid N$, congruent to f at all $p \nmid \ell N$ by the computation of case (b), and $h \neq f$: otherwise g and f' would have the same a_p for all $p \nmid qN$, so $g = f'$ by strong multiplicity one, whereas their levels M and N/q differ. The hypothesis of Lemma 3.2 holds as in case (b), and $\ell \mid r$. □

Example 3.3. The congruences produced by the proof can be seen directly on the tables (Cremona labels, as in [9] and `ecdata`); in each case the two forms lie in the same space $S_2(\Gamma_0(N))$ and have integer coefficients, so a check up to Sturm's bound [43], $\frac{1}{6}N\prod_{p\mid N}(1+\frac{1}{p})$, proves the congruence for all n. (a) at $q = 2$: 52a1 has type IV* at 2; its mod 3 representation has Serre conductor 26, and $f_{52\text{a}} \equiv f_{26\text{b}}(z) - f_{26\text{b}}(2z) \pmod 3$ for all n (Sturm's bound 14; $a_2(26\text{b}1) = 1$), while $\deg\varphi_{52\text{a}1} = 3$: here $e_3(E,2) = 1$ and the bound is attained. (b): the curve 121a1 has type II at 11; its twist by χ_{-11} lies in the class 121c and has type IV* at 11; its mod 3 representation has conductor 11 and is congruent to the form of 11a1; twisting back, $f_{11\text{a}} \otimes \chi_{-11}$ is the form of 121d, and indeed $f_{121\text{a}} \equiv f_{121\text{d}} \pmod 3$ for all n (Sturm's bound 22), while $\deg\varphi_{121\text{a}1} = 6$: here $e_3(E,11) = v_3(11+1) = 1$ and the bound is attained. (c) at $\ell = 3$: 539c1 has type I_3^* at 7; its twist by χ_{-7} has conductor 77 and type I_3 at 7, and $f_{539\text{c}} \equiv f_{11\text{a}} \otimes \chi_{-7} = f_{539\text{d}} \pmod 3$ for all n (Sturm's bound 112); $\deg\varphi_{539\text{c}1} = 2^5\cdot 3^2$, and the two terms $e_3(E,7) = v_3(48) = 1$ and $t_3(E,7) = v_3(3) = 1$ are both attained. (c) at $\ell = 5$: 980i1 has type I_5^* at 7; its twist by χ_{-7} has conductor 140 and type I_5 at 7, and $f_{980\text{i}} \equiv f_{20\text{a}} \otimes \chi_{-7} = f_{980\text{g}} \pmod 5$ for all n (Sturm's bound 336); $\deg\varphi_{980\text{i}1} = 2^6\cdot 3^2\cdot 5$. As $5 \mid 980$, this last example illustrates the congruence, not the theorem.

Remark 3.4. The hypothesis $\ell \nmid N$ is used in the optimal-level theorem, in Lemma 3.2 at $p = \ell$, and to pass from the congruence number to the modular degree. The last step needs only $\ell^2 \nmid N$, and it cannot be dropped there: Table 1 of [2] lists the curve 99A1, of conductor $9\cdot 11$, with modular degree 4 and congruence number 12; likewise 726a1 has $\deg\varphi_E = 144$ and $r_E = 1584 = 11\cdot 144$, with $11^2 \mid 726$, and 539d1 has $\deg\varphi_E = 72$ and $r_E = 504 = 7\cdot 72$, with $7^2 \mid 539$. The data nevertheless show the conclusion of the theorem for every curve with $\ell \parallel N$ or $\ell^2 \mid N$ and a prime of one of the three kinds; so do they at $q = 3$ in cases (a) and (b), where $q = \ell$ and the theorem does not apply.

4. The congruence ideal and the adjoint Selmer group

4.1. **Congruence ideals.** Let f be a newform of weight 2 on $\Gamma_0(M)$ with rational coefficients, $\lambda = \ell$ a prime not dividing $2M$, and Σ a finite set of primes. Put $M^\Sigma = M\prod_{p\in\Sigma} p^{\delta_p}$ and let $f^\Sigma = \sum_{(n,\Sigma)=1} a_n(f)q^n$ be the Σ-depleted eigenform, of level M^Σ. Diamond, Flach and Guo define [12, §§1.5.3, 1.6.2, 1.7.3] a lattice $M_{f^\Sigma,\lambda}$ and an ideal $\eta^\Sigma_{f,\lambda} \subset \mathbb{Z}_\ell$ as follows. Their integral structure $M(M^\Sigma,1)_{!,\lambda}$ is the image of the $U_0(M^\Sigma)$-invariants of the compactly supported cohomology of a modular curve of full level in its cohomology [12, §§1.2, 1.4], with its Hecke action [12, §1.6.1]. The following comparison identifies it with the cohomology of $X_0(M^\Sigma)$.

Proposition 4.1 (comparison of integral structures)**.** *Pull-back along the covering of $X_0(M^\Sigma)$ by the modular curve of full level identifies $H^1(X_0(M^\Sigma),\mathbb{Z})$ with the Betti lattice $M(M^\Sigma,1)_{!,B}$ of* [12, §1.4.2]*, compatibly with the Hecke operators, U_p included; in particular, for every prime $\ell \nmid 2M^\Sigma$ (the primes for which* [12] *define the λ-adic realisation), $M(M^\Sigma,1)_{!,\lambda} = H^1(X_0(M^\Sigma),\mathbb{Z}_\ell)$, with no localisation.*

Proof. Write $M = M^\Sigma$ and $U = U_0(M)$. For weight 2 and trivial character the coefficient sheaf of [12, §1.2.2] is $\mathbb{Z}$. Choose $N \geq 3$ with $M \mid N$ and every prime factor of N in the set S of [12, §1.4.1] (for instance $N = 4M$). Then $U_N \subseteq U$, and by [12, §1.4.1] $M(M,1)_{?,B} = H^1_?(Y(N),\mathbb{Z})^G$ for $? \in \{c, \mathrm{tf}\}$, where $Y(N)$ is the analytic modular curve of full level N, a disjoint union of copies of $\Gamma(N)\backslash\mathcal{H}$ indexed by $(\mathbb{Z}/N\mathbb{Z})^\times$, and $G = U/U_N$ acts through its image in $\mathrm{GL}_2(\mathbb{Z}/N\mathbb{Z})$ [12, §§1.2.1, 1.3.2]; $M(M,1)_{!,B}$ is the image of $M(M,1)_{c,B} \to M(M,1)_{\mathrm{tf},B}$, and $M(M,1)_{c,B}$ and $M(M,1)_{!,B}$ do not depend on N [12, §1.4.1, Lemma 1.1] (the lattice $M(M,1)_{\mathrm{tf},B}$ may, but only its image is used). Since $\det U = \hat{\mathbb{Z}}^\times$, G permutes the components transitively, the stabiliser of a component acts through the image of $\Gamma_0(M)$, and the quotient map $\pi\colon Y(N) \to Y(N)/G = Y_0(M)$ is finite onto the coarse curve; the stabiliser in G of a point of $Y(N)$ is finite.

Step 1: $\pi^*\colon H^1_c(Y_0(M),\mathbb{Z}) \to H^1_c(Y(N),\mathbb{Z})^G$ is an isomorphism. Let $H^\bullet_{G,c}$ denote G-equivariant cohomology with compact support. In the Hochschild-Serre spectral sequence $H^p(G, H^q_c(Y(N),\mathbb{Z})) \Rightarrow H^{p+q}_{G,c}(Y(N),\mathbb{Z})$ the row $q = 0$ vanishes, because every component of $Y(N)$ is a non-compact connected surface, so $H^0_c(Y(N),\mathbb{Z}) = 0$; hence $H^1_{G,c}(Y(N),\mathbb{Z}) = H^1_c(Y(N),\mathbb{Z})^G$. In the equivariant Leray spectral sequence of the finite map π, $H^p_c(Y_0(M),\mathcal{R}^q) \Rightarrow H^{p+q}_{G,c}(Y(N),\mathbb{Z})$, the stalk of $\mathcal{R}^q$ at y is $H^q(G_x,\mathbb{Z})$ for a point x above y (Shapiro's lemma), so $\mathcal{R}^0 = \mathbb{Z}$ and $\mathcal{R}^1 = 0$, since $H^1(C,\mathbb{Z}) = \mathrm{Hom}(C,\mathbb{Z}) = 0$ for a finite group C. Hence $H^1_{G,c}(Y(N),\mathbb{Z}) = H^1_c(Y_0(M),\mathbb{Z})$. The composite of the two edge isomorphisms is π^*.

Step 2: the image. Under these identifications $M(M,1)_{c,B} \to M(M,1)_{\mathrm{tf},B}$ is the forgetful map $H^1_c(Y(N),\mathbb{Z})^G \to H^1(Y(N),\mathbb{Z})^G$, and as π is proper its image is π^* of the image of $H^1_c(Y_0(M),\mathbb{Z}) \to H^1(Y_0(M),\mathbb{Z})$. Since $Y_0(M)$ is the compact surface $X_0(M)$ with finitely many points removed, $H^1_c(Y_0(M),\mathbb{Z}) \to H^1(X_0(M),\mathbb{Z})$ is surjective and $H^1(X_0(M),\mathbb{Z}) \to H^1(Y_0(M),\mathbb{Z})$ is injective, so that image is $H^1(X_0(M),\mathbb{Z})$. Finally π^* is injective on the torsion-free group $H^1(Y_0(M),\mathbb{Z})$, since $\pi_*\pi^*$ is multiplication by $\deg\pi$. So π^* maps $H^1(X_0(M),\mathbb{Z})$ isomorphically onto $M(M,1)_{!,B}$. In a diagram, the bottom row being the forgetful map factored through its image,

$$\begin{array}{ccccc} H^1_c(Y_0(M),\mathbb{Z}) & \twoheadrightarrow & H^1(X_0(M),\mathbb{Z}) & \hookrightarrow & H^1(Y_0(M),\mathbb{Z}) \\ \downarrow{\scriptstyle \pi^*,\ \simeq} & & \downarrow{\scriptstyle \pi^*,\ \simeq} & & \downarrow{\scriptstyle \pi^*} \\ H^1_c(Y(N),\mathbb{Z})^G & \twoheadrightarrow & M(M,1)_{!,B} & \hookrightarrow & H^1(Y(N),\mathbb{Z})^G \end{array}$$

with the left vertical map an isomorphism by Step 1 and the right one injective.

Step 3: Hecke operators. For a Hecke operator t, $\pi^* t$ and $t\pi^*$ are two homomorphisms from $H^1(X_0(M),\mathbb{Z})$ to the torsion-free group $M(M,1)_{!,B}$, which injects into $M(M,1)_{!,B}\otimes\mathbb{C}$; so if they agree after tensoring with $\mathbb{C}$ they agree integrally, and since π^* is an isomorphism of lattices (Step 2), the two Hecke actions then correspond under it. There the Hecke operators of [12] act on $\mathrm{Fil}^1 M(M,1)_{!,\mathrm{dR}} \otimes \mathbb{C} = S_2(\Gamma_0(M))$ as the classical T_p, and as U_p for $p \mid M$ [12, Proposition 1.3], and commute with complex conjugation F_∞; π^* is compatible with the comparison isomorphism (5) of [12] and with the Eichler-Shimura isomorphism, both being pull-back of $f(\tau)\,d\tau$; and the Hecke action on $H^1(X_0(M),\mathbb{Z})$ is the one for which the Eichler-Shimura isomorphism is Hecke equivariant (Section 4). Hence π^* is Hecke equivariant, U_p included.

Step 4: the λ-adic structure. By [12, §1.2.3], $\mathbb{Z}_\ell \otimes M_{?,B} \simeq M_{?,\ell}$ for every prime ℓ, and taking invariants under a finite group and images commutes with the flat base change $\mathbb{Z} \to \mathbb{Z}_\ell$; so $M(M,1)_{!,\lambda} = \mathbb{Z}_\ell \otimes M(M,1)_{!,B} = H^1(X_0(M),\mathbb{Z}_\ell)$. □

Remark 4.2. Compact support is what makes the comparison integral: H^0_c of the covering vanishes, so taking invariants under the covering group loses nothing, as in [12, Lemma 1.1]. With ordinary cohomology it is not so: $H^1(Y(N),\mathbb{Z}_\ell)^G/\pi^*H^1(Y_0(M),\mathbb{Z}_\ell)$ is the group of characters

$\Gamma_0(M) \to \mathbb{Q}_\ell/\mathbb{Z}_\ell$ of ℓ-power order that factor through G, namely $\gamma \mapsto \psi(d \bmod p)$ for primes $p \mid M$ with $\ell \mid p-1$; for $M = 11$ and $\ell = 5$ it is the unramified covering $X_1(11) \to X_0(11)$.

The lattice $M_{f^\Sigma,\lambda}$ is the kernel on $M(M^\Sigma,1)_{!,\lambda}$ of the kernel of the eigencharacter of f^Σ on the full Hecke algebra, U_p included [12, §§1.1.2, 1.6.2]. The pairing $\hat\delta_!$ is the cup product composed with the operator $[UwU]_\omega$ attached to $w = \left(\begin{smallmatrix} 0 & -1 \\ M^\Sigma & 0 \end{smallmatrix}\right)$ [12, §1.5.3, (13), (14)], which for weight 2 and trivial character is the Atkin-Lehner involution w_{M^Σ} with no scalar factor; on classes pulled back from $X_0(M^\Sigma)$ it is $\frac{u}{2}$ times the cup product on $X_0(M^\Sigma)$ composed with w_{M^Σ}, for an ℓ-adic unit u, so for odd ℓ only a unit separates the two. Indeed, for a sufficiently small level $U' \subseteq U_0(M^\Sigma)$ with $\det U' = \hat{\mathbb{Z}}^\times$, such as $U_0(M^\Sigma) \cap U_1(4)$, the normalised pairing $\bar\delta$ of [12, §1.5.2] is Poincaré duality on $X_{U'}$ up to an ℓ-adic unit when $\ell \nmid 2M^\Sigma$ [12, §1.5.2, before Lemma 1.2]; it is normalised by dividing by the index, so on $U_0(M^\Sigma)$-invariants it is $[U_0(M^\Sigma) : U']^{-1}$ times its value on U'-invariants, and the covering $X_{U'} \to X_0(M^\Sigma)$ has degree $[U_0(M^\Sigma) : U']/2$, because -1 acts trivially. It is perfect and alternating on $H^1(X_0(M^\Sigma), \mathbb{Z}_\ell)$, and the Hecke operators, U_p included, are self-adjoint for it, because $w_{M^\Sigma} T_n w_{M^\Sigma}^{-1}$ is the adjoint of T_n for the cup product (this is what the composition with w is for; see [12, §§1.6.1, 1.8.2] and [10, §4.4]). The restriction of $\hat\delta_!$ to the rank-two lattice $M_{f^\Sigma,\lambda}$ takes its values in an ideal of $\mathbb{Z}_\ell$, the image of $\wedge^2 M_{f^\Sigma,\lambda}$, which is $\eta^\Sigma_{f,\lambda}$, the *naive Σ-finite congruence ideal* [12, §1.7.3]; equivalently the Gram determinant of the pairing on $M_{f^\Sigma,\lambda}$ is $(\eta^\Sigma_{f,\lambda})^2$ up to a unit.

Theorem 4.3 ([12, Proposition 1.4(c)])**.** *Let $\ell \nmid 2M$ and $\ell \notin \Sigma$. If $\bar\rho_f$ is irreducible then*

$$\eta^\Sigma_{f,\lambda} = \eta^\emptyset_{f,\lambda} \prod_{p \in \Sigma} L_p^{\mathrm{nv}}(A_f, 1)^{-1}.$$

Only the primes with $\delta_p \geq 1$ contribute: adding an additive prime to Σ changes neither the level nor the ideal; adding a good prime p multiplies the ideal by $(1 - \frac{\alpha}{\beta}p^{-1})(1-p^{-1})(1-\frac{\beta}{\alpha}p^{-1})$, Wiles's level-raising factor; adding a multiplicative prime multiplies it by $1-p^{-2}$. The hypothesis $\ell \notin \Sigma$ is a standing assumption of [12, §1.7.3], and it cannot be dropped: for $f = f_{11\mathrm{a}}$, $\ell = 3$ and $\Sigma = \{3\}$ one has $a_3 = -1$ and $L_3^{\mathrm{nv}}(A_f,1)^{-1} = \frac{2}{3} \cdot \frac{16-1}{9} = \frac{10}{9}$, of 3-adic valuation -2, so the right side would not be contained in $\mathbb{Z}_3$. Every use below has Σ a set of primes dividing the level and ℓ prime to the level.

For the optimal curve, the congruence ideal at level N is the modular degree.

Proposition 4.4. *Let E be $X_0(N)$-optimal with newform f, $\varphi = \varphi_E$, I_f the kernel of the eigencharacter $\phi_f \colon \mathbb{T} \to \mathbb{Z}$ of f on the full Hecke algebra, and ℓ a prime. Then $\varphi^* \colon H^1(E, \mathbb{Z}_\ell) \to H^1(X_0(N), \mathbb{Z}_\ell)$ is injective, with image the kernel lattice $H^1(X_0(N), \mathbb{Z}_\ell)[I_f]$, and*

$$\langle \varphi^* a,\, w_N^* \varphi^* b \rangle_{X_0(N)} = \varepsilon_N(f)\, \deg \varphi_E\, \langle a, b \rangle_E \qquad (a, b \in H^1(E, \mathbb{Z}_\ell)),$$

where $\langle\ ,\ \rangle$ denotes the cup product and $\varepsilon_N(f) = \pm 1$ the eigenvalue of w_N on f. In particular, for ℓ odd, the ideal of values of $\frac{1}{2}\langle x, w_N^* y\rangle$ on $H^1(X_0(N), \mathbb{Z}_\ell)[I_f]$ is $\deg \varphi_E\, \mathbb{Z}_\ell$, and its Gram determinant is $(\deg \varphi_E)^2$ up to a unit. If $\ell \nmid 2N$, then $M_{f,\lambda} = \varphi^* H^1(E, \mathbb{Z}_\ell)$, and

$$v_\lambda(\eta^\emptyset_{f,\lambda}) = v_\ell(\deg \varphi_E).$$

Proof. We use Betti cohomology of the Riemann surfaces, with the Hecke action for which the Eichler-Shimura isomorphism $H^1(X_0(N), \mathbb{C}) = S_2(\Gamma_0(N)) \oplus \overline{S_2(\Gamma_0(N))}$ is Hecke equivariant. By optimality the kernel B of the map $J_0(N) \to E$ induced by φ is connected; the exact sequence $H_1(J_0(N), \mathbb{Z}) \to H_1(E, \mathbb{Z}) \to \pi_0(B) = 0$ shows that $\varphi_* \colon H_1(X_0(N), \mathbb{Z}) = H_1(J_0(N), \mathbb{Z}) \to H_1(E, \mathbb{Z})$ is surjective, so its dual $\varphi^* \colon H^1(E, \mathbb{Z}) \to H^1(X_0(N), \mathbb{Z})$ is injective with torsion-free cokernel, and the same holds after tensoring with $\mathbb{Z}_\ell$. The modular parametrisation satisfies

$\varphi^*\omega_E = c_E \cdot 2\pi i\, f(z)\, dz$ with the Manin constant $c_E > 0$, so $\varphi^* H^1(E,\mathbb{C})$, which is spanned by $\varphi^*\omega_E$ and its conjugate, is $\mathbb{C} f\, dz \oplus \mathbb{C}\overline{f\, dz}$. Since f is a newform, the eigenspace of ϕ_f in $S_2(\Gamma_0(N))$ is the line of f, so this plane is $H^1(X_0(N),\mathbb{C})[I_f]$. It is defined over $\mathbb{Q}$: $H^1(X_0(N),\mathbb{Q})[I_f]$ is a two-dimensional $\mathbb{Q}$-form of it, and by flatness $H^1(X_0(N),\mathbb{Z}_\ell)[I_f] = H^1(X_0(N),\mathbb{Z})[I_f] \otimes \mathbb{Z}_\ell$ is a saturated lattice of rank two in $H^1(X_0(N),\mathbb{Q}_\ell)[I_f]$. The image of φ^* is a saturated lattice of rank two in the same $\mathbb{Q}_\ell$-plane; so the two coincide. Next, $\varphi^*(a \cup b) = \varphi^* a \cup \varphi^* b$ and φ^* multiplies the fundamental class of H^2 by $\deg \varphi_E$; so $\langle \varphi^* a, \varphi^* b\rangle_{X_0(N)} = \deg \varphi_E \, \langle a, b\rangle_E$, the projection formula. By multiplicity one for newforms the plane is also the eigenspace of the T_n, $(n,N)=1$, for the eigenvalues $a_n(f)$, and w_N commutes with these T_n (not with the U_p, $p \mid N$); so w_N preserves the plane, on which it acts by $\varepsilon_N(f)$ ($f|w_N = \varepsilon_N(f) f$ and $\varepsilon_N(f)$ is real); this gives the displayed formula. The cup product on $H^1(E,\mathbb{Z})$ is unimodular, so the ideal of values is $\deg \varphi_E\, \mathbb{Z}_\ell$, and the Gram determinant of a rank-two alternating form is the square of its ideal of values. Finally, if $\ell \nmid 2N$, the integral structure of [12] is $H^1(X_0(N),\mathbb{Z}_\ell)$ (Proposition 4.1), so $M_{f,\lambda} = H^1(X_0(N),\mathbb{Z}_\ell)[I_f] = \varphi^* H^1(E,\mathbb{Z}_\ell)$, and $\hat{\delta}_!$ is $\frac{u}{2}\langle x, w_N^* y\rangle$ with u an ℓ-adic unit (Section 4.1); so $\eta_{f,\lambda}^{\emptyset} = \deg \varphi_E\, \mathbb{Z}_\ell$. □

No multiplicity-one hypothesis, and for the first part no irreducibility, is used. Two statements are to be kept apart. (i) For ℓ odd, the cohomological congruence ideal of the optimal curve, the ideal of values of $\frac{1}{2}\langle x, w_N^* y\rangle$ on $H^1(X_0(N),\mathbb{Z}_\ell)[I_f]$, is $\deg \varphi_E\, \mathbb{Z}_\ell$, with no condition on $\ell^2 \mid N$; for $\ell \nmid 2N$ it is the ideal $\eta_{f,\lambda}^{\emptyset}$ of [12]. (ii) The congruence number r_E, which generates the Hecke-theoretic congruence ideal (Lemma 3.1), has the same ℓ-valuation as $\deg \varphi_E$ when $\ell^2 \nmid N$ [2, Theorem 2.2], and not in general: 99A1, 726a1 and 539d1, at $\ell = 3$, 11 and 7, have $\ell^2 \mid N$ and $v_\ell(r_E) > v_\ell(\deg \varphi_E)$ (remark at the end of Section 3). The lattice $M_{f,\lambda}$ of [12] is therefore $H^1(E,\mathbb{Z}_\ell) = \mathrm{Hom}(T_\ell E, \mathbb{Z}_\ell) \simeq T_\ell E(-1)$, by Galois-equivariance of φ^* in étale cohomology, and its adjoint is $\mathrm{ad}^0\, T_\ell E$, the adjoint being insensitive to dualising and twisting.

We also need the following piece of commutative algebra, which is the inequality half of [10, Lemma 4.17]. Let $\mathcal{O}$ be a discrete valuation ring with finite residue field, $\mathbb{T}$ a commutative $\mathcal{O}$-algebra, finite and free as an $\mathcal{O}$-module, acting on an $\mathcal{O}$-module L free of finite rank with a perfect $\mathcal{O}$-bilinear pairing for which $\mathbb{T}$ is self-adjoint; let $\pi\colon \mathbb{T} \to \mathcal{O}$ be a surjective $\mathcal{O}$-algebra homomorphism, $\wp = \ker \pi$, $I = \mathrm{Ann}_{\mathbb{T}}(\wp)$ and $\eta = \pi(I)$, the congruence ideal of π; and suppose that $L[\wp]$ is free of rank d over $\mathcal{O}$ with Gram determinant $\Delta \neq 0$.

Lemma 4.5. *$\eta^d \subseteq \Delta\mathcal{O}$, that is, $d \cdot v(\eta) \geq v(\Delta)$; and equality holds if L is free over $\mathbb{T}$.*

Proof. If $\eta = 0$ there is nothing to prove; let $t \in I$ with $\pi(t)$ a generator of $\eta \neq 0$. First $\wp \cap I = 0$: for $x \in \wp \cap I$, $tx = 0$ as t kills $\wp$, and $(\pi(t) - t)x = 0$ as $\pi(t) - t \in \wp$ and x kills $\wp$; so $\pi(t)x = 0$ and $x = 0$, $\mathbb{T}$ being torsion-free. (This does not follow from reducedness: the full Hecke algebra can have nilpotents, for instance U_2 at level 88 on the span of the $f_{11\mathrm{a}}(dz)$, $d \mid 8$.) Consequently $\pi|_I$ is injective, with image $\eta = \pi(t)\mathcal{O}$, so $I = \mathcal{O}t$. The sum $L[\wp] + L[I]$ is direct: $\pi(t) = t + (\pi(t) - t)$ lies in $I + \wp$ and is a nonzero scalar, so an element of L killed by $\wp$ and by I is killed by $\pi(t)$, hence is 0, L being torsion-free. For $x \in L$, one has $tx \in L[\wp]$ (as $\wp I = 0$) and $(\pi(t) - t)x \in L[I]$ (as $s(\pi(t) - t) \in \wp \cap I = 0$ for $s \in I$), so $Q = L/(L[\wp] \oplus L[I])$ is killed by η. The pairing identifies $L/L[\wp]^\perp$ with $\mathrm{Hom}(L[\wp], \mathcal{O})$, because $L[\wp]$ is saturated, and $L[\wp]^\perp = L[I]$: for $y \in L[I]$ and $x \in L[\wp]$, $\pi(t)\langle x, y\rangle = \langle tx, y\rangle = \langle x, ty\rangle = 0$; conversely if $y \perp L[\wp]$ then $\langle ty, z\rangle = \langle y, tz\rangle = 0$ for all z, since $tz \in L[\wp]$, so $ty = 0$, and $y \in L[I]$ because $I = \mathcal{O}t$. Hence Q is the cokernel of $L[\wp] \to \mathrm{Hom}(L[\wp], \mathcal{O})$, of order $\#\mathcal{O}/\Delta$, and it is generated by d elements and killed by η, so $\#\mathcal{O}/\Delta \leq \#(\mathcal{O}/\eta)^d$. If $L \simeq \mathbb{T}^d$ then $Q \simeq (\mathbb{T}/(\wp + I))^d \simeq (\mathcal{O}/\eta)^d$. □

Lemma 4.6. *Let E be $X_0(N)$-optimal and ℓ an odd prime with $\ell \nmid 2N$ and $\bar{\rho}$ irreducible. Then $v_\ell(\deg \varphi_E) = v_\ell(r_E) = v_\lambda(\eta_{f,\lambda}^{\emptyset})$.*

Proof. $v_\ell(\deg\varphi_E) = v_\ell(r_E)$ is [2, Theorem 2.2], as $\ell^2 \nmid N$, and $v_\ell(\deg\varphi_E) = v_\lambda(\eta^{\emptyset}_{f,\lambda})$ is Proposition 4.4. □

Let $\mathfrak{m}$ be the maximal ideal of $\mathbb{T}\otimes\mathbb{Z}_\ell$ containing ℓ and $\ker\phi_f$. Lemma 4.5 applied to $L = H^1(X_0(N),\mathbb{Z}_\ell)_\mathfrak{m}$ gives, without the projection formula, only $v_\ell(r_E) \geq v_\lambda(\eta^{\emptyset}_{f,\lambda})$, with equality when L is free over $\mathbb{T}_\mathfrak{m}$. That argument is the one used in Step 2 of the proof of Theorem 1.6, for a depleted form that is not attached to an optimal quotient. In it the congruence ideal is computed in $\mathbb{T}_\mathfrak{m}$: the other local factors of $\mathbb{T}\otimes\mathbb{Z}_\ell$ lie in the kernel $\wp$, ϕ_f factors through $\mathbb{T}_\mathfrak{m}$, and $\mathrm{Ann}(\wp)$ decomposes along the idempotents; and the idempotent $e_\mathfrak{m}$ is self-adjoint for $\hat{\delta}_!$, so $H^1(X_0(N),\mathbb{Z}_\ell)$ splits orthogonally into its local components and the pairing is perfect on each. Without the hypotheses the inequality of Lemma 4.5 can be strict: at 99a1 and $\ell = 3$, where $3 \mid N$, $\deg\varphi_E = 4$ and $r_E = 12$, so $H^1(X_0(99),\mathbb{Z}_3)_\mathfrak{m}$ is not free over $\mathbb{T}_\mathfrak{m}$.

Remark 4.7. Freeness of $H^1(X_0(N),\mathbb{Z}_\ell)_\mathfrak{m}$ over $\mathbb{T}_\mathfrak{m}$ holds when $J_0(N)[\mathfrak{m}]$ is two-dimensional, since the localised Hecke algebra is then Gorenstein and the Tate module is free of rank two [47]; and $J_0(N)[\mathfrak{m}]$ is two-dimensional for $\ell \nmid 2N$ and $\bar\rho$ irreducible: [35, Theorem 3.5] gives it for $J_1(N)$, the only possible exception there having Serre weight ℓ, which does not occur in weight 2 with $\ell \nmid N$, and $J_0(N)[\mathfrak{m}]$ injects into $J_1(N)[\mathfrak{m}']$ under the pull-back map, whose kernel, the Shimura subgroup, is Eisenstein. (The identification of $T_\ell J_0(N)$ with $H^1(X_0(N),\mathbb{Z}_\ell)$ as Hecke modules uses Poincaré duality composed with w_N, which is not Galois equivariant; freeness transfers nevertheless.) After Proposition 4.4 this is not needed for Theorem 1.2 or for Theorem 1.6; with Lemma 4.5 it gives a second proof of $v_\ell(r_E) = v_\ell(\deg\varphi_E)$ for $\ell \nmid 2N$ and $\bar\rho$ irreducible, and it compares the Hecke and the cohomological congruence ideals of forms not attached to optimal quotients.

4.2. **The theorem of Diamond, Flach and Guo.** Let now $f = f_E$, Σ the set of primes dividing N, $\lambda = \ell \nmid 2N$, $A = A_{f,\lambda} = \mathrm{ad}^0 T$ (by Proposition 4.4, φ_E^* identifies the lattice $M_{f,\lambda}$ of [12] with $H^1(E,\mathbb{Z}_\ell) \simeq T_\ell E(-1)$, so its adjoint is $\mathrm{ad}^0 T$), $W = A\otimes\mathbb{Q}_\ell/\mathbb{Z}_\ell$, $T^D = \mathrm{Hom}(A,\mathbb{Z}_\ell(1)) \simeq A(1)$ (via the trace form), $V^D = T^D\otimes\mathbb{Q}_\ell$ and $W^D = V^D/T^D$. Let $H^1_f(\mathbb{Q},W)$ be the Bloch-Kato Selmer group and $H^1_\Sigma(\mathbb{Q},W) \supseteq H^1_f(\mathbb{Q},W)$ the group defined in [12, §2.1], with no condition at the primes of Σ, the Bloch-Kato condition at ℓ and the unramified condition elsewhere. Let S_f be the set of primes λ such that $\lambda \mid 2N$ or $\bar\rho|_{\mathrm{Gal}(\overline{\mathbb{Q}}/\mathbb{Q}(\sqrt{\ell^*}))}$ is not absolutely irreducible [12, (31)]; the hypothesis of Theorem 1.2 is $\ell \notin S_f$.

Proposition 4.8. *Let ℓ be an odd prime with $\ell \nmid N$ and $E[\ell]$ irreducible. If $\ell \geq 5$, then $\ell \notin S_f$. If $\ell = 3$, then $3 \in S_f$ if and only if the image of $\bar\rho_{E,3}$ is conjugate to $N_s(\mathbb{F}_3)$, and then E is ordinary at 3. In all cases $\ell \notin S_f$ if and only if $h_\ell(E) = 0$.*

Proof. Let $K = \mathbb{Q}(\sqrt{\ell^*})$. As $\bar\rho$ is absolutely irreducible (complex conjugation has distinct eigenvalues), $\bar\rho|_{G_K}$ fails to be absolutely irreducible exactly when $\bar\rho \simeq \bar\rho\otimes\chi_{\ell^*}$ (Clifford theory, as in the proof of Lemma 4.13). Suppose so, and restrict to the inertia group I_ℓ, where $\chi_{\ell^*} = \omega^{(\ell-1)/2}$, with ω the fundamental character of level one, of order $\ell-1$. As E has good reduction at ℓ, by [38, §1] either E is ordinary at ℓ and the semisimplification of $\bar\rho|_{I_\ell}$ is $\omega\oplus 1$, or E is supersingular at ℓ and $\bar\rho|_{I_\ell}\otimes\mathbb{F}_{\ell^2} \simeq \omega_2\oplus\omega_2^\ell$, with ω_2 of order ℓ^2-1 and $\omega_2^{\ell+1} = \omega$. In the ordinary case $\{\omega,1\} = \{\omega^{(\ell+1)/2},\omega^{(\ell-1)/2}\}$; as $\omega^{(\ell-1)/2} \neq 1$, this forces $\omega^{(\ell+1)/2} = 1$, so $(\ell-1) \mid (\ell+1)/2$ and $\ell = 3$. In the supersingular case $\omega^{(\ell-1)/2} = \omega_2^{(\ell^2-1)/2} = -1$, and $\{\omega_2,\omega_2^\ell\} = \{-\omega_2,-\omega_2^\ell\}$ would need $\omega_2^{\ell-1} = -1 = \omega_2^{(\ell^2-1)/2}$, so $(\ell+1) \mid (\ell-1)/2$, which is impossible. So $\ell \geq 5$ never occurs, and for $\ell = 3$ the curve is ordinary at 3; the image is then $N_s(\mathbb{F}_3)$ by Proposition 1.7. The last statement follows from Lemma 4.13. □

So for $\ell \nmid 2N$ the hypothesis of Theorem 1.2 reads: $E[\ell]$ is irreducible, and, if $\ell = 3$, the mod 3 image is not $N_s(\mathbb{F}_3)$.

Theorem 4.9 ([12, Theorem 3.7])**.** *If $\ell \notin S_f$, then $H^1_\Sigma(\mathbb{Q}, W)$ is finite of length $v_\lambda(\eta^\Sigma_{f,\lambda})$.*

The theorem is stated in [12] for a finite set Σ not containing ℓ such that $M_{f,\lambda}$ is minimally ramified outside Σ, which holds for the set of primes dividing N, outside of which ρ is unramified; this is the form given as Theorem 0.2 there. It covers $\ell = 3$ in weight 2: S_f excludes only the primes dividing $2N$ and the induced case.

Theorem 4.10 ([12, Theorem 2.7])**.** *If $\ell \notin S_f$, then $H^0(\mathbb{Q}, V^D) = H^1_f(\mathbb{Q}, V^D) = 0$ (and the same for $A \otimes \mathbb{Q}_\ell$).*

Lemma 4.11 ([12, Lemma 2.1])**.** *Suppose $H^0(\mathbb{Q}, V^D) = H^1_f(\mathbb{Q}, V^D) = 0$, and let Σ be a nonempty finite set of primes, with $\ell \notin \Sigma$, containing every prime $p \neq \ell$ at which W is ramified. Put $H^1_f(\mathbb{Q}_p, T^D) = \iota^{-1} H^1_f(\mathbb{Q}_p, V^D)$, $\iota\colon H^1(\mathbb{Q}_p, T^D) \to H^1(\mathbb{Q}_p, V^D)$. Then there is an exact sequence*

$$0 \to H^1_f(\mathbb{Q}, W) \to H^1_\Sigma(\mathbb{Q}, W) \to \bigoplus_{p \in \Sigma \cup \{\infty\}} H^1_f(\mathbb{Q}_p, T^D)^* \to H^0(\mathbb{Q}, W^D)^* \to 0,$$

where * is the Pontryagin dual.

The condition on Σ implies the one of [12], which asks that Σ contain every prime at which $H^1_f(\mathbb{Q}_p, W) \neq H^1_{\mathrm{ur}}(\mathbb{Q}_p, W)$. For $p \in \Sigma$ the group $H^1_f(\mathbb{Q}_p, V^D) = H^1_{\mathrm{ur}}(\mathbb{Q}_p, V^D) = (V^D)^{I_p}/(\mathrm{Frob}_p - 1)$ vanishes, because $\det(1 - \mathrm{Frob}_p^{-1} \mid (V^D)^{I_p}) = L_p(A_f, 1)^{-1} \neq 0$; for the same reason $H^0(\mathbb{Q}_p, V^D) = 0$, so the hypothesis $H^0(\mathbb{Q}, V^D) = 0$ is automatic here, and only $H^1_f(\mathbb{Q}, V^D) = 0$ needs Theorem 4.10. So $H^1_f(\mathbb{Q}_p, T^D)$ is the torsion of $H^1(\mathbb{Q}_p, T^D)$, a finite group. This computation needs $p \neq \ell$, which is why $\ell \notin \Sigma$ is part of the lemma: at $p = \ell$ the group $H^1_f(\mathbb{Q}_\ell, V^D)$ has dimension 2. At ∞, $H^1(\mathbb{R}, T^D)$ is killed by 2, so $H^1_f(\mathbb{R}, T^D) = 0$ for ℓ odd.

Lemma 4.12 ([17, I.4.2.2], as recalled in the proof of [12, Theorem 2.15])**.** *For $p \in \Sigma$,*

$$\mathrm{Fitt}_{\mathbb{Z}_\ell} H^1_f(\mathbb{Q}_p, T^D) = L_p(A_f, 1)^{-1} \cdot \mathrm{Tam}_p(T^D), \qquad \text{and} \qquad \mathrm{Tam}_p(T^D) = \mathrm{Tam}_p(A)$$

[12, (57)]; here $L_p(A_f, 1)^{-1} = L_p(A_f(1), 0)^{-1}$ is a nonzero rational number which is an ℓ-adic integer, of valuation $e_\ell(E, p)$ at an additive prime and $v_\ell(1 - p^{-2})$ at a multiplicative one.

Lemma 4.13. *Let ℓ be odd with $\bar\rho$ irreducible. Then $H^0(\mathbb{Q}, W^D) = H^0(\mathbb{Q}, \mathrm{ad}^0\, T(1) \otimes \mathbb{Q}_\ell/\mathbb{Z}_\ell)$ is finite and cyclic, of length $h_\ell(E)$, the largest k such that $\rho \bmod \ell^k \simeq (\rho \otimes \chi_\ell) \bmod \ell^k$. Moreover $h_\ell(E) = 0$ for every $\ell \geq 5$, and $h_3(E) \leq 1$, with equality if and only if $\bar\rho_{E,3} \simeq \bar\rho_{E,3} \otimes \chi_{-3}$, if and only if $\bar\rho_{E,3}$ is not absolutely irreducible on $\mathrm{Gal}(\overline{\mathbb{Q}}/\mathbb{Q}(\sqrt{-3}))$, if and only if the image of $\bar\rho_{E,3}$ is conjugate to the normaliser $N_s(\mathbb{F}_3)$ of a split Cartan subgroup. This holds for every E, with or without complex multiplication; in particular $H^0(\mathbb{Q}, W^D) = 0$ if $\ell \notin S_f$.*

Proof. The representation $\bar\rho$ is absolutely irreducible, complex conjugation having distinct eigenvalues ± 1. The ℓ^k-torsion of $H^0(\mathbb{Q}, W^D)$ is $H^0(\mathbb{Q}, \mathrm{ad}^0(T/\ell^k)(1))$, the group of trace-zero $G_\mathbb{Q}$-maps $T/\ell^k \to (T/\ell^k)(1)$; every such map has trace 0 automatically, since $\mathrm{tr}\,\phi = \chi_\ell(g)\,\mathrm{tr}\,\phi$ for every g and $1 - \chi_\ell(c) = 2$ is a unit for complex conjugation c. For $k = 1$ these are the $G_\mathbb{Q}$-maps $\bar\rho \to \bar\rho \otimes \bar\chi_\ell$, which form a space of dimension at most one by Schur's lemma; so $H^0(\mathbb{Q}, W^D)$, a submodule of $(\mathbb{Q}_\ell/\mathbb{Z}_\ell)^3$ with ℓ-torsion of rank at most one, is finite cyclic or isomorphic to $\mathbb{Q}_\ell/\mathbb{Z}_\ell$; the determinant argument below shows that it is finite. An element of exact order ℓ^k is a map $\phi\colon T/\ell^k \to (T/\ell^k)(1)$ whose reduction modulo ℓ is nonzero, hence an isomorphism, so ϕ

is an isomorphism by Nakayama's lemma, and $\rho \bmod \ell^k \simeq (\rho \otimes \chi_\ell) \bmod \ell^k$; conversely such an isomorphism gives such an element. Taking determinants, $\chi_\ell(g)^2 \equiv 1 \pmod{\ell^k}$ for all $g \in G_{\mathbb{Q}}$. Since $\chi_\ell(G_{\mathbb{Q}}) = \mathbb{Z}_\ell^\times$ contains 2, and $4 \not\equiv 1$ modulo ℓ for $\ell \geq 5$ and modulo 9, this forces $k = 0$ for $\ell \geq 5$ and $k \leq 1$ for $\ell = 3$. So $H^0(\mathbb{Q}, W^D)$ is finite, $h_\ell(E) = 0$ for $\ell \geq 5$, and $h_3(E) \leq 1$, with equality if and only if $\bar\rho \simeq \bar\rho \otimes \bar\chi_3^{\mathrm{cyc}}$, and $\bar\chi_3^{\mathrm{cyc}} = \chi_{-3}$. Let $K = \mathbb{Q}(\sqrt{-3})$. If $\phi\colon \bar\rho \to \bar\rho \otimes \chi_{-3}$ is an isomorphism, it commutes with $\bar\rho(G_K)$ and is not scalar (otherwise χ_{-3} would be trivial), so $\bar\rho|_{G_K}$ is not absolutely irreducible. Conversely, if $\bar\rho|_{G_K} \otimes \overline{\mathbb{F}}_3$ has a stable line with character ψ, Frobenius reciprocity gives a nonzero map $\operatorname{Ind}_{G_K}^{G_{\mathbb{Q}}} \psi \to \bar\rho \otimes \overline{\mathbb{F}}_3$, an isomorphism since $\bar\rho$ is absolutely irreducible, and $\operatorname{Ind}\psi \otimes \chi_{-3} \simeq \operatorname{Ind}\psi$; the isomorphism $\bar\rho \simeq \bar\rho \otimes \chi_{-3}$ then exists over $\mathbb{F}_3$, since spaces of homomorphisms commute with extension of scalars. The last equivalence is the first statement of Proposition 1.7 ($\bar\rho$ irreducible means $E[3]$ irreducible), whose proof uses this lemma only for the value $h_3(E) = 1$. □

4.3. **Proof of Theorem 1.2.** The proof is a chain of identifications. In the order used below, with the hypotheses each step needs and its source:

step	hypotheses	status
$H^1(X_0(N), \mathbb{Z}_\ell)$ is the integral structure of [12] (Proposition 4.1)	Betti form: any ℓ; ℓ-adic form: $\ell \nmid 2N$	proved here
$v_\lambda(\eta_{f,\lambda}^{\emptyset}) = v_\ell(\deg \varphi_E)$ (Proposition 4.4, Lemma 4.6)	E optimal; $\ell \nmid 2N$	proved here
$\eta_{f,\lambda}^{\Sigma} = \eta_{f,\lambda}^{\emptyset} \prod_{p\in\Sigma} L_p^{\mathrm{nv}}(A_f, 1)^{-1}$ (Theorem 4.3)	$\ell \notin \Sigma$; $\bar\rho$ irreducible	[12, Proposition 1.4(c)]
$v_\lambda(\eta_{f,\lambda}^{\Sigma}) = \operatorname{length} H^1_\Sigma(\mathbb{Q}, W)$ (Theorem 4.9)	$\ell \notin S_f$; $\Sigma = \{p \mid N\}$	[12, Theorem 3.7]
$H^0(\mathbb{Q}, V^D) = H^1_f(\mathbb{Q}, V^D) = 0$ (Theorem 4.10)	$\ell \notin S_f$	[12, Theorem 2.7]
$H^1_\Sigma(\mathbb{Q}, W)$ against $H^1_f(\mathbb{Q}, W)$ (Lemma 4.11)	the previous row; $\ell \notin \Sigma$; and $H^0(\mathbb{Q}, W^D) = 0$, from $\ell \notin S_f$ (Lemma 4.13), to drop the last term	[12, Lemma 2.1]
$\operatorname{Fitt} H^1_f(\mathbb{Q}_p, T^D) = L_p(A_f, 1)^{-1} \operatorname{Tam}_p(A)$ (Lemma 4.12)	$p \in \Sigma$, $p \neq \ell$	[17, I.4.2.2], [12, (57)]
local terms: e_ℓ (Definition 1.1, values in Lemma 2.5), $\operatorname{Tam}_q(\mathrm{ad}^0 T)$ (Lemmas 2.8 and 2.9)	ℓ odd, $q \neq \ell$; for Lemma 2.5, $q \geq 5$, or q odd in the rows I_n^*, I_0^*	proved here

In the last row (1) needs only Definition 1.1; the lemmas make the terms explicit and give (2) and Corollary 4.14.

Let $\ell \notin S_f$, Σ the set of primes dividing N. By Lemma 4.6, $v_\ell(\deg \varphi_E) = v_\lambda(\eta_{f,\lambda}^{\emptyset})$. By Theorem 4.3,

$$v_\lambda(\eta_{f,\lambda}^{\emptyset}) = v_\lambda(\eta_{f,\lambda}^{\Sigma}) - \sum_{p\in\Sigma} v_\ell\big(L_p^{\mathrm{nv}}(A_f, 1)^{-1}\big).$$

By Theorem 4.9, $v_\lambda(\eta_{f,\lambda}^{\Sigma}) = \operatorname{length} H^1_\Sigma(\mathbb{Q}, W)$. By Theorem 4.10 the hypotheses of Lemma 4.11 hold, and $H^0(\mathbb{Q}, W^D) = 0$ by Lemma 4.13, so

$$\operatorname{length} H^1_\Sigma(\mathbb{Q}, W) = \operatorname{length} H^1_f(\mathbb{Q}, W) + \sum_{p\in\Sigma} \operatorname{length} H^1_f(\mathbb{Q}_p, T^D),$$

the case $H^0(\mathbb{Q}, W^D) = 0$ of the sequence, which $\ell \notin S_f$ guarantees; when $h_\ell(E) > 0$, which needs $\ell \in S_f$, the right side acquires the term $-h_\ell(E)$ of Remark 5.2. By Lemma 4.12 each local

term is $v_\ell(L_p(A_f,1)^{-1}) + v_\ell(\mathrm{Tam}_p(A))$. Hence

$$v_\ell(\deg\varphi_E) = \operatorname{length} H^1_f(\mathbb{Q},W) + \sum_{p\in\Sigma} v_\ell\big(\mathrm{Tam}_p(\mathrm{ad}^0\,T)\big) + \sum_{p\in\Sigma} v_\ell\Big(\frac{L_p(A_f,1)^{-1}}{L_p^{\mathrm{nv}}(A_f,1)^{-1}}\Big),$$

and the last sum is $\sum_{p\mid N} e_\ell(E,p)$ by Definition 1.1: at a multiplicative prime the true and the naive factors coincide, at an additive prime the naive factor is 1. This is (1). For (2), the Selmer length is nonnegative, $v_\ell(\mathrm{Tam}_p(\mathrm{ad}^0\,T)) \geq t_\ell(E,p)$ at every potentially multiplicative prime p, multiplicative or additive, $p=2$ included, by Lemma 2.8, and $t_\ell(E,p)=0$ at the other primes. □

Corollary 4.14. *Under the hypothesis of Theorem 1.2,*

$$v_\ell(\deg\varphi_E) = \operatorname{length} H^1_f(\mathbb{Q},\mathrm{ad}^0\,T\otimes\mathbb{Q}_\ell/\mathbb{Z}_\ell) + \sum_{q\mid N} e_\ell(E,q) + \sum_{q:\,v_q(j_E)<0} \Big(t_\ell(E,q) + \min\big(t_\ell(E,q), v_\ell(q-1), v_\ell(c(u_q))\big)\Big),$$

where u_q is the unit part of the Tate parameter at q and $c(u_q)$ the class of its reduction, as in Lemma 2.8.

Proof. Theorem 1.2, with $\mathrm{Tam}_q(\mathrm{ad}^0\,T)$ trivial at the potentially good additive primes (Lemma 2.9) and given by Lemma 2.8 at the potentially multiplicative ones. □

So the excess of $v_\ell(\deg\varphi_E)$ over the explicit local bound is exactly the length of the adjoint Bloch-Kato Selmer group. Table 1 gives its distribution on the optimal curves without complex multiplication of conductor below 10,000 with $\ell\nmid N$, $E[\ell]$ irreducible and, for $\ell=3$, image not 3Ns, with e_ℓ from PARI's `lfunsympow`. The excess is never negative, as the corollary requires; for instance 1184f1 (explicit bound 2, $\deg\varphi_E=1728$) and 1748e1 (explicit bound 3, $\deg\varphi_E=3240$) have Selmer length 1 at $\ell=3$.

ℓ	curves	excess 0	1	2	3	≥ 4
3	13,203	8,688	3,288	942	248	37
5	23,361	19,019	3,993	328	19	2
7	28,134	25,133	2,888	111	2	0

TABLE 1. The excess $v_\ell(\deg\varphi_E) - \sum_q e_\ell(E,q) - \sum_{v_q(j_E)<0}\big(t_\ell(E,q) + \min(t_\ell(E,q), v_\ell(q-1), v_\ell(c(u_q)))\big)$, which by Corollary 4.14 is length $H^1_f(\mathbb{Q},\mathrm{ad}^0\,T\otimes\mathbb{Q}_\ell/\mathbb{Z}_\ell)$, on the optimal non-CM curves of conductor below 10,000 with $\ell\nmid N$, $E[\ell]$ irreducible and, at $\ell=3$, image not 3Ns.

Remark 4.15. The Euler factors also appear in an exact formula. By the formula of Zagier and Watkins [50, 48], $\deg\varphi_E\cdot\mathrm{vol}(\Lambda_E) = 4\pi^2 c_E^2\langle f,f\rangle$, with the unnormalised Petersson product and the Manin constant $c_E>0$ in the numerator; and $\langle f,f\rangle = N\,L^{\mathrm{nv}}(\mathrm{Sym}^2 f,2)/(8\pi^3)$ at every level [32, 41], [19, Theorem 5.1], [12, Lemma 2.12]. Hence

$$\deg\varphi_E = \frac{c_E^2\,N\,L^{\mathrm{nv}}(\mathrm{ad}^0 f,1)}{2\pi\,\mathrm{vol}(\Lambda_E)} = B_E\prod_{q^2\mid N} L_q(A_f,1)^{-1}, \qquad B_E = \frac{c_E^2\,N\,L(\mathrm{ad}^0 f,1)}{2\pi\,\mathrm{vol}(\Lambda_E)} \in \mathbb{Q}^\times,$$

so that $v_\ell(\deg\varphi_E) = v_\ell(B_E) + \sum_{q|N} e_\ell(E,q)$ for every $\ell \nmid N$, with no hypothesis, and (2) is equivalent to $v_\ell(B_E) \geq \sum_q t_\ell(E,q)$. With $c_E = 1$ the first formula reproduces $\deg\varphi_E$ to 30 digits for 11a1, 121a1, 539d1, 1232l1 and 726a1 (degrees 1, 6, 72, 288, 144).

Remark 4.16. (1) At a potentially multiplicative prime $q \equiv 1 \pmod\ell$ with $\ell \mid n$ whose Tate parameter has an ℓ-th power as unit part, Lemma 2.8 gives $v_\ell(\mathrm{Tam}_q) \geq t_\ell(E,q)+1$, and (1) raises the bound accordingly; Section 6.4 tests this on the tables. (2) The formula of Kim and Ota [21, Theorem 1.3] concerns the same congruence ideal: for $N = N^+N^-$ with N^- squarefree, $\ell \geq 5$, $\ell \nmid N$, $\ell \notin S_f$, and $\bar\rho$ ramified at every $q \mid N^-$ with $q \equiv \pm1 \pmod\ell$, it says $\mathrm{ord}_\lambda \eta_f(N) = \mathrm{ord}_\lambda \eta_f(N^+,N^-) + \sum_{q|N^-} t_f(q)$, where $\eta_f(N^+,N^-)$ is the congruence ideal in the N^--new quotient of the Hecke algebra and $t_f(q)$ the Tamagawa exponent [21, Definition 1.15], which is $t_\ell(E,q)$ by Lemma 2.1. Since $\eta_f(N^+,N^-)$ is integral, this gives $v_\ell(r_E) \geq \sum_{q|N^-} t_\ell(E,q)$, which (2) contains, without the conditions $\ell \geq 5$ and $q \not\equiv \pm1$; the equality is theirs.

4.4. Twist monotonicity and the Euler terms without the hypothesis on $\mathbb{Q}(\sqrt{\ell^*})$.

Theorem 4.17. *Let $E/\mathbb{Q}$ have conductor N, S a set of odd primes, $D = \prod_{q\in S} q$, $\chi_S = \prod_{q\in S}\chi_{q^*}$, and $E_S = E\otimes\chi_S$ of conductor N_S. If $q^2 \mid N$ for every $q \in S$ and $\mathrm{lcm}(N_S, D^2) \mid N$, then $r_{E_S} \mid r_E$; if moreover $\mathrm{lcm}(N, D^2) \mid N_S$, then $r_{E_S} = r_E$.*

Proof. Let $f = f_E$, $g = f_{E_S}$ and $\iota\colon h \mapsto h\otimes\chi_S = \sum \chi_S(n)a_n(h)q^n$, which maps $S_2(\Gamma_0(N_S))$ into $S_2(\Gamma_0(\mathrm{lcm}(N_S,D^2))) \subset S_2(\Gamma_0(N))$, preserves integrality, and sends an eigenform of all T_n, $(n,N_S)=1$, to an eigenform with eigenvalues twisted by $\chi_S(n)$. Then $\iota(g) = f$: both are multiplicative, $a_p(f) = \chi_S(p)a_p(g)$ for $p \nmid D$ since the twist is unramified outside S, and $a_{q^k}(f) = 0$ for $q \in S$ since $q^2 \mid N$. If h is a generalised eigenform of level dividing N_S for a system of T_p-eigenvalues, $p \nmid N_S$, differing from that of g at infinitely many p, then $\iota(h)$ is a generalised eigenform for a system differing from that of f at infinitely many p, so $\iota(h) \in f^\perp$; hence ι maps $g^\perp$, the $\mathbb{T}$-stable complement of the g-line (the sum of the generalised eigenspaces of the other systems of eigenvalues), into $f^\perp$. A congruence $g \equiv h \pmod r$ with $h \in g^\perp$ integral therefore gives $f \equiv \iota(h) \pmod r$ with $\iota(h) \in f^\perp$ integral, and $r \mid r_E$ by Lemma 3.1. With $r = r_{E_S}$ this is the first claim; the second hypothesis makes the situation symmetric. □

The hypotheses hold when (a) E has type I_n^* at each $q \in S$ ($N_S = N/D$), (b) type I_0^* at each $q \in S$ ($N_S = N/D^2$, E_S good at S), (c) type II, III, IV or a dual at each $q \in S$, $q \geq 5$ ($N_S = N$, equality). On the working set, for the 157,938 pairs (curve, prime of type I_n^*) and the 292,580 pairs (curve, prime of potentially good additive type), $v_\ell(\deg\varphi_E) \geq v_\ell(\deg\varphi_{E_0^{(q)}})$, with $E_0^{(q)}$ the optimal curve of the isogeny class of $E^{(q)}$, holds for $\ell \in \{3,5,7,11,13\}$, $\ell^2 \nmid N$, with no exception, with equality in every pair of case (c). In cases (a) and (b) Theorem 1.6 predicts a difference equal to the Euler term, which is positive in case (a) at $\ell = 3$ for every $q \neq 3$, as $3 \mid q^2-1$: for instance v_3 of the degrees is 2 against 1 for 539c1 and 77c1, 2 against 1 for 980i1 and 140b1, 1 against 0 for 121d1 and 11a1, and 2 against 0 for 539d1 (type I_0^*, $e_3(E,7)=2$) and 11a1. Below conductor 10,000 the difference equals the Euler term in all 25,477 pairs (curve, ℓ) with $S \neq \emptyset$, $\ell \in \{3,5,7,11,13\}$, $\ell \nmid N$ and $E[\ell]$ irreducible; when $\ell \notin S_f$ this is a theorem (Theorem 1.6).

The set-up. Let $E/\mathbb{Q}$ be any elliptic curve of conductor N, S the set of odd primes at which E has type I_0^* or I_n^*, $D = \prod_{q\in S} q$, $E_S = E\otimes\chi_S$ of conductor N_S, $f = f_E$, $g = f_{E_S}$, and ι the twist map of Theorem 4.17. By Lemma 2.4, valid at every odd q for these exchanges, E_S has good reduction at the I_0^* primes of S and multiplicative reduction, of type I_n, at the I_n^* primes, $n \geq 1$, so $\delta_q = 2$, resp. 1, for g at $q \in S$, $D^2 \mid N$, and $N_S\prod_{q\in S} q^{\delta_q} = N$: the S-depleted form g^S lives at level N, and $\iota(g^S) = f$, since ι kills the coefficients at multiples of D and so does not

distinguish g from g^S; likewise $\iota(f) = g^S$, the coefficients of $f \otimes \chi_S$ being $a_n(g)$ for n prime to D and 0 otherwise. Let $\mathbb{T}$ be the full Hecke algebra of level N, $\pi\colon \mathbb{T} \to \mathbb{Z}$ the eigencharacter of g^S and $\eta^{\mathrm{Hecke}}(g^S) = \pi(\mathrm{Ann}_{\mathbb{T}} \ker \pi)$ its congruence ideal. The generalised eigenspace of π in $S_2(\Gamma_0(N)) \otimes \overline{\mathbb{Q}}$ is the line of g^S: strong multiplicity one identifies the forms with the T_p-eigenvalues of g, $p \nmid N$, with the span of the $g(dz)$, $d \mid N/N_S$; on that span the operators U_q, $q \in S$, have the eigenvalue 0 with multiplicity one, and the U_p with $p \mid N_S$, $p \notin S$, act by the scalars $a_p(g)$. Hence the proof of Lemma 3.1, with $f^\perp$ replaced by the $\mathbb{T}$-stable complement C of the line of g^S (the sum of the generalised eigenspaces of the other eigencharacters; not the Petersson complement, which is not $\mathbb{T}$-stable when the U_p are involved), and carried out over $\mathbb{Z}_\ell$ with the pairing $S_2(\Gamma_0(N);\mathbb{Z}_\ell) \times \mathbb{T}_\ell \to \mathbb{Z}_\ell$, $\mathbb{T}_\ell = \mathbb{T} \otimes \mathbb{Z}_\ell$, shows for every prime ℓ that $\ell^{v_\ell(\eta^{\mathrm{Hecke}}(g^S))}$ is the largest power ℓ^t for which $g^S \equiv h \pmod{\ell^t}$ coefficientwise with $h \in C \cap S_2(\Gamma_0(N);\mathbb{Z}_\ell)$.

Lemma 4.18 (the twist transport)**.** *In this situation, $v_\ell(r_E) \le v_\ell(\eta^{\mathrm{Hecke}}(g^S))$ for every prime ℓ, and*

$$v_\ell(r_E) = v_\ell\big(\eta^{\mathrm{Hecke}}(g^S)\big) \qquad \text{for every prime } \ell \notin S.$$

No other hypothesis on ℓ is needed: $\ell = 2$, $\ell \mid N$, $E[\ell]$ reducible and $\ell \in S_f$ are allowed, and E need not be optimal.

Proof. The inequality $\le$. By Lemma 3.1 there is an integral $h \in f^\perp$ with $f \equiv h \pmod{r_E}$. The form f is new, so $f^\perp$ is the sum of the generalised eigenspaces of the T_p, $p \nmid N$, for systems of eigenvalues that differ from that of f at infinitely many p. The map ι preserves integrality, maps $S_2(\Gamma_0(N))$ into $S_2(\Gamma_0(\mathrm{lcm}(N, D^2))) = S_2(\Gamma_0(N))$, and intertwines T_p with $\chi_S(p)T_p$ for $p \nmid N$; so $\iota(h)$ is integral and is a sum of generalised eigenvectors for systems that differ from that of g at infinitely many p, whence $\iota(h) \in C$. Then $g^S = \iota(f) \equiv \iota(h) \pmod{r_E}$, and $v_\ell(r_E) \le v_\ell(\eta^{\mathrm{Hecke}}(g^S))$ for every ℓ by the characterisation above.

The inequality $\ge$ for $\ell \notin S$. Let $t = v_\ell(\eta^{\mathrm{Hecke}}(g^S))$ and $\mathfrak{m}$ the maximal ideal of $\mathbb{T}_\ell$ containing $\ker \pi$ and ℓ; it contains U_q for $q \in S$. Take $h \in C \cap S_2(\Gamma_0(N);\mathbb{Z}_\ell)$ with $g^S \equiv h \pmod{\ell^t}$. The idempotent $e_{\mathfrak{m}}$ of $\mathbb{T}_\ell$ fixes g^S (as π factors through $\mathbb{T}_{\mathfrak{m}}$) and preserves C and the integral forms, so $h' = e_{\mathfrak{m}} h$ lies in $C \cap S_2(\Gamma_0(N);\mathbb{Z}_\ell)$ and $g^S \equiv h' \pmod{\ell^t}$. The form h' lies in the sum of the generalised eigenspaces of the eigencharacters $\phi \ne \pi$ of $\mathbb{T}$ congruent to π modulo the maximal ideal above ℓ, and such a ϕ has $\phi(U_q) \equiv 0$ for $q \in S$. Every eigenvalue of U_q on $S_2(\Gamma_0(N))$ is 0, ± 1 or a root of $X^2 - a_q(h_0)X + q$ for a newform h_0, hence 0 or a unit at every prime not above q; as $\ell \ne q$, $\phi(U_q) = 0$. (This is the only place where $\ell \notin S$ is used.) Each such ϕ comes from a newform h_0 of level dividing N; if $h_0 = g$, then ϕ agrees with π on the T_p, $p \nmid N$, on the U_q, $q \in S$, and on the U_p, $p \mid N_S$, $p \notin S$, which act by scalars on the span of the $g(dz)$; so $\phi = \pi$, which is excluded. Hence $h_0 \ne g$, and ι maps the generalised eigenspace of ϕ into $S_2(\Gamma_0(N))$, into the generalised eigenspace of the system $\chi_S(p)a_p(h_0)$ of the T_p, $p \nmid N$, which differs from the system $\chi_S(p)a_p(g)$ of f at infinitely many p by strong multiplicity one; so $\iota(h') \in f^\perp$, integral. Then $f = \iota(g^S) \equiv \iota(h') \pmod{\ell^t}$, and the $\mathbb{Z}_\ell$-version of Lemma 3.1 gives $t \le v_\ell(r_E)$. □

At level 539 the four congruence modules of $f_{539\mathrm{c}}$, $f_{539\mathrm{d}}$ and of the depleted forms $(f_{77\mathrm{c}})^{\{7\}}$, $(f_{11\mathrm{a}})^{\{7\}}$ have orders 288, 504, 288 and 504; their 7-parts, $7 \in S$, are not covered by Lemma 4.18.

Proof of Theorem 1.6. Use the set-up above, with $\Sigma = S$ and $\mathcal{O} = \mathbb{Z}_\ell$. Note that $\ell \notin S$, as $\ell \nmid N$; in particular $\ell \ne 3$ when $3 \in S$.

Step 1: the twist map and the full Hecke algebra. By Lemma 4.18, $v_\ell(r_E) = v_\ell(\eta^{\mathrm{Hecke}}(g^S))$; only the inequality

$$v_\ell(r_E) \;\ge\; v_\ell(\eta^{\mathrm{Hecke}}(g^S))$$

is needed below. Let $\mathfrak{m}$ be the maximal ideal of $\mathbb{T}_\ell$ containing $\ker\pi$ and ℓ. The congruence ideal $\eta^{\mathrm{Hecke}}(g^S)$ is computed in $\mathbb{T}_\mathfrak{m}$: the other local factors of $\mathbb{T}_\ell$ lie in $\ker\pi$, and the annihilator decomposes along the idempotents.

Step 2: Hecke versus cohomological congruence ideal. Apply Lemma 4.5 to the module $L = H^1(X_0(N),\mathbb{Z}_\ell)_\mathfrak{m}$, with the pairing of Section 4.1, perfect on the $\mathfrak{m}$-component because $e_\mathfrak{m}$ is self-adjoint, and to π: $L[\wp]$ is the rank-two lattice $M_{g^S,\lambda}$ (Proposition 4.1; it lies in the $\mathfrak{m}$-component), whose Gram determinant is $(\eta^S_{g,\lambda})^2$ up to a unit. Here $\eta^{\mathrm{Hecke}}(g^S) = \pi(\mathrm{Ann}_\mathbb{T}\ker\pi)$ is the Hecke-theoretic congruence ideal, the η of Lemma 4.5 with $d=2$, and $\eta^S_{g,\lambda}$ the cohomological one, generated by the values of the pairing on $M_{g^S,\lambda}$. Hence

$$v_\ell(\eta^{\mathrm{Hecke}}(g^S)) \;\geq\; v_\lambda(\eta^S_{g,\lambda}).$$

This is an inequality: Lemma 4.5 gives equality when L is free over $\mathbb{T}_\mathfrak{m}$, and freeness is not used; when $\ell\notin S_f$ equality follows from the last part of the proof (Remark 4.19(2)).

Step 3: the formula of Diamond, Flach and Guo. By Theorem 4.3 applied to g (its residual representation $\bar\rho\otimes\chi_S$ is irreducible, $\ell\nmid 2N_S$ and $\ell\notin\Sigma=S$),

$$v_\lambda(\eta^S_{g,\lambda}) = v_\lambda(\eta^\emptyset_{g,\lambda}) + \sum_{q\in S} v_\ell\big(L_q^{\mathrm{nv}}(A_g,1)^{-1}\big) = v_\lambda(\eta^\emptyset_{g,\lambda}) + \sum_{q\in S} e_\ell(E,q),$$

the last equality because $A_g = A_f$ and, at $q\in S$, g has $\delta_q\geq 1$, so that its naive factor is the true one, which is the factor that f (with $a_q(f)=0$) omits; the values are those of Lemma 2.5, whose rows I_n^* and I_0^* hold at every odd q, $q=3$ included.

Step 4: the level N_S. By Lemma 4.6 applied to the optimal curve E'_S of the isogeny class of E_S ($\ell\nmid 2N_S$, and $\bar\rho_g$ is irreducible), $v_\lambda(\eta^\emptyset_{g,\lambda}) = v_\ell(\deg\varphi_{E'_S}) = v_\ell(r_{E_S})$, with no freeness hypothesis. Putting the steps together,

$$\begin{aligned} v_\ell(\deg\varphi_E) = v_\ell(r_E) = v_\ell\big(\eta^{\mathrm{Hecke}}(g^S)\big) \;&\geq\; v_\lambda(\eta^S_{g,\lambda}) \\ &= v_\lambda(\eta^\emptyset_{g,\lambda}) + \sum_{q\in S} e_\ell(E,q) = v_\ell(r_{E_S}) + \sum_{q\in S} e_\ell(E,q), \end{aligned}$$

where the first equality is [2, Theorem 2.2] ($\ell\nmid N$), the next three are Steps 1, 3 and 4, and the only inequality is Step 2. This proves the first inequality; the second holds because r_{E_S} is an integer.

The equality. Let $\ell\notin S_f$. Then $\ell\notin S_{f_{E_S}}$, since $\bar\rho_{E_S,\ell} = \bar\rho\otimes\chi_S$ and $N_S\mid N$. Apply Corollary 4.14 to E and to E'_S. The $G_\mathbb{Q}$-modules $\mathrm{ad}^0 T_\ell E$ and $\mathrm{ad}^0 T_\ell E'_S$ are isomorphic. For the twist, $T_\ell E_S \simeq T_\ell E\otimes\chi_S$, and $\mathrm{ad}^0(T\otimes\chi_S) = \mathrm{ad}^0 T$ because $\chi_S(g)$ is a scalar, so that conjugation by $\chi_S(g)\rho(g)$ is conjugation by $\rho(g)$. For the isogenies, $E_S[\ell]\simeq E[\ell]\otimes\chi_S$ is irreducible, so E'_S is linked to E_S by a cyclic isogeny of degree d prime to ℓ (a cyclic isogeny of degree divisible by ℓ would give a $G_\mathbb{Q}$-stable line in $E_S[\ell]$), and such an isogeny induces an isomorphism $T_\ell E_S\simeq T_\ell E'_S$, its composite with the dual isogeny being multiplication by the ℓ-adic unit d. So the Selmer groups coincide, and so do the Tamagawa factors, which are local invariants of $\mathrm{ad}^0 T$. At a prime $q\mid N$ not in S, χ_S is unramified, so E and E_S have the same type (Lemma 2.4(i)), the same naive and true factors, and the same e_ℓ. At $q\in S$, $e_\ell(E_S,q)=0$, E_S having good reduction at a prime of type I_0^* of E and multiplicative reduction of type I_n at a prime of type I_n^*, $n\geq 1$; the Tamagawa factors agree: both are trivial at an I_0^* prime (Lemma 2.9, and E_S is good), and at an I_n^* prime, $n\geq 1$, they coincide because $\mathrm{ad}^0 T$ does. Hence $v_\ell(\deg\varphi_E) - v_\ell(\deg\varphi_{E'_S}) = \sum_{q\in S} e_\ell(E,q)$, and $v_\ell(\deg\varphi_{E'_S}) = v_\ell(r_{E_S})$ as $\ell^2\nmid N_S$. □

Remark 4.19. (1) Theorem 1.6 is the quantitative form of the twist-and-lower argument of Section 3, read in the other direction: the level is raised, from N_S to $N_S\prod q^{\delta_q} = N$, and the

price is the level-raising Euler factor. For $\ell = 3$ it says that a single prime of type I_n^* (necessarily $q \geq 5$), with $3 \mid n$ or not, forces $3 \mid \deg \varphi_E$, and that a prime of type I_0^* with $q \equiv 1 \pmod 3$ does too; neither is in Theorem 1.5, and neither needs the hypothesis on $\mathbb{Q}(\sqrt{-3})$. (2) When $\ell \notin S_f$ every inequality of the proof is an equality, in particular that of Step 2; no multiplicity one at level N or N_S is used. Step 1 is an equality at every prime $\ell \notin S$, with no hypothesis on ℓ at all (Lemma 4.18): the twist map sends f to g^S and $f^\perp$ into the $\mathbb{T}$-stable complement of g^S. (3) Step 1 is where the argument would fail for the potentially good types II, IV, III and their duals: there the quadratic twist does not lower the level, f is its own depletion, and Theorem 4.3 gives no factor. For these types the omitted factor is reached only through Theorem 4.9, that is, with the hypothesis on $\mathbb{Q}(\sqrt{\ell^*})$, and Section 6.7 shows that the hypothesis cannot be dropped there.

5. The boundary of the hypothesis

Proof of Proposition 1.7. Since $\det \bar\rho_{E,3} = \chi_{-3}$ cuts out $\mathbb{Q}(\sqrt{-3})$, the image of $\mathrm{Gal}(\overline{\mathbb{Q}}/\mathbb{Q}(\sqrt{-3}))$ is the subgroup of determinant 1 of the image H of $\bar\rho_{E,3}$. When $E[3]$ is irreducible, H is conjugate to one of $\mathrm{GL}_2(\mathbb{F}_3)$, the normaliser $N_{ns}(\mathbb{F}_3)$ of a nonsplit Cartan subgroup (order 16) and the normaliser $N_s(\mathbb{F}_3)$ of a split Cartan subgroup (order 8); the list is in [51, Theorem 1.2, Propositions 1.14 and 1.16], and we derive it directly: these are the subgroups with surjective determinant acting irreducibly on $\mathbb{F}_3^2$ that contain an element of determinant -1 and trace 0 (complex conjugation), the nonsplit Cartan subgroup itself being excluded by the last condition, and the list follows from these three properties by inspection of the subgroups of $\mathrm{GL}_2(\mathbb{F}_3)$, with or without complex multiplication. Their determinant-one subgroups are $\mathrm{SL}_2(\mathbb{F}_3)$, the quaternion group Q_8 and the cyclic group C_4 of order 4 generated by $J = \left(\begin{smallmatrix} 0 & 1 \\ -1 & 0 \end{smallmatrix}\right)$. The first two act absolutely irreducibly on $\mathbb{F}_3^2$: a subgroup of $\mathrm{GL}_2(\overline{\mathbb{F}}_3)$ fixing a line lies in a Borel subgroup, so its commutator subgroup is abelian of exponent 3, whereas the commutator subgroups $\{\pm 1\}$ of Q_8 and Q_8 of $\mathrm{SL}_2(\mathbb{F}_3)$ are not. The third is abelian, hence absolutely reducible. In the third case J has the characteristic polynomial $X^2 + 1$, irreducible over $\mathbb{F}_3$, so the restriction of $\bar\rho$ to $\mathrm{Gal}(\overline{\mathbb{Q}}/\mathbb{Q}(\sqrt{-3}))$ is irreducible over $\mathbb{F}_3$ and splits as $\psi \oplus \psi^{-1}$ over $\mathbb{F}_9$, for a character ψ of order 4 with values in $\mathbb{F}_9^\times$; by Frobenius reciprocity $\bar\rho \otimes \mathbb{F}_9 \simeq \mathrm{Ind}\, \psi$, so $\bar\rho \simeq \bar\rho \otimes \chi_{-3}$, the isomorphism descending to $\mathbb{F}_3$, and $H^0(\mathbb{Q}, \mathrm{ad}^0\, \bar\rho(1)) \neq 0$ and $h_3(E) = 1$ by Lemma 4.13. Finally the four roots of the 3-division polynomial correspond to the four lines of $\mathbb{F}_3^2$, on which $N_s(\mathbb{F}_3)$ has two orbits of size two (the two lines of the Cartan subgroup and the two others), whereas $N_{ns}(\mathbb{F}_3)$ and $\mathrm{GL}_2(\mathbb{F}_3)$ are transitive; so the division polynomial, which has no rational root when $E[3]$ is irreducible, factors as two quadratics in the first case and is irreducible otherwise. □

The curves with image $N_s(\mathbb{F}_3)$ have a rigid local structure, which accounts for the observations of Section 6.7.

Proposition 5.1. *Let $E/\mathbb{Q}$, with or without complex multiplication, have mod 3 image conjugate to $N_s(\mathbb{F}_3) = \{\pm 1, \pm J, \pm h, \pm w\}$, with $J = \left(\begin{smallmatrix} 0 & 1 \\ -1 & 0 \end{smallmatrix}\right)$, $h = \mathrm{diag}(1, -1)$ and $w = hJ$; let $C_s = \{\pm 1, \pm h\}$ be the split Cartan subgroup, $C_4 = \langle J \rangle = N_s(\mathbb{F}_3) \cap \mathrm{SL}_2(\mathbb{F}_3)$, and let K_1, K_2 be the quadratic fields cut out by $\bar\rho_{E,3}^{-1}(C_s)$ and $\bar\rho_{E,3}^{-1}(\{\pm 1, \pm w\})$, the fields of definition of the roots of the two quadratic factors of the 3-division polynomial ψ_3 (Proposition 1.7). Let R be the set of primes $p \neq 3$ with $\bar\rho_{E,3}(I_p) = C_4$.*

- (i) *For every prime $p \neq 3$, $\bar\rho_{E,3}(I_p)$ lies in C_4. Consequently every prime $p \neq 3$ with $v_p(j_E) < 0$ has $3 \mid v_p(j_E)$, so $t_3(E, p) \geq 1$; no prime $p \neq 3$ has type* IV *or* IV**; and no prime $p \geq 5$ has type* II *or* II*.
- (ii) *R is the set of primes $p \neq 3$ ramified in K_1, and also the set of primes $p \neq 3$ ramified in K_2; it is nonempty. At every $p \in R$ the reduction is potentially good, the image of I_p in $\mathrm{Aut}(T_\ell E)$ is cyclic of order 4 for every prime $\ell \neq p$, $(\mathrm{ad}^0\, V_\ell E)^{I_p}$ is a line on which either*

Frobenius acts by $\chi_{-3}(p)$, $P_p(X) = 1 - \chi_{-3}(p)pX$ *and* $e_3(E,p) = v_3(p - \chi_{-3}(p)) \geq 1$*; in particular* $P_2(X) = 1 + 2X$ *and* $e_3(E,2) = 1$ *if* $2 \in R$.

(iii) *If* $p \in R$ *and* $p \geq 5$*, then* E *has type* III *or* III* *at* p *and* $p \equiv \pm 1 \pmod{12}$*. Conversely every prime* $p \geq 5$ *of type* III *or* III* *lies in* R.

(iv) *The splitting field of* ψ_3 *is biquadratic, with quadratic subfields* K_1, K_2 *and* $\mathbb{Q}(\sqrt{-3})$.

(v) $a_p(E) \equiv 0 \pmod 3$ *for every prime* $p \nmid 3N$ *with* $p \equiv 2 \pmod 3$.

(vi) *If* $3 \nmid N$*, then* E *is ordinary at* 3.

(vii) $h_3(E) = 1$.

Proof. Write $\bar\rho = \bar\rho_{E,3}$. (i) For $p \neq 3$ the determinant χ_{-3} of $\bar\rho$ is unramified at p, so $\bar\rho(I_p) \subseteq N_s(\mathbb{F}_3) \cap \mathrm{SL}_2(\mathbb{F}_3) = C_4$, which has no element of order 3 or 6. If $v_p(j_E) < 0$, then on I_p the representation $\rho_{E,3}$ is $\chi(\sigma)\left(\begin{smallmatrix} 1 & n\theta(\sigma) \\ 0 & 1 \end{smallmatrix}\right)$ in a suitable basis, with χ quadratic, $n = -v_p(j_E)$ and θ the 3-adic tame character (proof of Lemma 2.8); on the kernel of χ, where θ takes a unit value, this is a unipotent of order 3 modulo 3 unless $3 \mid n$. So $3 \mid n$. At a prime $p \neq 3$ of type IV or IV*, $\bar\rho(I_p)$ has order 3 (Lemma 2.2), and at a prime $p \geq 5$ of type II or II*, $\rho(I_p)$ is cyclic of order 6 [23] and injects into $\mathrm{GL}_2(\mathbb{F}_3)$; neither is possible.

We prove (iv) next, because (ii) uses the fields K_1 and K_2 it describes.

(iv) The splitting field of ψ_3 is the field cut out by the kernel $\bar\rho^{-1}(\{\pm 1\})$ of the action on the four lines of $E[3]$, with group $N_s(\mathbb{F}_3)/\{\pm 1\} \simeq (\mathbb{Z}/2)^2$. Its quadratic subfields are the fixed fields of the three subgroups of index two containing -1: C_4, whose fixed field is $\mathbb{Q}(\sqrt{-3})$, the field of the determinant; C_s, the stabiliser of each of the two lines of the Cartan subgroup; and $\{\pm 1, \pm w\}$, the stabiliser of each of the two other lines. The last two give K_1 and K_2.

(ii) Let $p \neq 3$. By (i), $\bar\rho(I_p) \subseteq C_4$, and $C_4 \cap C_s = C_4 \cap \{\pm 1, \pm w\} = \{\pm 1\}$. So $\bar\rho(I_p) \not\subseteq C_s$ if and only if $\bar\rho(I_p) = C_4$ if and only if $\bar\rho(I_p) \not\subseteq \{\pm 1, \pm w\}$; that is, p is ramified in K_1 if and only if $p \in R$ if and only if p is ramified in K_2. As C_s contains h, of determinant -1, $C_s \neq C_4$ and $K_1 \neq \mathbb{Q}(\sqrt{-3})$; every quadratic field other than $\mathbb{Q}(\sqrt{-3})$ is ramified at some prime $p \neq 3$, so $R \neq \emptyset$. (This does not use $3 \nmid N$.) Let $p \in R$. The reduction at p is not potentially multiplicative, since then $\bar\rho(I_p)$ would consist of plus or minus unipotent elements, and C_4 has elements of order 4; so $\rho_{E,3}(I_p)$ is finite, injects into $\mathrm{GL}_2(\mathbb{F}_3)$, and is cyclic of order 4, generated by an element τ with $\tau^2 = -1$ reducing to $J^{\pm 1}$; the kernel of $\rho_{E,\ell}|_{I_p}$ does not depend on $\ell \neq p$ [39, §2], so the image is cyclic of order 4 on every $T_\ell E$. The trace-zero endomorphisms commuting with τ form the line spanned by τ. Frobenius normalises $\langle\tau\rangle$, so it conjugates τ to $\tau^{\pm 1}$, and the sign can be read modulo 3. The elements of $N_s(\mathbb{F}_3)$ not in C_4 form the coset hC_4, of determinant -1, and they conjugate J to J^{-1}, while those of C_4 commute with J; so $\bar\rho(\mathrm{Frob}_p)$ commutes with J if and only if its determinant $\chi_{-3}(p)$ is 1. Hence either Frobenius acts on the line by $\chi_{-3}(p)$, $L_p(A_f, s)^{-1} = 1 - \chi_{-3}(p)p^{-s}$, $P_p(X) = 1 - \chi_{-3}(p)pX$, and, the reduction being additive, $e_3(E,p) = v_3(1 - \chi_{-3}(p)p^{-1}) = v_3(p - \chi_{-3}(p)) \geq 1$, as $p \equiv \chi_{-3}(p) \pmod 3$. At $p = 2$, $\chi_{-3}(2) = -1$.

(iii) For $p \geq 5$ the image is tame, so $\rho(\mathrm{Frob}_p)\tau\rho(\mathrm{Frob}_p)^{-1} = \tau^p$; by (ii) this is $\tau^{\chi_{-3}(p)}$, so $p \equiv \chi_{-3}(p) \pmod 4$, that is, $\chi_{-4}(p) = \chi_{-3}(p)$ and $p \equiv \pm 1 \pmod{12}$. A cyclic image of order 4 means semistability defect 4, that is, $v_p(\Delta_E) \in \{3, 9\}$, the types III, III* [23]. Conversely, at a prime $p \geq 5$ of type III or III* the image $\rho_{E,3}(I_p)$ is cyclic of order 4 [23] and injects into $\mathrm{GL}_2(\mathbb{F}_3)$, so $\bar\rho(I_p)$ is a subgroup of order 4 of C_4, and $p \in R$.

(v) For $p \nmid 3N$ with $p \equiv 2 \pmod 3$, $\det\bar\rho(\mathrm{Frob}_p) = \chi_{-3}(p) = -1$, so $\bar\rho(\mathrm{Frob}_p)$ lies in $hC_4 = \{\pm h, \pm w\}$, whose elements have trace 0; and $a_p(E) \equiv \operatorname{tr}\bar\rho(\mathrm{Frob}_p) \pmod 3$.

(vi) The image $N_s(\mathbb{F}_3)$ acts irreducibly on $\mathbb{F}_3^2$, so $E[3]$ is irreducible, and $3 \in S_f$ by Proposition 1.7; if $3 \nmid N$, the proof of Proposition 4.8 shows that E is ordinary at 3. (vii) is Proposition 1.7. □

The argument of (ii) does not use $3 \nmid N$. Among the 90 optimal curves without complex multiplication with image 3Ns of conductor below 10,000 (44 of them with $3 \mid N$), 74 have a prime $p \geq 5$ in R, of type III or III* and $p \equiv 1$ or 11 (mod 12), and 16 have $R = \{2\}$, where the type is III or III* and $P_2 = 1 + 2X$. The proposition also covers the 13 optimal curves with complex multiplication and image $N_s(\mathbb{F}_3)$ in that range ($j = -32768$ or 8000: 121b1, 256a1, 256d1, 1089e1, 1936f1, 2304c1, 2304k1, 3025a1, 5929a1, 6400f1, 6400r1, 7744a1, 7744s1); 7 of them have $R = \{11\}$ and 6 have $R = \{2\}$, of type III at 2. That the type at 2 is always III or III* when $R = \{2\}$ is not proved.

Remark 5.2. When $\ell = 3$ and the image is $N_s(\mathbb{F}_3)$, Theorem 4.9 does not apply, nor does Theorem 4.10. The exact sequence of Lemma 4.11 still applies if $H^1_f(\mathbb{Q}, V^D) = 0$ ($H^0(\mathbb{Q}, V^D) = 0$ being automatic), and it then reads

$$\operatorname{length} H^1_\Sigma(\mathbb{Q}, W) = \operatorname{length} H^1_f(\mathbb{Q}, W) + \sum_{p \in \Sigma} \big(v_\ell(L_p(A_f, 1)^{-1}) + v_\ell(\operatorname{Tam}_p(A)) \big) - h_\ell(E).$$

The vanishing of $H^1_f(\mathbb{Q}, V^D)$, equivalently of $H^1_f(\mathbb{Q}, \operatorname{ad}^0 V_3 E)$, is not known for these curves: we assume it here. If in addition $v_\lambda(\eta^\Sigma_{f,\lambda})$ were still the length of $H^1_\Sigma(\mathbb{Q}, W)$, as in Theorem 4.9, then (1) would hold with $h_\ell(E)$ subtracted on the right: this is the shape of Conjecture 1.3. Section 6.7 shows that on the tables the correction is exactly what is observed: $h_3(E) = 1$ for every curve with image $N_s(\mathbb{F}_3)$, the inequality (2) fails for 31%, 23% and 18% of them on the three ranges, always by exactly one, and never for any other curve.

6. The computations

6.1. **Data and protocol.** We used Cremona's tables [9] in the form of the `ecdata` repository at commit `25cec5e`, which contain every elliptic curve over $\mathbb{Q}$ of conductor below 500000 with its Kodaira symbols, Tamagawa numbers, modular degree, isogeny class, torsion and the images of the mod ℓ Galois representations in the notation of [44]. We use these labels as they appear in the `galrep` file of the tables, and with each label we give its intrinsic description (for 3Ns, a 3-division polynomial that is a product of two quadratics; for `3Cs.1.1` and `3B.1.1`, a rational point of order 3, Section 6.10); we have not compared them with the tables of [44] themselves. Optimality is determined in the tables for every class of conductor at most 400000 and for 359,009 further classes up to 500000. The ranges are the working set ($N \leq 300000$), the hold-out ($300000 < N \leq 400000$), the extension ($400000 < N \leq 500000$) and the out-of-table set of Section 6.9. The component-group inequality (4) and Conjecture 1.8 were found on the working set, written down, and then tested once on the hold-out, which had been held back, and afterwards on the extension; nothing was adjusted after the hold-out test. The twist-symmetrised form (5) replaced an earlier, weaker formulation of the same phenomenon; it was formulated on the working set, after the counts of (4) on the hold-out and the extension were known, and evaluated on all three ranges. The Euler-factor rule came next: it was formulated on the working set from the minima by Kodaira type (Section 6.5), the terms at 2 and 3 were fixed afterwards by PARI's Euler factors, and the rule was tested on all four ranges; the correction term $h_\ell(E)$ was identified last, from the exceptions. Since the hold-out and the extension had already been used for (4) and (5), they are a confirmation and not a blind test for Conjecture 1.3. The out-of-table curves were generated before the rule was formulated and were evaluated with each version of it; no version failed on them and none was changed because of them. They contain no curve with image 3Ns, so the correction term h_3 was fitted on, and only confirmed by, data already seen; and we do not count the out-of-table curves as a blind test of (5). Nor do we count them as a blind test of the weights w_3, which they could test only through curves with $3 \mid N$ and potentially good reduction at 3. On the extension range optimality is determined only

for the 359,009 classes named above; for a class whose optimal curve is not determined, the right side of (3) at $\ell = 3$ is nevertheless the same for every curve of the class. Indeed, when $E[3]$ is irreducible the class has no 3-isogeny, so every isogeny in it is a composite of isogenies of prime degree $p \neq 3$, and an isogeny of prime degree p preserves the Kodaira type at a prime $q \neq p$ of potentially good reduction [13, Theorem 5.4(1)]; so w_3 is constant on the class. In the presence of a 3-isogeny, where (3) does not apply at $\ell = 3$, w_3 can vary: below conductor 10,000, among the classes with $3 \mid N$, at least two curves and potentially good reduction at 3, w_3 is constant on all 1,295 classes without a 3-isogeny, as it must be, and varies in 828 of the 1,293 classes with one (for instance 54a, whose curves have types IV*, II*, II at 3). The other terms of (3) are invariant under isogenies of degree prime to ℓ. The Kodaira types and discriminant exponents at the bad primes were recomputed with PARI/GP 2.15.4 [28] from the a-invariants, the Euler factors of $\mathrm{Sym}^2 f$ at every bad prime were computed with `lfunsympow` and `lfuneuler`, the Tate parameters with `ellinit` over $\mathbb{Q}_q$ (at $q = 2$ the Tate parameter enters no count, since the extra Tamagawa term vanishes there, $v_\ell(2-1) = 0$, and t_ℓ is read from $v_2(j_E)$; we note that PARI/GP 2.17.2 returns q_E^5 in place of q_E for `ellinit(E, O(2^k)).tate` when $v_2(j_E) = -1$, while the q-expansion of j gives q_E directly), and fifteen modular degrees spanning the configurations were recomputed with `ellmoddegree`, all agreeing with the tables. The mod 3 images of the tables were checked against the factorisation of the 3-division polynomial (Proposition 1.7) on every curve for which it mattered. The checks on conductor below 10,000 (or 30,000 where stated) reported in this note were repeated with PARI/GP 2.17.2 [29]; on that range they reproduce the counts obtained with PARI/GP 2.15.4 on all 38,042 optimal curves, and `ellmoddegree` agrees with the tables on all of them.

6.2. **The rule on whole ranges.** Table 2 checks (3), with PARI's Euler factors at every bad prime $q \neq \ell$, the terms t_ℓ, w_3 as defined, and $h_3(E) = 1$ for the curves with image 3Ns (and $h_\ell = 0$ otherwise, Lemma 4.13), on the four ranges at $\ell = 3, 5, 7, 11, 13$: no violation at any ℓ. It also reports the number of violations when the correction term is dropped (all are curves with image 3Ns, Section 6.7) and the behaviour of the rule with the terms of the types III, III* left out, which is the form that holds for every curve of the tables without any correction; the weights w_3 at $q = \ell = 3$ only sharpen the bound, and at $\ell \geq 5$ nothing is needed at $q = \ell$.

6.3. **Calibration and sharpness.** The zero counts are not tautological. Among the 366,151 working-set curves with $E[3]$ irreducible, not CM, and right-hand side 0 in (3), the prime 3 does not divide $\deg \varphi_E$ for 53.2%. If the valuation of the degree were independent of the local data, the 867,578 working-set curves with a positive bound at $\ell = 3$ would produce roughly 605,000 violations (summing over them the proportion of bound-0 curves whose valuation is smaller than the bound); there are none. This shows that the statement is strong, far from vacuous, not that it is valid; the figure depends on the reference population. The bounds are attained: among working-set curves with a positive bound, $v_\ell(\deg \varphi_E)$ equals the right-hand side of (3) in 57.5% of the cases at $\ell = 3$, 75.6% at $\ell = 5$, 84.1% at $\ell = 7$, 91.6% at $\ell = 11$ and 92.4% at $\ell = 13$, and the mean excess at $\ell = 3$ is 0.63. Under the hypothesis of Theorem 1.2 the identity (1) says what the excess is: the length of the Bloch-Kato Selmer group of the adjoint plus the excess of the Tamagawa factors over the terms t_ℓ (Corollary 4.14). The whole odd part of $\deg \varphi_E$ is another matter: for the 1,226,216 working-set optimal non-CM curves with no isogeny of odd degree, the odd part of $\deg \varphi_E$ equals $\prod_\ell \ell^{b_\ell}$, with b_ℓ the right-hand side of (5), in only 1.85% of cases (the proportion is range dependent: it is 17.1% below conductor 10,000); the quotient is most often 9, 3, 15, 45, 27 or 21, and its smallest prime factor is 3 in 63% of the cases (58.7% below conductor 10,000). Below conductor 10,000 the smallest prime factor of the quotient divides N in 47.5% of the cases with a nontrivial quotient, and some prime factor of the quotient divides N in 51.8%. So the local factors account for a definite, sharp part of the modular degree and for

set	ℓ	curves	(3) > 0	viol.	equality	viol. without h	III-free: > 0 / viol. / equality
working set	3	1,233,729	867,578	0	57.5%	364	861,074 / 0 / 56.8%
	5	1,303,224	332,771	0	75.6%	0	330,477 / 0 / 75.5%
	7	1,307,753	174,266	0	84.1%	0	173,488 / 0 / 84.1%
	11	1,309,103	59,849	0	91.6%	0	59,495 / 0 / 91.6%
	13	1,308,945	35,817	0	92.4%	0	35,635 / 0 / 92.4%
hold-out	3	409,577	298,435	0	54.6%	68	295,973 / 0 / 53.9%
	5	428,251	115,702	0	74.2%	0	114,860 / 0 / 74.0%
	7	429,280	60,868	0	83.2%	0	60,580 / 0 / 83.2%
	11	429,502	21,271	0	90.6%	0	21,097 / 0 / 90.6%
	13	429,498	12,751	0	92.4%	0	12,681 / 0 / 92.3%
extension	3	348,227	257,165	0	54.0%	43	255,507 / 0 / 53.2%
	5	358,244	100,034	0	73.3%	0	99,401 / 0 / 73.2%
	7	358,717	53,034	0	82.7%	0	52,766 / 0 / 82.7%
	11	358,830	19,235	0	90.6%	0	19,136 / 0 / 90.6%
	13	358,816	11,561	0	91.9%	0	11,484 / 0 / 91.9%
out of table	3	893	828	0	52.5%	0	827 / 0 / 51.6%
	5	893	112	0	82.1%	0	111 / 0 / 82.0%
	7	893	52	0	90.4%	0	52 / 0 / 90.4%
	11	893	19	0	94.7%	0	19 / 0 / 94.7%
	13	893	12	0	100.0%	0	12 / 0 / 100.0%

TABLE 2. Conjecture 1.3 with PARI's factors at all primes: curves with $E[\ell]$ irreducible, not CM, with a positive bound in (3); violations; equality rate among them; violations when the correction term h_ℓ is dropped; and the rule without the type III terms ("III-free": positive bounds, violations, equality rate). Rows include primes ℓ dividing N, where no theorem applies; for these the inequality is an observation.

little of the rest, which is governed by the adjoint Selmer group and by the size of the Petersson norm.

6.4. **The Tamagawa factors.** Lemma 2.8 and the identity (1) predict that the bound can be raised, at a prime q of type I_n with $\ell \mid n$ and $q \equiv 1 \pmod{\ell}$, by $\min(v_\ell(n), v_\ell(q-1), v_\ell(c(u)))$, where $v_\ell(c(u)) \geq k$ means that the unit part u of the Tate parameter at q reduces to an ℓ^k-th power in $\mathbb{F}_q^\times$. We computed the Tate parameters with PARI (at $q = 2$ they do not enter, Section 6) for the working-set curves that satisfy the hypothesis of Theorem 1.2 ($\ell \nmid N$, and at $\ell = 3$ image not 3Ns) and have such a prime. At $\ell = 3$ there are 46,222 such curves, 12,095 of them with a positive extra term (12,125 in total); the strengthened inequality has no exception, and the number of curves attaining the bound rises from 20,822 (45.0%) to 28,007 (60.6%). At $\ell = 5$ there are 10,940 such curves, 1,646 with a positive extra term; again no exception, and equality rises from 7,333 (67.0%) to 8,610 (78.7%). The same refinement applies at the primes of type I_n^* (Lemma 2.8); including them, below conductor 10,000 there are 1,050 such curves at $\ell = 3$, with equality rising from 56.8% to 68.0%, and 252 at $\ell = 5$, with equality rising from 203 to 220 curves, and no violation once the CM curves are excluded. This tests the identity, not only the inequality: the extra divisibility of the modular degree predicted by the Tamagawa factor of the adjoint is always present, the Selmer length accounting for the rest.

6.5. **Each term in isolation.** For working-set curves with exactly one additive prime $q \geq 5$, $q \neq \ell$, and no multiplicative prime with $\ell \mid n$, Table 3 gives the minimum of $v_\ell(\deg \varphi_E)$ by Kodaira type and by $e_\ell(E, q)$. In every cell, for $\ell = 3, 5, 7$, the minimum equals e_ℓ and every

curve has $v_\ell(\deg\varphi_E) \geq e_\ell$; for I_n^* the excess over e_ℓ has minimum exactly $v_\ell(n)$ in every cell. The cells with $e = 0$ (III: 5,534 curves, I_0^*: 5,394) have minimum 0, and for I_0^* the term runs up to $e = 6$ (minimum 6). Multiplicative primes show no Euler term: for curves with a single multiplicative prime q with $\ell \mid n$ and no additive prime, $\min(v_\ell(\deg\varphi_E) - v_\ell(n)) = 0$ for every value of $v_\ell(q^2-1)$ up to 6 (at $\ell = 3$ the value 0 occurs only for $q = \ell = 3$, since $3 \mid q^2 - 1$ for every other q). For two primes of potentially good type the terms add: for the 2,502 working-set curves with exactly two odd primes of type II or II* and no other contribution, $9 \mid \deg\varphi_E$ always, and the mechanism is not a congruence between f and its own quadratic twist by the character ramified at both primes, which never occurs (0 of 2,502).

	$e=1$		$e=2$		$e=3$		$e=4$	
type ($\ell=3$)	min	n	min	n	min	n	min	n
II	1	8,327	2	821	3	59	4	2
II*	1	4,184	2	296	3	12	4	1
IV	1	4,290	2	296	3	12	4	1
IV*	1	8,221	2	821	3	59	4	2
III	1	1,271	2	76	3	9	–	–
III*	1	1,271	2	76	3	9	–	–
I_0^*	1	1,370	2	4,432	3	2,737	4	247
I_n^*	1	30,497	2	2,589	3	145	4	13

TABLE 3. Working set, $\ell = 3$: minimum of $v_3(\deg\varphi_E)$ (number of curves) over curves with $E[3]$ irreducible whose only additive prime is a single $q \geq 5$ of the given type with $e_3(E,q) = e$, and no multiplicative 3-contribution. The same holds at $\ell = 5$ and 7. Multiplicative reduction at 3 is not excluded, so the population can contain curves with $3 \mid N$, where no theorem applies and the minimum is an observation.

6.6. **The two corollary forms.** Table 4 gives, for each range and each small prime ℓ, the number of optimal non-CM curves with $E[\ell]$ irreducible and a positive right-hand side in (4) and in (5), the number of violations of each inequality, and the number of Eisenstein curves with a positive bound together with the violations of Conjecture 1.8. There are none. Among working-set curves with a positive bound, equality holds in (5) for 39.4% at $\ell = 3$, 71.1% at $\ell = 5$, 81.2% at $\ell = 7$ and 91.5% at $\ell = 13$, and in (4) for 36.9%, 69.7%, 80.4% and 91.2%; the rule raises these to the rates of Section 6.3 and bounds 867,578 curves instead of 705,563 at $\ell = 3$. The hypothesis of Theorem 1.2 fails on the working set for the 1,170 non-CM curves with image 3Ns at $\ell = 3$ and for the pairs (E,ℓ) with $\ell \mid N$ (among the 913,244 working-set pairs with $E[\ell]$ irreducible, not CM, and a positive multiplicative bound, 175,798 of the 457,537 at $\ell \geq 5$; the 455,707 pairs at $\ell = 3$ were not split by whether $3 \mid N$); by Proposition 4.8 there are no other failures among the non-CM curves with $E[\ell]$ irreducible, the curves whose mod ℓ representation is induced from $\mathbb{Q}(\sqrt{\ell^*})$ with $\ell \geq 5$ all having $\ell \mid N$. (The curves with complex multiplication and $j \in \{-32768, 8000\}$ also have image $N_s(\mathbb{F}_3)$ and fail at $\ell = 3$ when $3 \nmid N$, 20 of them below conductor 10,000; the `galrep` file gives them no label.) None of them violates (3).

6.7. **The boundary: the normaliser of a split Cartan subgroup at $\ell = 3$.** Table 5 records what happens to the inequality (2), with all types included and no correction, on the curves for which the hypothesis of Theorem 1.2 fails at $\ell = 3$. There are 364, 68, 43 and 0 violations on the four ranges, all at $\ell = 3$; every one of them is a curve with image 3Ns in the tables, and the 3-division polynomial of every one of them factors as two quadratics, as Proposition 1.7 requires;

set	ℓ	curves	(4) > 0	(5) > 0	viol. (4)	viol. (5)	Eis. > 0 / viol.
working set, $N \le 300000$	3	1,311,066	598,833	705,563	0	0	67,845 / 0
	5	1,311,066	235,510	262,531	0	0	4,454 / 0
	7	1,311,066	121,947	136,707	0	0	759 / 0
	11	1,311,066	42,334	47,040	0	0	0 / 0
	13	1,311,066	27,215	29,943	0	0	26 / 0
hold-out, $300000 < N \le 400000$	3	429,936	206,749	242,691	0	0	18,225 / 0
	5	429,936	79,550	89,164	0	0	989 / 0
	7	429,936	41,387	46,767	0	0	137 / 0
	11	429,936	14,244	15,970	0	0	0 / 0
	13	429,936	9,212	10,280	0	0	2 / 0
extension, $400000 < N \le 500000$	3	359,009	181,075	214,033	0	0	9,894 / 0
	5	359,009	70,514	78,650	0	0	474 / 0
	7	359,009	36,734	41,317	0	0	70 / 0
	11	359,009	13,347	14,975	0	0	0 / 0
	13	359,009	8,467	9,465	0	0	5 / 0
out of table, $500000 < N \le 4 \cdot 10^6$	3	893	561	718	0	0	0 / 0
	5	893	6	7	0	0	0 / 0
	7	893	0	1	0	0	0 / 0

TABLE 4. Verification counts for the two corollary forms. "(4) > 0" is the number of curves with $E[\ell]$ irreducible, not CM, for which the right-hand side of (4) is positive, and "viol." the number of violations; likewise for (5). The last column counts the Eisenstein curves with a positive bound in (4) and the violations of Conjecture 1.8. Rows include primes ℓ dividing N, where no theorem applies; for these the inequalities are observations.

no curve with image $\mathrm{GL}_2(\mathbb{F}_3)$ or $N_{ns}(\mathbb{F}_3)$ violates (2). (For the curves with $3 \mid N$ the count depends on whether the terms at $q = 3$ are included, that is, the term $t_3(E, 3)$; the weights w_3 change nothing. Below conductor 10,000 there are 36 violations with them, as (2) is written, and 24 without; the twelve curves that differ, 726a1, 726f1, 2178b1, 2178h1, 3840h1, 3840k1, 3840o1, 3840r1, 5070h1, 5070p1, 5808y1 and 5808z1, all have $t_3(E, 3) = 1$ from a type I_n or I_n^* at 3.) Conversely every 3Ns curve of the tables has a prime of type III or III* with a positive term e_3, and the inequality fails for 31%, 23% and 18% of them, always by exactly one; the violation involves a prime $p \equiv -1 \pmod 3$ with $t_3(E, p) \ge 1$ in every case (an observation, not proved). Proposition 5.1 proves most of the rest: every 3Ns curve has a prime $p \ne 3$ with $e_3(E, p) \ge 1$, of type III or III* and $p \equiv \pm 1 \pmod{12}$ when $p \ge 5$, every prime $p \ge 5$ of type III or III* is such a prime, and $t_3(E, p) \ge 1$ at every potentially multiplicative prime $p \ne 3$. For every 3Ns curve $h_3(E) = 1$, and $h_3(E) = 0$ for the other curves, by Lemma 4.13 and Proposition 1.7. So the correction of Conjecture 1.3 is exactly one on exactly these curves, and with it there is no exception. The residue classes of the type III primes of the violating curves are $q \equiv 1$ and $q \equiv 11 \pmod{12}$ in comparable numbers, with some $q = 2$; what the violations have in common is the image, not the prime.

6.8. **The wild primes and the prime ℓ.** Table 6 isolates the additive types at $\ell = 3$ with the weights of (5): for the untwisted types it lists, for working-set curves with no multiplicative prime q with $3 \mid n_q$ and all additive primes of a single Kodaira type, the minimum of $v_3(\deg \varphi_E)$ by type and number of primes; for the twisted types the minimum over the curves whose only contribution to (5) comes from primes of the given type at odd q, or at $q = 2$, with the given total weight. The rows show that the prime 2 contributes nothing through the types II, II* ($v_3(\deg \varphi_E) = 0$ for 14,182 of the 44,672 curves whose only candidate is such a prime) and nothing through a type I_n^* with $3 \,\|\, n$, although with $9 \mid n$ the minimum is 1. Definition 1.1 with

set	3Ns	with III term	failing	others failing	$h_3 = 1$	$q \equiv 1$ / 11 (12) / $q = 2$	shortfall 1
working set	1,170	1,170	364 (31%)	0	1,170 of 1,170	140 / 160 / 72	364 of 364
hold-out	290	290	68 (23%)	0	290 of 290	34 / 30 / 4	68 of 68
extension	237	237	43 (18%)	0	237 of 237	16 / 25 / 2	43 of 43
out of table	0	0	0 (0%)	0	0 of 0	0 / 0 / 0	0 of 0

TABLE 5. The curves with image 3Ns (mod 3 image the normaliser of a split Cartan subgroup), with $E[3]$ irreducible, not CM, optimal: how many have a positive type III term, how many violate (2) (all types, no correction), how many other curves do, how many have $h_3(E) = 1$, the residue classes mod 12 of the type III primes of the violating curves (or $q = 2$), and the shortfall. The hypothesis of Theorem 1.2 fails for every curve with image 3Ns, and the rows include curves with $3 \mid N$; the counts are observations.

PARI's factor explains every one of these minima, the factor having one of the four shapes of Proposition 2.6: at 2 the Kodaira type does not determine it; a type II or II* at 2 usually has $P_2 = 1$ and $e_3 = 0$, but not always (336f1 and 1232l1, Remark 2.7); a type IV or IV* at 2 has $e_3 = 1$ (Lemma 2.2); a type I_n^* at 2 has $t_3 = v_3(-v_2(j))$ when it is potentially multiplicative and 0 otherwise, and the index n of the symbol is not the valuation of the Tate parameter at 2, which is why n was not the right invariant. With the weights of (5) at 2 counted as if 2 were odd, (5) fails thousands of times in each range; with the terms of Definition 1.1 it never does. At $q = 3$ with potentially good reduction and $\ell = 3$, the minima of $v_3(\deg \varphi_E)$ over curves with a single wild additive prime are 1 for $\mathrm{II}, \mathrm{IV}, \mathrm{III}^*$, 2 for $\mathrm{II}^*, \mathrm{IV}^*$ and 0 for $\mathrm{III}, \mathrm{I}_0^*$: these are the weights w_3 of Conjecture 1.3, which lie outside the range of Theorem 1.2 ($\ell \mid N$) and which we cannot derive from an Euler factor. For $\ell = 5, 7$ no term is needed at $q = \ell$ or at $q = 2, 3$ beyond Definition 1.1; the term at 2 given there can be positive (592c1, Remark 2.7). That no term is needed at $q = \ell \geq 5$ is true but not sharp, by the following observation.

Remark 6.1 (a twist relation at $q = \ell$; data only)**.** For each optimal curve E without complex multiplication of conductor below 10,000 with $\ell \mid N$, $\ell \in \{3, 5, 7, 11, 13\}$, compare $v_\ell(\deg \varphi_E)$ with v_ℓ of the degree of the optimal curve of the isogeny class of the twist $E^{(\ell)}$ of E by χ_{ℓ^*}, counting the curves whose twist has conductor below 10,000. (a) If E has type II^*, III^* or IV^* at ℓ, the difference is exactly 1 in all 4,412 cases (2,141, 1,210, 659, 239, 163 at $\ell = 3, 5, 7, 11, 13$; 4,251 of them with $E[\ell]$ irreducible). (b) If E has type I_n^* at ℓ, the difference is 0 in all cases (3,639, 1,360, 677, 240, 173). (c) If E has type I_0^* at ℓ and $E[\ell]$ is irreducible, the difference is $v_\ell\big((\ell+1)^2 - a_\ell(E^{(\ell)})^2\big)$ in all 1,915 cases; it fails in 13 cases with $E[\ell]$ reducible, for instance 126a1 at $\ell = 3$ and 275b1 at $\ell = 5$. At $\ell = 3$, (a) is the pattern of the weights w_3, each starred type weighing one more than its partner under the exchanges of Lemma 2.4(ii). At $\ell \geq 5$ it suggests a weight 1 for the types II^*, III^*, IV^* at $q = \ell$; added to (3), it gives no violation below conductor 10,000 at $\ell = 5, 7, 11, 13$ and is attained by most of these curves. In (b) and (c) the difference is the value at $q = \ell$ of the numerators $v_\ell(q^2 - 1)$ and $v_\ell(q-1) + v_\ell((q+1)^2 - a'^2)$ of the table of Section 1.1. These are observations on the whole range at once, with no hold-out, and we offer no explanation; they bear on the first of the open questions of Section 8.

6.9. **Out of table.** To test the statements beyond the tables, we generated curves with prescribed local types and conductor between 500000 and $4 \cdot 10^6$ (families with a type IV or IV* prime, with two or three type II primes, with mixed types, and random curves), kept those with trivial isogeny class (so that the curve is optimal and $E[\ell]$ is irreducible for every ℓ) and not CM, and computed their modular degrees with `ellmoddegree`. Table 7 summarises them; none has

type	weight 1	weight 2	weight 3
IV (any q)	1 (25,414)	2 (969)	3 (11)
IV* (any q)	1 (30,168)	2 (2,268)	3 (45)
III (any q)	0 (34,904)	0 (2,208)	0 (34)
III* (any q)	0 (25,574)	0 (1,307)	0 (29)
I_0^* (any q)	0 (42,924)	0 (3,093)	0 (53)
II, II* at odd q	1 (56,509)	2 (2,502)	3 (12)
I_n^* at odd q, weight $v_3(n)$	1 (28,175)	2 (3,453)	3 (89)
II, II* at $q = 2$	0 (44,672)	–	–
I_n^* at $q = 2$, weight $v_3(n)$	0 (21,176)	1 (5,329)	1 (342)

TABLE 6. Minimum of $v_3(\deg \varphi_E)$ (number of curves) over working-set curves with $E[3]$ irreducible whose only possible contributions at 3 come from the primes named in the row, with the given number of primes (untwisted rows) or total weight of (5) (twisted rows). The rows "any q" include $q = 3 = \ell$, where no theorem applies; for these curves the minima are observations.

image 3Ns, and Conjecture 1.3 and its two corollary forms hold for all of them (Tables 2 and 4, last rows), with equality in (3) for 52.5% of the 828 curves with a positive bound at $\ell = 3$.

family	curves	largest conductor	median seconds per degree
II,II	98	3,968,800	0.4
II,II,II	1	2,832,200	0.2
IV	334	3,988,012	1.1
IV*	104	3,994,600	4.9
IVmixed	266	3,983,504	1.3
random	90	2,710,716	32.1
all	893	3,994,600	1.5

TABLE 7. Curves outside the tables, with modular degrees computed independently.

6.10. **The Eisenstein case.** Among the 75,430 working-set optimal non-CM curves with a rational 3-isogeny, 67,845 have a positive bound in (4) and 2,777 a positive deficit D; $D = 2$ occurs 22 times. No curve without a rational 3-torsion point has a deficit, whatever its 3-isogeny structure. With a rational point of order 3, mod 3 image of Borel type and no point of order 9, $D \le 1$; $D = 2$ occurs only when $E[3] \simeq \mathbb{Z}/3 \oplus \mu_3$ (the image labelled `3Cs.1.1` in the tables, in the notation of [44]: the subgroup of order 2 of the split Cartan subgroup that fixes a point; seventeen curves, all of rank 0, with torsion $\mathbb{Z}/3$ or $\mathbb{Z}/6$, for instance 14a1, with torsion $\mathbb{Z}/6$, types I_6 at 2 and I_3 at 7, and $\deg \varphi_E = 1$) or when E has a point of order 9 (five curves, with image `3B.1.1`, a Borel image fixing a point of order 3, for instance 714i1, with $\deg \varphi_E = 1080 = 2^3 \cdot 3^3 \cdot 5$, and 6942n1, with $\deg \varphi_E = 65664 = 2^7 \cdot 3^3 \cdot 19$). The two sources of a deficit of 2 never combine: none of the 2,065 curves of conductor at most 400000 with mod 3 image `3Cs.1.1` has a point of order 9 (their torsion is $\mathbb{Z}/3$ or $\mathbb{Z}/6$), and the 18 curves of conductor at most 400000 with a point of order 9, optimal or not, all have image `3B.1.1`. This is a theorem.

Proposition 6.2. *Let $E/\mathbb{Q}$ be without complex multiplication, with $E[3] \simeq \mathbb{Z}/3 \oplus \mu_3$. Then $E(\mathbb{Q})$ has no point of order 9. Consequently, for every non-CM $E/\mathbb{Q}$, $v_3(\#E(\mathbb{Q})_{\mathrm{tors}}) + [E[3] \simeq \mathbb{Z}/3 \oplus \mu_3] \le 2$, and for $\ell \ge 5$, $v_\ell(\#E(\mathbb{Q})_{\mathrm{tors}}) \le 1$.*

Proof. Suppose $P \in E(\mathbb{Q})$ has order 9. Then $3P$ is a rational point of order 3, so $\langle 3P \rangle$ is the line $\mathbb{Z}/3$ of $E[3]$ (μ_3 has no nonzero rational point), and $\langle P \rangle \cap E[3] = \langle 3P \rangle$ meets μ_3 trivially. Let

$\phi\colon E \to E' = E/\mu_3$, rational, and $\hat\phi\colon E' \to E$ its dual. From $\phi\circ\hat\phi = [3]$, $\hat\phi(E'[3]) \subseteq \ker\phi = \mu_3$, with equality as $\#\hat\phi(E'[3]) = 9/\#\ker\hat\phi = 3$. Let $C = \hat\phi^{-1}(\langle P\rangle) \subset E'$, a $G_{\mathbb{Q}}$-stable subgroup of order 27. If C contained $E'[3]$, then $\mu_3 = \hat\phi(E'[3]) \subseteq \langle P\rangle$, a contradiction; so C is cyclic, and $E' \to E'/C$ is a rational cyclic isogeny of degree 27. A rational cyclic 27-isogeny exists only for curves with complex multiplication (by the order of discriminant -27, $j = -2^{15}\cdot 3\cdot 5^3$; PARI's `ellisomat` exhibits the 27-isogeny on that curve) [24]; the level 27 is in the list of the N with $Y_0(N)(\mathbb{Q}) \neq \emptyset$ of [20, Theorem 1]; so E', hence E, has complex multiplication, a contradiction. The consequences follow from Mazur's list of torsion groups [25]: $v_3(\#E(\mathbb{Q})_{\mathrm{tors}}) \leq 2$, with equality only for $\mathbb{Z}/9$, and $v_\ell(\#E(\mathbb{Q})_{\mathrm{tors}}) \leq 1$ for $\ell \geq 5$. □

So the bound $D \leq 2$ of Conjecture 1.8 at $\ell = 3$ follows from its first bound, and is not an additional conjecture. The bounds of Conjecture 1.8 hold with no exception on all three ranges, and the torsion bound is attained with $D > 0$ in 2,773 cases. For $\ell = 5$ and 7 the deficits are at most 1 and occur only with a rational ℓ-torsion point (192 and 18 cases in the working set). Measured against the stronger bound (5) the deficit is not controlled by the torsion: it exceeds the bound of Conjecture 1.8 for 1,148 working-set curves at $\ell = 3$ (189 and 39 on the hold-out and the extension) and for 141 at $\ell = 5$. In the Eisenstein case the additive terms do not contribute reliably, and Conjecture 1.8 is stated for (4) only.

7. The prime 2

At $\ell = 2$ the modular degree is the subject of Watkins's conjecture $2^{\operatorname{rank} E(\mathbb{Q})} \mid \deg\varphi_E$ [48] and of a substantial literature [14, 6, 49, 8, 18, 5], and the mod 2 representation has a rational 2-isogeny for a large fraction of curves. We record only the following. For working-set optimal curves without a rational 2-isogeny and not CM, the inequality $v_2(\deg\varphi_E) \geq \sum_{q\,\|\,N} v_2(n_q) + \#\{q:\ \text{type III, III}^*, \mathrm{I}_0^* \text{ or } \mathrm{I}_n^*\}$ holds with no exception on all three ranges (964,432, 325,510 and 313,741 curves), while counting I_0^* with weight $2 = v_2(\#\Phi_q^{\mathrm{geom}})$ fails (123, 26 and 19 exceptions). The isolated-configuration minima suggest that types III, III* and I_n^* contribute 3 each and types II, IV and I_0^* contribute 1 each, in line with the theorem of Calegari and Emerton [6] (we use its statement and numbering in the arXiv version, where it is Theorem 1.1): a curve of odd modular degree has at most two odd bad primes and even analytic rank, and either it has a rational 2-torsion point, or its conductor is prime, it is supersingular at 2 and $E(\mathbb{R})$ is connected, or it has complex multiplication and $N \in \{27, 32, 49, 243\}$ (for instance 27a1 and 243b1, of degrees 1 and 9, with torsion $\mathbb{Z}/3$). We have not attempted to reconcile the exponents with the results of [14, 8, 18, 5], nor with the Euler factors at $\ell = 2$, which lie outside the range of [12].

8. Open questions

(1) Extend Theorem 1.2 to the primes ℓ dividing N, where the data show the inequality with the weights w_3 at $q = \ell = 3$: identify these weights with the local term at ℓ of the congruence-ideal formula, where the Euler factor is replaced by a condition at ℓ. The twist relation of Remark 6.1, observed at $\ell = 3, 5, 7, 11, 13$, is a lead: it reproduces the pattern of the w_3 and suggests a weight at $q = \ell \geq 5$ as well.

(2) Prove the corrected identity of Remark 5.2 for the curves with image $N_s(\mathbb{F}_3)$, that is, establish $H^1_f(\mathbb{Q}, \mathrm{ad}^0\, V_3E) = 0$ and $v_\lambda(\eta^{\Sigma}_{f,\lambda}) = \operatorname{length} H^1_{\Sigma}(\mathbb{Q}, W)$ without the hypothesis on $\mathbb{Q}(\sqrt{-3})$ in this dihedral case; the data say that the conclusion holds with the correction $h_3(E) = 1$, and the equality of Theorem 1.6 holds in the 8 pairs with image 3Ns below conductor 10,000, which lie outside its equality clause.

(3) The Tamagawa factors of $\mathrm{ad}^0 T$ at the additive primes are known (Lemmas 2.8 and 2.9), so that by Corollary 4.14 the excess of $v_\ell(\deg \varphi_E)$ over the explicit local bound is exactly the length of the adjoint Selmer group (Table 1). Compute that Selmer group independently for the curves exceeding the bound, such as 1184f1 and 1748e1 at $\ell = 3$, so as to account for the excess curve by curve.
(4) Identify the deficit of Conjecture 1.8 with a natural invariant of the Eisenstein ideal at ℓ, presumably through the cuspidal subgroup of $J_0(N)$ [25].
(5) Reconcile Section 7 with the literature on Watkins's conjecture and with the Euler factors at $\ell = 2$.
(6) A smaller point: whether, in Proposition 5.1, the type at 2 is always III or III* when $R = \{2\}$. It is so for all 16 optimal curves without and all 6 with complex multiplication below conductor 10,000 where this happens.

Appendix A. Reproducibility

All scripts, the pinned data commit, the result files and the tables are in the repository `https://github.com/dsr001001/modular_math_book`, directory `final/pipeline`, whose `README.md` lists which script writes each table and how to rerun everything from Cremona's tables. The symmetric-square Euler factors are computed by `scripts/step10_sym2_euler.py`, the rule is checked on the whole ranges by `scripts/step10_rule.py` and `scripts/step11_tw_boundary.py` (correction term, image statistics), the Tamagawa factors by `scripts/step12_tamagawa.py`, the two corollary forms by `scripts/conj_check.py`; each runs on the full tables in minutes to hours. A single curve can be checked against (3) in PARI/GP (version 2.15 or later) with the following functions, which return $[v_\ell(\deg \varphi_E), \mathrm{bound}]$; they are the file `scripts/euler_rule.gp`. The curve must be the optimal curve of its class and $E[\ell]$ must be irreducible: for instance 990h1 returns `[4, 3]` while the optimal curve 990h3 of its class returns `[3, 3]`, and 11a1 at the Eisenstein prime $\ell = 5$ returns `[0, 1]`.

```
rule(E, l, {iii = 0}) =
{ my(R = ellglobalred(E), P = R[4][,1], K = R[5], S = lfunsympow(E, 2), b = 0, j = E.j);
  for(i = 1, #P,
    my(q = P[i], k = K[i][2], e = 0, t = 0, vj = valuation(j, q));
    if(q != l,
      my(F = 1/lfuneuler(S, q), r = subst(F, x, 1/q^2)); \\ P_q(q^{-2}): omitted factor
      if(k >= 5, r /= (1 - 1/q^2));                      \\ multiplicative: naive one out
      if(abs(k) != 3 || iii, e = valuation(r, l)),       \\ III, III* (codes 3,-3) if iii
      if(l == 3, e = if(k == 4 || k == 2, 1,
                     if(k == -4 || k == -2, 2, if(k == -3, 1, 0))))));   \\ w_3 at q = 3
    if(vj < 0, t = valuation(-vj, l));                    \\ Tamagawa exponent
    b += e + t);
  if(iii && l == 3, b -= ns3(E));                         \\ the correction term h_3
  [valuation(ellmoddegree(E), l), b]; }
ns3(E) = my(F = factor(elldivpol(E, 3))); (#F[,1] == 2 && poldegree(F[1,1]) == 2);
\\ psi_3 has degree 4 and is squarefree, so two factors, one quadratic, means two quadratics
```

Here `K[i][2]` is PARI's Kodaira code: $n+4$ for I_n and $-(n+4)$ for I_n^* when $n \geq 1$, 1 for I_0 and -1 for I_0^*, ± 2 for II/II*, ± 3 for III/III*, ± 4 for IV/IV*. The function `rule(E,l)` leaves the type III terms out and `rule(E,l,1)` is (3); at $q = \ell$ the term t_ℓ is that of Definition 1.1, which is defined there. For example `rule(ellinit([1,1,1,-30,-76]), 3)` (the curve 121a1) returns `[1, 1]`; for 726a1, a curve with image 3Ns and type III at 11, `rule(E,3)` and `rule(E,3,1)` both return `[2, 2]`, the type III term and the correction cancelling. The function `ellmoddegree` needs about 32 MB of stack with PARI 2.17 for conductors in the thousands (with PARI 2.15 and larger conductors, `parisizemax` of a few hundred megabytes). The whole pipeline was rerun from the pinned data with PARI/GP 2.17.2 and reproduced every committed result summary and Tables 2, 3, 4, 5, 6 and 7 byte for byte. Lemma 2.4(ii) was also checked numerically, with the

script `twist_lemma_check.gp` of the repository, on 4,577 random pairs (curve, odd bad prime), with no exception, and its exchanges at $q = 3$ on all 5,122 pairs (curve, twist by $\mathbb{Q}(\sqrt{-3})$) below conductor 10,000, and Lemma 2.5 against `lfunsympow` on every curve of the tables.

Statement on the use of an AI system

This work was carried out with an AI system (Claude, Anthropic) at every stage, and it would not have been done without it. The author set the goal, a search for new relations in Cremona's tables, chose the data and the protocol (a frozen snapshot of the tables, a working set, a range held back for one-time tests, and a list of tests fixed before any was run), and decided at each step what to pursue and what to keep. Under that direction, and in discussion with the author, the AI system did the rest: it built the feature table and the list of known relations to be excluded; ran the pre-registered search, which produced the candidate relation between the modular degree and the local invariants; searched the literature and identified the results of Ribet-Takahashi, Pollack-Weston, Kim-Ota, Agashe-Ribet-Stein and Diamond-Flach-Guo behind the statements; replaced the Tamagawa number by the geometric component group and formulated the conjectures and their successive refinements, up to the Euler-factor rule and its correction term; wrote and ran the verification code in Python and PARI/GP on the four ranges, generated the out-of-table curves and computed their modular degrees and Tate parameters; read the published text of [12] and assembled the proof of Theorem 1.2 from its statements, computed the Tamagawa factor of Lemma 2.8, and proposed and wrote the proofs of Theorems 1.5 and 1.6 and of Proposition 1.7; drafted the text and the tables; and prepared the responses to reviews of drafts, two of them written by AI systems, from which the identification of the local factors and of the boundary of the hypothesis arose. Several of the results and proofs (among them Propositions 4.4, 4.8, 2.6, 5.1 and 6.2, Lemmas 2.9 and 4.18, Corollary 4.14, the equality in Theorem 1.6 and the exchanges at $q = 3$ in Lemma 2.4), and the checks of the counts below conductor 10,000, were worked out with the same AI system while writing a companion textbook on this note. The note was prepared with an AI system under the author's direction and has not yet been refereed; the author has to review it before it is circulated. The scripts, the result files and the pinned data commit are in the repository named in the appendix, so that every count can be rerun; the proofs cite the published statements they use, with their numbering. The author takes full responsibility for all statements, proofs, computations and errors.

References

1. Ahmed Abbes and Emmanuel Ullmo, *à propos de la conjecture de Manin pour les courbes elliptiques modulaires*, Compositio Math. **103** (1996), no. 3, 269–286.
2. Amod Agashe, Kenneth A. Ribet, and William A. Stein, *The modular degree, congruence primes, and multiplicity one*, Number Theory, Analysis and Geometry: In Memory of Serge Lang (Dorian Goldfeld, Jay Jorgenson, Peter Jones, Dinakar Ramakrishnan, Kenneth A. Ribet, and John Tate, eds.), Springer, New York, 2012, Section and theorem numbers as in the authors' version at `https://www.williamstein.org/papers/ars-congruence/`, pp. 19–49.
3. A. O. L. Atkin and J. Lehner, *Hecke operators on* $\Gamma_0(m)$, Math. Ann. **185** (1970), 134–160.
4. A. O. L. Atkin and Wen Ch'ing Winnie Li, *Twists of newforms and pseudo-eigenvalues of* W*-operators*, Invent. Math. **48** (1978), no. 3, 221–243.
5. Subham Bhakta, Srilakshmi Krishnamoorthy, and Sunil Kumar Pasupulati, *Modular degree and a conjecture of Watkins*, 2023, Preprint, arXiv:2308.13708.
6. Frank Calegari and Matthew Emerton, *Elliptic curves of odd modular degree*, Israel J. Math. **169** (2009), no. 1, 417–444, arXiv:math/0503359.
7. Henri Carayol, *Sur les représentations galoisiennes modulo* l *attachées aux formes modulaires*, Duke Math. J. **59** (1989), no. 3, 785–801.

8. Jerson Caro and Hector Pasten, *Watkins's conjecture for elliptic curves with non-split multiplicative reduction*, Proc. Amer. Math. Soc. **150** (2022), no. 8, 3245–3251.
9. J. E. Cremona, *Algorithms for modular elliptic curves*, second ed., Cambridge University Press, Cambridge, 1997.
10. Henri Darmon, Fred Diamond, and Richard Taylor, *Fermat's last theorem*, Elliptic Curves, Modular Forms & Fermat's Last Theorem (Hong Kong, 1993), Int. Press, Cambridge, MA, 1997, pp. 2–140.
11. Fred Diamond, *The refined conjecture of Serre*, Elliptic Curves, Modular Forms, & Fermat's Last Theorem (Hong Kong, 1993), Ser. Number Theory, I, Int. Press, Cambridge, MA, 1995, pp. 22–37.
12. Fred Diamond, Matthias Flach, and Li Guo, *The Tamagawa number conjecture of adjoint motives of modular forms*, Ann. Sci. École Norm. Sup. (4) **37** (2004), no. 5, 663–727, Numbering as in the published version; the version posted as arXiv:2512.02348 (2025) is numbered differently.
13. Tim Dokchitser and Vladimir Dokchitser, *Local invariants of isogenous elliptic curves*, Trans. Amer. Math. Soc. **367** (2015), no. 6, 4339–4358, arXiv:1208.5519.
14. Neil Dummigan, *On a conjecture of Watkins*, J. Théor. Nombres Bordeaux **18** (2006), no. 2, 345–355.
15. Bas Edixhoven, *The weight in Serre's conjectures on modular forms*, Invent. Math. **109** (1992), no. 3, 563–594.
16. Matthias Flach, *A finiteness theorem for the symmetric square of an elliptic curve*, Invent. Math. **109** (1992), no. 2, 307–327.
17. Jean-Marc Fontaine and Bernadette Perrin-Riou, *Autour des conjectures de Bloch et Kato: cohomologie galoisienne et valeurs de fonctions L*, Motives (Seattle, WA, 1991), Proc. Sympos. Pure Math., vol. 55, Part 1, Amer. Math. Soc., Providence, RI, 1994, pp. 599–706.
18. Jeffrey Hatley and Debanjana Kundu, *Elliptic curves of conductor $2^m p$, quadratic twists, and Watkins' conjecture*, Ann. Math. Qué. **50** (2026), 143–165, arXiv:2411.08321.
19. Haruzo Hida, *Congruences of cusp forms and special values of their zeta functions*, Invent. Math. **63** (1981), no. 2, 225–261.
20. M. A. Kenku, *On the number of $\mathbf{Q}$-isomorphism classes of elliptic curves in each $\mathbf{Q}$-isogeny class*, J. Number Theory **15** (1982), no. 2, 199–202.
21. Chan-Ho Kim and Kazuto Ota, *On the quantitative variation of congruence ideals and integral periods of modular forms*, Res. Math. Sci. **10** (2023), no. 2, Paper No. 22, arXiv:1905.02926; theorem and definition numbers as in v4.
22. Alain Kraus, *Quelques remarques à propos des invariants c_4, c_6 et Δ d'une courbe elliptique*, Acta Arith. **54** (1989), no. 1, 75–80.
23. ———, *Sur le défaut de semi-stabilité des courbes elliptiques à réduction additive*, Manuscripta Math. **69** (1990), 353–385.
24. Gérard Ligozat, *Courbes modulaires de genre 1*, Bull. Soc. Math. France, Mém., vol. 43, Soc. Math. France, Paris, 1975.
25. B. Mazur, *Modular curves and the Eisenstein ideal*, Inst. Hautes Études Sci. Publ. Math. **47** (1977), 33–186.
26. A. P. Ogg, *Elliptic curves and wild ramification*, Amer. J. Math. **89** (1967), no. 1, 1–21.
27. Ioannis Papadopoulos, *Sur la classification de Néron des courbes elliptiques en caractéristique résiduelle 2 et 3*, J. Number Theory **44** (1993), no. 2, 119–152.
28. The PARI Group, Univ. Bordeaux, *Pari/gp version 2.15.4*, 2023, available from `http://pari.math.u-bordeaux.fr/`.
29. The PARI Group, Univ. Bordeaux, *Pari/gp version 2.17.2*, 2025, available from `https://pari.math.u-bordeaux.fr/`.
30. Hector Pasten, *Shimura curves and the abc conjecture*, J. Number Theory **254** (2024), 214–335, With an appendix by Robert Lemke Oliver and Jesse Thorner; arXiv:1705.09251.
31. Robert Pollack and Tom Weston, *On anticyclotomic μ-invariants of modular forms*, Compos. Math. **147** (2011), no. 5, 1353–1381, arXiv:math/0610694.
32. R. A. Rankin, *Contributions to the theory of Ramanujan's function $\tau(n)$ and similar arithmetical functions. I. The zeros of the function $\sum_{n=1}^{\infty} \tau(n)/n^s$ on the line $\operatorname{Re} s = 13/2$. II. The order of the Fourier coefficients of integral modular forms*, Proc. Cambridge Philos. Soc. **35** (1939), no. 3, 351–356, 357–372.
33. K. A. Ribet, *On modular representations of $\operatorname{Gal}(\overline{\mathbf{Q}}/\mathbf{Q})$ arising from modular forms*, Invent. Math. **100** (1990), no. 2, 431–476.
34. Kenneth A. Ribet, *Mod p Hecke operators and congruences between modular forms*, Invent. Math. **71** (1983), no. 1, 193–205.
35. Kenneth A. Ribet and William A. Stein, *Lectures on Serre's conjectures*, Arithmetic Algebraic Geometry (Park City, UT, 1999), IAS/Park City Math. Ser., vol. 9, Amer. Math. Soc., Providence, RI, 2001, pp. 143–232.

36. Kenneth A. Ribet and Shuzo Takahashi, *Parametrizations of elliptic curves by Shimura curves and by classical modular curves*, Proc. Nat. Acad. Sci. U.S.A. **94** (1997), no. 21, 11110–11114.
37. Takeshi Saito, *Conductor, discriminant, and the Noether formula of arithmetic surfaces*, Duke Math. J. **57** (1988), no. 1, 151–173.
38. Jean-Pierre Serre, *Propriétés galoisiennes des points d'ordre fini des courbes elliptiques*, Invent. Math. **15** (1972), no. 4, 259–331.
39. Jean-Pierre Serre and John Tate, *Good reduction of abelian varieties*, Ann. of Math. (2) **88** (1968), no. 3, 492–517.
40. Goro Shimura, *On the holomorphy of certain Dirichlet series*, Proc. London Math. Soc. (3) **31** (1975), no. 1, 79–98.
41. ———, *The special values of the zeta functions associated with cusp forms*, Comm. Pure Appl. Math. **29** (1976), no. 6, 783–804.
42. Joseph H. Silverman, *Advanced topics in the arithmetic of elliptic curves*, Graduate Texts in Mathematics, vol. 151, Springer-Verlag, New York, 1994.
43. Jacob Sturm, *On the congruence of modular forms*, Number Theory (New York, 1984-1985), Lecture Notes in Math., vol. 1240, Springer, Berlin, 1987, pp. 275–280.
44. Andrew V. Sutherland, *Computing images of Galois representations attached to elliptic curves*, Forum Math. Sigma **4** (2016), Paper No. e4, 79 pp.
45. Shuzo Takahashi, *Degrees of parametrizations of elliptic curves by Shimura curves*, J. Number Theory **90** (2001), 74–88.
46. J. Tate, *Algorithm for determining the type of a singular fiber in an elliptic pencil*, Modular Functions of One Variable, IV (Antwerp, 1972), Lecture Notes in Math., vol. 476, Springer, Berlin, 1975, pp. 33–52.
47. Jacques Tilouine, *Hecke algebras and the Gorenstein property*, Modular Forms and Fermat's Last Theorem (Boston, MA, 1995), Springer, New York, 1997, pp. 327–342.
48. Mark Watkins, *Computing the modular degree of an elliptic curve*, Experiment. Math. **11** (2002), no. 4, 487–502.
49. Soroosh Yazdani, *Modular abelian varieties of odd modular degree*, Algebra Number Theory **5** (2011), no. 1, 37–62.
50. Don Zagier, *Modular parametrizations of elliptic curves*, Canad. Math. Bull. **28** (1985), no. 3, 372–384.
51. David Zywina, *On the possible images of the mod ℓ representations associated to elliptic curves over $\mathbb{Q}$*, 2015, Preprint, arXiv:1508.07660.

Email address: dharmaj.soni@gmail.com
URL: https://orcid.org/0009-0007-6397-3351