

Determining experimental plasma wave dispersion relations in the solar wind

David Stansby, Tim Horbury, Christopher Chen and Lorenzo Matteini
Space and Atmospheric Physics Group, Imperial College London
david.stansby14@imperial.ac.uk

Stansby et. al. The Astrophysical Journal, 829(1), L16. <http://doi.org/10.3847/2041-8205/829/1/L16>

Introduction

- The solar wind hosts short and intense coherent magnetic field fluctuations at multiple scales up to 10% of the time [1, 2]:
 - what type of waves are they?
 - under what conditions do they form?
- Solar Orbiter and Solar Probe Plus will be able to answer these questions for the first time in the inner heliosphere
- Here we look at waves between ion and electron scales at 1AU as a case study

[1] Lacombe et. al. (2014) ApJ doi:10.1088/0004-637X/796/1/5
[2] Jian et. al. (2014) ApJ doi:10.1088/0004-637X/786/2/123


Measuring dispersion relations

- To experimentally derive a dispersion relation, need to measure frequency (ω) and wave-vector (k)
- ω directly from \underline{B} waveform
- \hat{k} from minimum variance on \underline{B} waveform and direction of $\delta \underline{E} \times \delta \underline{B}$
- $|k|$ from $\frac{\omega}{|k|} = \frac{|\delta \underline{E}_\perp|}{|\delta \underline{B}|}$
- This method works with elliptically polarised monochromatic waves (eg. ion cyclotron, whistler waves)

With both \underline{E} and \underline{B} waveforms, possible to calculate plasma frame frequencies, wavenumbers, and an experimental **dispersion relation**.

Electric field processing


- Using electric field measurements taken in the solar wind is challenging due to spacecraft wake effects and spin tones
- A new method has been developed to remove these effects from Electric Field Instrument measurements on ARTEMIS




- This is used to extract high frequency (128 samples/s) 2D spin plane electric field waveforms

Wave packet observations


- 2.5 years of ARTEMIS data in solar wind (2009 - mid 2011)
- Burst mode data at 128 samples/s with strong magnetic fluctuation power in the range [4, 64]Hz selected
- Automatically detect wave packets



- 289 wave packets with reliable data detected over three separate days
- > 95% are **RH circularly polarised**




Travel predominantly **anti-sunward**



Travel **along B₀**

Experimental dispersion relation



- Agreement with warm dispersion relation and RH polarisation
 - These are electron whistler waves
- Systematic variability in dispersion relation with local $\beta_{\parallel e}$
 - Agrees qualitatively with bi-Maxwellian dispersion relations

Conclusions

- Developed a new method to isolate high frequency component of solar wind electric field data
- For the first time have experimentally measured dispersion relation of whistler waves at electron scales in the solar wind
- Warm plasma effects play a significant role in whistler wave dispersion
- These whistler waves are travelling anti-sunward → **cannot** undergo linear wave-particle interaction to scatter electron strahl

Further work

- Combine experimental dispersion relations with theoretical predictions derived from in-situ particle measurements
- Apply method to fluctuations near ion scales

Acknowledgements

D. Stansby is supported by STFC studentship ST/N504336/1.
T. S. Horbury and L. Matteini are supported by STFC grant ST/N000692/1.
C. H. K. Chen. is supported by a STFC Ernest Rutherford Fellowship.