

Math 128A, Summer 2019

Midterm Study Guide

1 Topics

conditions for convergence (fixed point iteration) - contractive - invariance
 added conditions for quadratic convergence ; other big-O practice
 - bounding error of a step of fixed point iteration by max of derivative on that interval (examples!!)
 iee delta/epsilon error bounds
 [fixed point, iteration, big-O, floating point]

2 Key Results

2.1 Convergence of Fixed Point Iteration

Theorem 2.1. If $a \leq x \leq b$ implies:

1. $a \leq g(x) \leq b$ [Invariant]
2. $|g'(x)| \leq \frac{1}{2}$ [Contractive],

then for any $x_n \in [a, b]$, we have:

$$x_{n+1} = g(x_n); x = g(x) \implies |x_n - x| \leq 2^{-n} |x_0 - x|$$

Moreover, this gives a **unique** solution.

2.2 Rounding in Floating Point

Remark: Rounding is “monotonic”; that is, it preserves order (inequalities), with possible equality.

$$a < b \implies fl(a) \leq fl(b)$$

The ‘Mantra’ of FP arithmetic is it delivers the exact result, correctly rounded (for binary operations). Note that we write $|\delta| < \frac{\epsilon}{2}$ as relative error, completely independent of the dimensions of x .

2.3 Quadratic Convergence: Fixed Point

Theorem 2.2. If $x \in [a, b]$ implies:

1. $g(x) \in [a, b]$
2. $|g'(x)| \leq \frac{1}{2}$
3. $|g''(x)| < C; C|b - a| \leq 1$

If $p \in [a, b]$ has the property $|g''(x)| < M$ for $x \in [a, b]$ and $g'(p) = 0$ (derivative is zero at 2nd derivative), we have convergence at least order of 2 (quadratic).

For derivation of the error bound here, look at the equations in this text:

$$|p_{n+1} - p| < \frac{M}{2}|p_n - p|^2.$$

$$g(x) = g(p) + g'(p)(x - p) + \frac{g''(\xi)}{2}(x - p)^2,$$

then x and p . The hypotheses $g(p) = p$ and $g'(p) =$

$$g(x) = p + \frac{g''(\xi)}{2}(x - p)^2.$$

if $x = p_n$,

$$p_{n+1} = g(p_n) = p + \frac{g''(\xi_n)}{2}(p_n - p)^2,$$

ξ_n and p . Thus,

$$p_{n+1} - p = \frac{g''(\xi_n)}{2}(p_n - p)^2.$$

< 1 on $[p - \delta, p + \delta]$ and g maps $[p - \delta, p + \delta]$ into itself. By the Contraction Mapping Theorem that $\{p_n\}_{n=0}^\infty$ converges to p . But ξ_n is between p_n and p , so ξ_n also converges to p , and

$$\lim_{n \rightarrow \infty} \frac{|p_{n+1} - p|}{|p_n - p|^2} = \frac{|g''(p)|}{2}.$$

1, or duplicated, in whole or in part. Due to electronic rights, some third party content may be suppressed from the eBook and/or eChapter(s). Editorial review has determined that any suppressed content does not materially affect the overall learning experience. Cengage Learning reserves the right to remove additional content at any time if subsequent rights restrictions require it.

2.4 Error Analysis for Iteration

shows that the sequence $\{p_n\}_{n=0}^\infty$ is quadratically convergent to p if $g''(p) \neq 0$.

g'' is continuous and strictly bounded by M on the interval $[p - \delta, p + \delta]$. That is, for sufficiently large values of n ,

$$|p_{n+1} - p| < \frac{M}{2}|p_n - p|^2.$$

3 Lagrange Interpolation and Error

Suppose x_0, x_1, \dots, x_n are distinct numbers in the interval $[a, b]$ and $f \in C^{n+1}[a, b]$. Then, for each x in $[a, b]$, a number $\xi(x)$ (generally unknown) between $\min\{x_0, x_1, \dots, x_n\}$, and the $\max\{x_0, x_1, \dots, x_n\}$ and hence in (a, b) , exists with

$$f(x) = P(x) + \frac{f^{(n+1)}(\xi(x))}{(n+1)!} (x - x_0)(x - x_1) \cdots (x - x_n), \quad (3.3)$$

where $P(x)$ is the interpolating polynomial given in Eq. (3.1).

4 Hermite Interpolation

If $f \in C^1[a, b]$ and $x_0, \dots, x_n \in [a, b]$ are distinct, the unique polynomial of least degree agreeing with f and f' at x_0, \dots, x_n is the Hermite polynomial of degree at most $2n + 1$ given by

$$H_{2n+1}(x) = \sum_{j=0}^n f(x_j) H_{n,j}(x) + \sum_{j=0}^n f'(x_j) \hat{H}_{n,j}(x),$$

where, for $L_{n,j}(x)$ denoting the j th Lagrange coefficient polynomial of degree n , we have

$$H_{n,j}(x) = [1 - 2(x - x_j)L'_{n,j}(x_j)]L_{n,j}^2(x) \quad \text{and} \quad \hat{H}_{n,j}(x) = (x - x_j)L_{n,j}^2(x).$$

Moreover, if $f \in C^{2n+2}[a, b]$, then

$$f(x) = H_{2n+1}(x) + \frac{(x - x_0)^2 \cdots (x - x_n)^2}{(2n+2)!} f^{(2n+2)}(\xi(x)),$$

for some (generally unknown) $\xi(x)$ in the interval (a, b) . ■

z	$f(z)$	First divided differences	Second divided differences
$z_0 = x_0$	$f[z_0] = f(x_0)$		
		$f[z_0, z_1] = f'(x_0)$	
$z_1 = x_0$	$f[z_1] = f(x_0)$		$f[z_0, z_1, z_2] = \frac{f[z_1, z_2] - f[z_0, z_1]}{z_2 - z_0}$
		$f[z_1, z_2] = \frac{f[z_2] - f[z_1]}{z_2 - z_1}$	
$z_2 = x_1$	$f[z_2] = f(x_1)$		$f[z_1, z_2, z_3] = \frac{f[z_2, z_3] - f[z_1, z_2]}{z_3 - z_1}$
		$f[z_2, z_3] = f'(x_1)$	
$z_3 = x_1$	$f[z_3] = f(x_1)$		$f[z_2, z_3, z_4] = \frac{f[z_3, z_4] - f[z_2, z_3]}{z_4 - z_2}$
		$f[z_3, z_4] = \frac{f[z_4] - f[z_3]}{z_4 - z_3}$	
$z_4 = x_2$	$f[z_4] = f(x_2)$		$f[z_3, z_4, z_5] = \frac{f[z_4, z_5] - f[z_3, z_4]}{z_5 - z_3}$
		$f[z_4, z_5] = f'(x_2)$	
$z_5 = x_2$	$f[z_5] = f(x_2)$		

5 Other Practice Exams

5.1 Hare F2003

: **Problem 1.** (a) Let $f(x)$ be a continuous function on $[2, 3]$. Further, assume that for all $x \in [2, 3]$ that $f(x) \in [2, 3]$. Prove that $f(x)$ has a fixed point $p \in [2, 3]$. (b) Further assume that $|f'(x)| \leq \frac{1}{11}$ for all $x \in [2, 3]$. How many steps of fixed point iteration are needed to guarantee accuracy of within 10^{-3} ?

Write $g(x) := f(x) - x$. Because $f(2) \in [2, 3]$, we have

$$g(2) = f(2) - 2 \geq 0.$$

Further, because $f(3) \in [2, 3]$, we have

$$g(3) = f(3) - 3 \leq 0.$$

Hence by the IVT, $g(x)$ has a root $p \in [2, 3]$. So $0 = g(p) = f(p) - p$, which gives us $f(p) = p$ as desired as a fixed point.

Let $p_0 \in [2, 3]$ be our initial guess and let p be the true unique fixed point. Then we know, for all n :

$$|p_n - p| < \left(\frac{1}{11}\right)^n |p_0 - p| < \left(\frac{1}{11}\right)^n$$

So to ensure precision within 10^{-3} , $n \geq 3$ should suffice.

Remark: The right inequality follows from $|p_0 - p| < 3 - 2$, but why do we have the left inequality?

Give an example of a, b, c where three-digit rounding fails associativity; that is,

$$a + (b + c) \neq (a + b) + c$$

Solution. Define $a := 1.00, b := 0.004, c := 0.004$. □

Let $g(x) := ax + b$ for some fixed nonzero a, b . How many steps of Newton's method are needed to find a root of $g(x)$ to within 10^{-6} ?

Solution. One step guarantees the exact solution. Intuitively, Newton's method finds the intercept (root) of the tangent line of a function. Because our function itself is a line, for any starting point, we find the actual root to the line $g(x)$.

To see this precisely, consider:

$$\begin{aligned} x - \frac{g(x)}{g'(x)} &= x - \frac{ax + b}{a} \\ &= x - x - \frac{b}{a} \\ &= -\frac{b}{a}, \end{aligned}$$

which is the exact solution (root). □

Define what it means for a sequence $\{p_n\}_{n=0}^{\infty}$ to converge **linearly** to p .

Solution. We say the sequence p_n converges linearly to p if there exists some $\lambda \in (0, 1)$ such that:

$$\lim_{n \rightarrow \infty} \frac{|p_{n+1} - p|}{|p_n - p|^1} = \lambda.$$

□