

TECHNICAL UNIVERSITY OF DENMARK

CONSTRAINED OPTIMIZATION

COURSE 02612

Assignment 1

Authors:

Jakub CZERNY, s99999

Oskar HINT, s161559

Joachim Finn JENSEN, s134052

March 14, 2017



Contents

1	Introduction	1
2	Assignment	1
2.1	Problem 1 - Quadratic Optimization	1
2.2	Problem 2 - Equality Constrained Quadratic Optimization	2
2.3	Problem 3 - Inequality Constrained Quadratic Programming	2
2.4	Problem 4 - Markowitz Portfolio Optimization	4
2.5	Problem 5 - Interior-Point Algorithm for Convex Quadratic Programming . .	4
3	Conclusion	4

1 Introduction

this is some report

2 Assignment

2.1 Problem 1 - Quadratic Optimization

From the problem 1 it is seen that the system can be written as:

$$H = \begin{bmatrix} 6 & 2 & 1 \\ 2 & 5 & 2 \\ 1 & 2 & 4 \end{bmatrix}$$

$$g = \begin{bmatrix} -8 \\ -3 \\ -3 \end{bmatrix}$$

$$A = \begin{bmatrix} 1 & 0 \\ 0 & 1 \\ 1 & 1 \end{bmatrix}$$

$$b = \begin{bmatrix} 3 \\ 0 \end{bmatrix}$$

Then the general KKT system can be written as:

$$\begin{bmatrix} H & -A \\ A^T & 0 \end{bmatrix} \times \begin{bmatrix} x \\ \lambda \end{bmatrix} = \begin{bmatrix} -g \\ b \end{bmatrix}$$

The system is solve with a function in MatLab, see the code in ?X?X. The result shows that:

$$x = \begin{bmatrix} 0.5 \\ 0.25 \\ 4 \end{bmatrix}$$

$$\lambda = \begin{bmatrix} 0.7 \\ 0.5 \end{bmatrix}$$

To test the solver random values for x and λ are generated, from these the H , g , A and b are determined. With these new matrix the system are solved again to see if the x and λ are the same. The solve works as expected and gives the same values for x and λ as there were generated.

To see the sensitivity of the problem the parameter sensitivity analyse is made. The general function is as followed.

$$F(x, \lambda; p) = \begin{bmatrix} \nabla_x f(x, p) - \nabla_x c(x, p) \lambda \\ c(x, p) \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

Where the implicit function theorem gives parameter sensitivities

$$[\nabla_x(p) \lambda(p)] = -[W_{xp} - \nabla_p c(x, p)] \begin{bmatrix} W_{xx} & -\nabla_x c(x, p) \\ -\nabla_x c(x, p)^T & 0 \end{bmatrix}^{-1}$$

$$W_{xx} = \nabla_{xx}^2 f(x, p) - \sum_{i \in \varepsilon} \lambda_i \nabla_{xx}^2 c_i(x, p)$$

$$W_{xp} = \nabla_{xp}^2 f(x, p) - \sum_{i \in \varepsilon} \lambda_i \nabla_{xp}^2 c_i(x, p)$$

From this the sensitivity of this problem can be written as:

$$f(x, p) = \frac{1}{2} x^T H x + (g + \begin{bmatrix} p_1 \\ p_2 \\ p_3 \end{bmatrix})^T x$$

$$c(x, p) = A^T x - b - \begin{bmatrix} p_5 \\ p_6 \end{bmatrix}$$

$$w_{xp} = \begin{bmatrix} 1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

$$W_{xx} = H$$

$$[\nabla x(p) \lambda(p)] = - \begin{bmatrix} H & -A^T \\ -A & 0 \end{bmatrix}$$

This are added to the solver and the sensitivity are found as:

2.2 Problem 2 - Equality Constrained Quadratic Optimization

blablablablab

2.3 Problem 3 - Inequality Constrained Quadratic Programming

From page 475 in Nocedal and Wright the following system is given.

$$\begin{aligned} \min_x q(x) &= (x_1 - 1)^2 + (x_2 - 2.5)^2 \\ s.t. \quad &x_1 - 2x_2 + 2 \geq 0, \\ &-x_1 - 2x_2 + 6 \geq 0, \\ &-x_1 + 2x_2 + 2 \geq 0, \\ &x_1 \geq 0, \\ &x_2 \geq 0. \end{aligned} \tag{1}$$

in MatLab a contour plot of this is made and seen in figure 1.




Figure 1: A contour plot of the problem.

From the line it is seen that the feasible region is a pentagon.
The problem can be written in the standart martix way as:

$$\begin{aligned}
 H &= \begin{bmatrix} 2 & 0 \\ 0 & 2 \end{bmatrix} \\
 g &= \begin{bmatrix} -2 \\ -5 \end{bmatrix} \\
 A &= \begin{bmatrix} 1 & -1 & -1 & 1 & 0 \\ -2 & -2 & 2 & 0 & 1 \end{bmatrix} \\
 b &= \begin{bmatrix} -2 \\ -6 \\ -2 \\ 0 \\ 0 \end{bmatrix}
 \end{aligned}$$

Then the general KKT system can be written as:

$$\begin{bmatrix} H & -A \\ A^T & 0 \end{bmatrix} \times \begin{bmatrix} x \\ \lambda \end{bmatrix} = \begin{bmatrix} -g \\ b \end{bmatrix}$$

When the system can be solve with the same QP solver as used for problem 1. That gives:

$$x = \begin{bmatrix} -2 \\ 0 \end{bmatrix}$$

$$\lambda = \begin{bmatrix} 0 \\ 1.8 \\ -0.3 \\ -2.1 \\ 3.0 \end{bmatrix}$$

Where all the values are multiply with 10 in the power of 16. When a active set algorithm can be apply for the problem by..???

2.4 Problem 4 - Markowitz Portfolio Optimization

blablablablbal

2.5 Problem 5 - Interior-Point Algorithm for Convex Quadratic Programming

blablablablbal

3 Conclusion

Some conclusions things