

Homework G.7 - Solution

The linearized equations are

$$m_1 \ddot{z}_1 + m_1 g \tilde{\theta} = 0$$

$$\left(\frac{m_2 l^2}{3} + m_1 z_e^2 \right) \ddot{\tilde{\theta}} + m_1 g (\tilde{z}) = l \tilde{F}.$$

Solving for $\ddot{\tilde{z}}$ and $\ddot{\tilde{\theta}}$ gives

$$\ddot{\tilde{z}} = -g \tilde{\theta}$$

$$\ddot{\tilde{\theta}} = \frac{l}{\frac{m_2 l^2}{3} + m_1 z_e^2} \tilde{F} - \frac{m_1 g}{\frac{m_2 l^2}{3} + m_1 z_e^2} \tilde{z}.$$

Let

$$\mathbf{x} = \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{bmatrix} \triangleq \begin{bmatrix} \tilde{\theta} \\ \tilde{z} \\ \dot{\tilde{\theta}} \\ \dot{\tilde{z}} \end{bmatrix}$$

$$u \triangleq \tilde{F}$$

$$\mathbf{y} \triangleq \begin{bmatrix} \tilde{z} \\ \tilde{\theta} \end{bmatrix}$$

then

$$\dot{x}_1 = \dot{\tilde{\theta}} = x_3$$

$$\dot{x}_2 = \dot{\tilde{z}} = x_4$$

$$\dot{x}_3 = \ddot{\tilde{\theta}} = \frac{l}{\frac{m_2 l^2}{3} + m_1 z_e^2} \tilde{F} - \frac{m_1 g}{\frac{m_2 l^2}{3} + m_1 z_e^2} \tilde{z}$$

$$= \frac{l}{\frac{m_2 l^2}{3} + m_1 z_e^2} u - \frac{m_1 g}{\frac{m_2 l^2}{3} + m_1 z_e^2} x_2$$

$$\dot{x}_4 = \ddot{\tilde{z}} = -g \tilde{\theta} = -g x_1.$$

In matrix form

$$\begin{aligned}
 \begin{bmatrix} \dot{x}_1 \\ \dot{x}_2 \\ \dot{x}_3 \\ \dot{x}_4 \end{bmatrix} &= \underbrace{\begin{bmatrix} 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ -g & 0 & 0 & 0 \\ 0 & -\frac{m_1 g}{\frac{m_2 l^2}{3} + m_1 z_e^2} & 0 & 0 \end{bmatrix}}_{\triangleq \mathbf{A}} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{bmatrix} + \underbrace{\begin{bmatrix} 0 \\ 0 \\ \frac{l}{\frac{m_2 l^2}{3} + m_1 z_e^2} \\ 0 \end{bmatrix}}_{\triangleq \mathbf{B}} u \\
 \mathbf{y} &= \underbrace{\begin{bmatrix} 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 \end{bmatrix}}_{\triangleq \mathbf{C}} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{bmatrix} + \underbrace{\begin{bmatrix} 0 \\ 0 \end{bmatrix}}_{\triangleq \mathbf{D}} u.
 \end{aligned}$$