

Algorithms for Programming Contests - Week 12

Tobias Meggendorfer, Philipp Meyer,
Christian Müller, Gregor Schwarz
`conpra@in.tum.de`

23.01.2019

Geometry

Classical geometry

- Points as (x, y) coordinates.
- Lines represented by two points, or e.g. with Hesse normal form.

Problems

- Special cases: e.g. meet of parallel vs. non-parallel lines
- No simple systematic representation of all objects and operations.

Towards projective geometry

Idea of projective geometry

- Represent all objects uniformly through (higher-dimensional) vectors.
- Construct new objects with vector products.
- Represent transformations through matrix multiplications.

Projective geometry

Homogeneous coordinates

- Drawing plane is embedded into \mathbb{R}^3 at $z = 1$.
- Homogenization:

$$\begin{pmatrix} x \\ y \end{pmatrix} \rightarrow \begin{pmatrix} x \\ y \\ 1 \end{pmatrix} \sim \lambda \begin{pmatrix} x \\ y \\ 1 \end{pmatrix} \quad \lambda \in \mathbb{R} \setminus \{0\}$$

- Identify scalar multiples as the same point.
- Normalization/Dehomogenization:


$$\begin{pmatrix} x \\ y \\ z \end{pmatrix} \sim \begin{pmatrix} x/z \\ y/z \\ 1 \end{pmatrix} \rightarrow \begin{pmatrix} x/z \\ y/z \end{pmatrix}$$

- What happens when $z = 0$?

Projective geometry

Points at infinity

- Consider $p(t) = (x \cdot t, y \cdot t, 1)^T \sim (x, y, 1/t)^T$.
- We have $\lim_{t \rightarrow \infty} p(t) = (x, y, 0)^T$.
- $(x, y, 0)^T$ is the point at infinity in the direction (x, y) .
- Every vector in $\mathbb{R}^3 \setminus \{(0, 0, 0)^T\}$ represents a point.



Projective geometry

Lines in projective geometry

- Every vector $l = (x, y, z)^T \neq (0, 0, 0)^T$ also represents a line.
- Line given by intersection of plane through $(0, 0, 0)^T$ where l is a normal with drawing plane.
- Scalar multiples represent the same line.

Projective geometry

Lines in projective geometry

- Every vector $l = (x, y, z)^T \neq (0, 0, 0)^T$ also represents a line.
- Line given by intersection of plane through $(0, 0, 0)^T$ where l is a normal with drawing plane.
- Scalar multiples represent the same line.

Lines at infinity

- One line not contained in \mathbb{R}^2 : $(0, 0, 1)^T$.
- Contains all the points at infinity.

Operations in projective space

Point-on-line

- A point p is on a line l iff the vectors are orthogonal.
- Vectors p and l orthogonal iff *dot/scalar product* $p \cdot l = 0$.

Operations in projective space

Point-on-line

- A point p is on a line l iff the vectors are orthogonal.
- Vectors p and l orthogonal iff *dot/scalar product* $p \cdot l = 0$.

Line through two points

- Line through points p and q has to be orthogonal to both vectors.
- Line given by *cross product* $p \times q$.

Operations in projective space

Point-on-line

- A point p is on a line l iff the vectors are orthogonal.
- Vectors p and l orthogonal iff *dot/scalar product* $p \cdot l = 0$.


Line through two points

- Line through points p and q has to be orthogonal to both vectors.
- Line given by *cross product* $p \times q$.


Intersection of two lines

- Intersection point of lines l and m has to be on l and m .
- Orthogonal to both: point given by $l \times m$.


Example: intersection of diagonals




Example: intersection of diagonals



Example: intersection of diagonals




Example: intersection of diagonals




Points at infinity revisited

What happens if we take the intersection of two parallel lines?



Points at infinity revisited


What happens if we take the intersection of two parallel lines?



$$p = l \times m = (3, 0, 0)^T \sim (1, 0, 0)^T \sim (-1, 0, 0)^T$$

Points at infinity revisited


What happens if we take the intersection of two parallel lines?



$$p = l \times m = (3, 0, 0)^T \sim (1, 0, 0)^T \sim (-1, 0, 0)^T$$

Points at infinity revisited


What happens if we take the intersection of two parallel lines?



$$p = l \times m = (3, 0, 0)^T \sim (1, 0, 0)^T \sim (-1, 0, 0)^T$$

Points at infinity revisited


What happens if we take the intersection of two parallel lines?



$$p = l \times m = (3, 0, 0)^T \sim (1, 0, 0)^T \sim (-1, 0, 0)^T$$

Points at infinity revisited


What happens if we take the intersection of two parallel lines?



$$p = l \times m = (3, 0, 0)^T \sim (1, 0, 0)^T \sim (-1, 0, 0)^T$$

Points at infinity revisited

Connection of points at infinity gives line at infinity:



$$l = p \times q = (1, 0, 0)^T \times (0, 1, 0)^T = (0, 0, 1)^T$$

Constructions with points at infinity

Parallel

- Line m parallel to l through a point p .
- Infinite point in direction of l : $q = l \times (0, 0, 1)^T$. Then $m = p \times q$.

Constructions with points at infinity

Parallel

- Line m parallel to l through a point p .
- Infinite point in direction of l : $q = l \times (0, 0, 1)^T$. Then $m = p \times q$.

Perpendicular

- Line m perpendicular to l through a point p .
- Infinite point in direction of l : $q = l \times (0, 0, 1)^T = (x, y, 0)^T$.
- Orthogonal direction to q : $q^\perp = (y, -x, 0)^T$. Then $m = p \times q^\perp$.

Constructions with points at infinity

Parallel

- Line m parallel to l through a point p .
- Infinite point in direction of l : $q = l \times (0, 0, 1)^T$. Then $m = p \times q$.

Perpendicular

- Line m perpendicular to l through a point p .
- Infinite point in direction of l : $q = l \times (0, 0, 1)^T = (x, y, 0)^T$.
- Orthogonal direction to q : $q^\perp = (y, -x, 0)^T$. Then $m = p \times q^\perp$.


Projection

- Projection of point p on line l .
- Compute perpendicular m , then $r = m \times l$.

Area of a triangle

- Find area of the triangle given by three points p , q and r .
- First normalize p, q, r (scale to $z = 1$).
- Obtain matrix $M = \begin{pmatrix} p & q & r \end{pmatrix}$.
- Area is then $\frac{1}{2} |\det M| = \frac{1}{2} |p \cdot (q \times r)|$.
- $p \cdot (q \times r) = (p \times q) \cdot r$ is called the *triple product* of p, q, r .
- Gives value of counter-clockwise function: $\text{CCW}(p, q, r) = p \cdot (q \times r)$.

Area of a triangle



- Volume of parallelepiped spanned by p, q, r : $V_{\text{par}} = |p \cdot (q \times r)|$
- Volume of tetrahedron spanned by p, q, r : $V_{\text{tetra}} = \frac{1}{6} V_{\text{par}}$.
- Volume of tetrahedron spanned by p, q, r : $V_{\text{tetra}} = \frac{1}{3} A_{\text{triangle}} \cdot h$, where A_{triangle} is area of base triangle (p, q, r) and h is the height.
- $h = 1$ by normalization $\Rightarrow A_{\text{triangle}} = \frac{1}{2} |p \cdot (q \times r)|$.

Transformations in Euclidean and projective space

Rotations

$$\begin{pmatrix} x \\ y \end{pmatrix} \mapsto \begin{pmatrix} \cos \alpha & -\sin \alpha \\ \sin \alpha & \cos \alpha \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix}$$

Translations

$$\begin{pmatrix} x \\ y \end{pmatrix} \mapsto \begin{pmatrix} x \\ y \end{pmatrix} + \begin{pmatrix} t_x \\ t_y \end{pmatrix}$$

Transformations in Euclidean and projective space

Rotations

$$\begin{pmatrix} x \\ y \end{pmatrix} \mapsto \begin{pmatrix} \cos \alpha & -\sin \alpha \\ \sin \alpha & \cos \alpha \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix}$$

Translations

$$\begin{pmatrix} x \\ y \end{pmatrix} \mapsto \begin{pmatrix} x \\ y \end{pmatrix} + \begin{pmatrix} t_x \\ t_y \end{pmatrix}$$

Problem: concatenation of operations

Transformations in Euclidean and projective space

Rotations

$$\begin{pmatrix} x \\ y \end{pmatrix} \mapsto \begin{pmatrix} \cos \alpha & -\sin \alpha \\ \sin \alpha & \cos \alpha \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix}$$

Translations

$$\begin{pmatrix} x \\ y \end{pmatrix} \mapsto \begin{pmatrix} x \\ y \end{pmatrix} + \begin{pmatrix} t_x \\ t_y \end{pmatrix}$$

Problem: concatenation of operations

In projective space

$$\begin{pmatrix} x \\ y \\ 1 \end{pmatrix} \mapsto \begin{pmatrix} \cos \alpha & -\sin \alpha & 0 \\ \sin \alpha & \cos \alpha & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} x \\ y \\ 1 \end{pmatrix}$$

$$\begin{pmatrix} x \\ y \\ 1 \end{pmatrix} \mapsto \begin{pmatrix} 1 & 0 & t_x \\ 0 & 1 & t_y \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} x \\ y \\ 1 \end{pmatrix}$$

Affine transformations

Transformations preserving parallelity

- Shifting
- Scaling
- Rotating
- Mirroring
- Shearing
- Combinations of all these operations

$$M = \begin{pmatrix} a & b & c \\ d & e & f \\ 0 & 0 & 1 \end{pmatrix}, \det(M) \neq 0 \qquad \begin{pmatrix} x \\ y \\ 1 \end{pmatrix} \mapsto M \cdot \begin{pmatrix} x \\ y \\ 1 \end{pmatrix}$$


Projective transformations

Arbitrary projective transformation


$$M = \begin{pmatrix} a & b & c \\ d & e & f \\ g & h & i \end{pmatrix}, \det(M) \neq 0 \qquad \begin{pmatrix} x \\ y \\ z \end{pmatrix} \mapsto M \cdot \begin{pmatrix} x \\ y \\ z \end{pmatrix}$$

- M and λM for $\lambda \neq 0$ represent the same transformation.
- All affine transformations and perspective distortions.
- Collinearity and point-on-line property is preserved.
- Every transformation preserving collinearity is projective.
- Uniquely determined by four non-collinear points and their images.

Example: projective transformation




Example: projective transformation



$$M = \begin{pmatrix} 77 & -11 & -20 \\ -8 & 97 & 3 \\ -26 & 29 & 89 \end{pmatrix}$$

Example: projective transformation



Computing projective transformations

Computing M

Given four points a, b, c, d and their respective images a', b', c', d' by a projective transformation, what is a matrix M for this transformation?

Assumption: no three points of a, b, c, d or a', b', c', d' are collinear.

Computing projective transformations

Computing M

Given four points a, b, c, d and their respective images a', b', c', d' by a projective transformation, what is a matrix M for this transformation?

Assumption: no three points of a, b, c, d or a', b', c', d' are collinear.

Solve linear equation system to find M and $\lambda_a, \lambda_b, \lambda_c, \lambda_d$ such that:

$$M \cdot a = \lambda_a a' \quad M \cdot b = \lambda_b b' \quad M \cdot c = \lambda_c c' \quad M \cdot d = \lambda_d d'$$

M unique up to scalar multiple: one λ can be fixed.

Solve linear equation system with $9 + 3 = 12$ variables and 12 equations.

Computing projective transformations

Fix $\lambda_d = 1$, and consider the following special case:

$$a = (1, 0, 0)^T \quad b = (0, 1, 0)^T \quad c = (0, 0, 1)^T \quad d = (1, 1, 1)^T$$

$$M \cdot \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix} = \lambda_a a' \quad M \cdot \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix} = \lambda_b b' \quad M \cdot \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix} = \lambda_c c' \quad M \cdot \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix} = d'$$

Computing projective transformations

Fix $\lambda_d = 1$, and consider the following special case:

$$a = (1, 0, 0)^T \quad b = (0, 1, 0)^T \quad c = (0, 0, 1)^T \quad d = (1, 1, 1)^T$$


$$M \cdot \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix} = \lambda_a a' \quad M \cdot \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix} = \lambda_b b' \quad M \cdot \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix} = \lambda_c c' \quad M \cdot \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix} = d'$$

$$M = \begin{pmatrix} | & | & | \\ \lambda_a a' & \lambda_b b' & \lambda_c c' \\ | & | & | \end{pmatrix} \quad \begin{pmatrix} | & | & | \\ a' & b' & c' \\ | & | & | \end{pmatrix} \cdot \begin{pmatrix} \lambda_a \\ \lambda_b \\ \lambda_c \end{pmatrix} = d'$$

Only need to invert single 3×3 matrix to obtain $\lambda_a, \lambda_b, \lambda_c$ and then M .

Computing projective transformations

General case:



Spot the point at infinity



More projective geometry

There is a lot more

- Circles, conic sections.
- Measurements of lengths and angles.
- Cross-ratios and projective invariants.
- Higher-dimensional projective geometry.
- \Rightarrow lectures “Geometriekalküle” and “Projective Geometry” by Prof. Richter-Gebert.