

1 Title:
2 Incorporation of water-derived hydrogen into methane during artificial maturation of kerogen under hydrothermal
3 conditions
4

5 A manuscript prepared for submission to *Organic Geochemistry* on 15 July 2021
6
7

8 Authors and affiliations:

9 David T. Wang^{a,b,c*}, Jeffrey S. Seewald^b, Eoghan P. Reeves^d, Shuhei Ono^a, and Sean P. Sylva^b

10 ^aDepartment of Earth, Atmospheric and Planetary Sciences, Massachusetts Institute of Technology, Cambridge, Massachusetts 02139, USA.

11 ^bMarine Chemistry and Geochemistry Department, Woods Hole Oceanographic Institution, Woods Hole, Massachusetts 02543, USA.

12 ^cExxonMobil Upstream Research Company, Spring, Texas 77389, USA.

13 ^dDepartment of Earth Science and Centre for Geobiology, University of Bergen, Bergen N-5020, Norway.
14

15 * Corresponding author at: Esso Exploration and Production Guyana Ltd, 86 Duke Street, Georgetown, Guyana.
16 E-mail address: dtw@alum.mit.edu or david.t.wang@exxonmobil.com (D.T. Wang).
17

18 **Keywords:**

19 methane, natural gas generation kinetics, D/H ratios, kerogen, clumped isotopologues, hydrogen isotope exchange,
20 water isotopes
21

22 **Abstract**

23 To investigate the origin of H in thermogenic methane, a sample of organic-rich Eagle Ford shale was reacted with
24 D₂O under hydrothermal conditions in a flexible Au-Ti cell hydrothermal apparatus in a water-to-rock ratio of
25 approximately 5:1. Temperatures were increased from 200 to 350 °C over the course of one month, maintaining
26 pressure at 350 bar, and the concentrations of aqueous species and methane isotopologues produced were quantified.
27 Production of H₂, CO₂, alkanes, and alkenes was observed. Methane formed during the early stages of the experi-
28 ment at 200 °C was primarily CH₄ with some CH₃D, whereas at higher temperatures, increasing proportions of
29 deuterated isotopologues were produced. Near the end of the experiment, the concentration of CD₄ exceeded that
30 of all other isotopologues combined. These results suggest that competition between rates of kerogen-water isotopic
31 exchange and natural gas generation may govern the D/H ratio of thermogenic gases.

32

33 Abstract: 147 words

34 Main Text: XXXX words

35

1. INTRODUCTION

Variation in δD values of thermogenic natural gases is often attributed to kinetically-controlled fractionation during pyrolysis of kerogen or oils. A number of studies have investigated how D/H ratios of methane and other hydrocarbons evolve with increasing maturity (Sackett, 1978; Berner et al., 1995; Sackett and Conkright, 1997; Tang et al., 2005; Ni et al., 2011). However, kinetic isotope effects involving hydrogen addition or abstraction are often large and by themselves do not explain the geologically-reasonable apparent equilibrium temperatures of ~ 150 to 220°C obtained for reservoir gases that have been studied for their clumped isotopologue compositions (Stolper et al., 2014, 2015; Wang et al., 2015a; Douglas et al., 2017; Young et al., 2017; Shuai et al., 2018; Giunta et al., 2019; Labidi et al., 2020; Thiagarajan et al., 2020). There is also evidence that δD values of CH_4 approach values expected for isotopic equilibrium between CH_4 and H_2O in formation waters at temperatures characterizing reservoirs and/or mature source rocks (~ 150 to 250°C) (Clayton, 2003; Wang et al., 2015a; Xie et al., 2021), although findings of insignificant hydrogen exchange occurring under these conditions also exist (Yeh and Epstein, 1981). In order for methane samples to have approached or attained equilibrium values of $\Delta^{13}\text{CH}_3\text{D}$ and $\Delta^{12}\text{CH}_2\text{D}_2$ —parameters that describe the abundance of clumped isotopologues relative to a population of molecules containing isotopes randomly distributed amongst them [e.g., Young et al. (2017)]—there must be a pathway by which either (i) isotopes can be exchanged amongst methane isotopologues alone, (ii) methane isotopologues exchange hydrogen with water or organic molecules, or (iii) methane isotopologues are derived from methyl moieties which contain C–H bonds that have pre-exchanged with water prior to forming methane (Hoering, 1984; Smith et al., 1985; Schimmelmann et al., 1999, 2006; Lis et al., 2006).

Here, we study the origin of C–H bonds in thermogenic methane by heating kerogen in the presence of D_2O and examining the degree of deuteration in the generated methane. This experiment is conceptually very similar to those conducted by Hoering (1984), Lewan (1997), and Schimmelmann et al. (2001). However, none of these workers quantified the extent of deuteration in the produced natural gases, though Lewan mentioned that methane formed in his experiments contained deuterium [Lewan (1997) and M. D. Lewan (pers. comm.)].

2. METHODS

2.1. Experimental methods

Experiments were conducted in a gold-titanium reaction cell housed within a flexible cell hydrothermal apparatus (Seyfried et al., 1987) at WHOI. The reaction cell was pre-treated prior to loading by soaking in concentrated HCl for 4 hours, followed by rinsing with water to pH neutral and drying in the oven. The exit tube of the apparatus was cleaned by forcing ~ 20 ml of MilliQ deionized water ($18.2\text{ M}\Omega$) through, followed by ~ 20 ml conc. HCl, ~ 100 ml water, ~ 20 ml conc. HNO_3 , and then ~ 100 ml of water until the pH tested 7 using pH paper.

The source material for this experiment was a hand sample of Upper Cretaceous Eagle Ford Shale taken from an outcrop in Uvalde County, Texas, USA (Hentz and Ruppel, 2010). The sample was kindly provided to J. Seewald by Keith F. M. Thompson (PetroSurveys, Inc.). There is no known oil or gas production from the Eagle Ford in Uvalde County [(Tian et al., 2013); and IHS (2019)]. The Eagle Ford here is thermally-immature [$R_o = 0.40$ – 0.55% , Cardneaux (2012); Cardneaux and Nunn (2013); and Harbor (2011)]. The sample was powdered to $<250\text{ }\mu\text{m}$ and Soxhlet-extracted (by Carl Johnson, WHOI) to remove bitumen and free hydrocarbons. In a subsequent step, the solvent-extracted residue was subjected to hydrochloric acid treatment to remove carbonate minerals. Elemental analysis (Table 1) of the original rock sample (UNEX), the Soxhlet-extracted rock sample (EX), and the decalci-

fied+extracted rock sample (DECA) indicates a total organic carbon (TOC) content of ~2.5% and a carbonate content of ~80% by weight. The H/C atomic ratio of the decalcified rock is 2.4. This value is probably several tens of percent higher than the actual H/C ratio of isolated kerogen (not determined) given that substantial amounts of H are likely borne by clays and other minerals that were not removed (Whelan and Thompson-Rizer, 1993; Baskin, 1997). The reaction cell was loaded with 10.03 grams of the EX powder.

Geochemical data for the Eagle Ford sample can be drawn from neighboring Kinney County, Texas, where complete sections of immature Eagle Ford have been recovered by the USGS (drill core GC-3; French et al., 2020) and Shell (Iona-1 drill core; Eldrett et al., 2014, 2015; Sun et al., 2016); there, the Eagle Ford also crops out, is immature, and is presumed to be geochemically similar. The high calcium carbonate content and relatively lower organic enrichment is consistent with data from the Upper Eagle Ford in the Shell Iona-1 core from neighboring Kinney County, Texas (Eldrett et al., 2015).

The starting fluid was heavy water (D₂O, 99% purity, Cambridge Isotope Laboratories, Inc.) containing some NaCl (0.497 mol/kg). The added NaCl allows for detection of dilution of the fluid by deionized water from the pressure vessel in the case of a leak in the reaction cell. The reaction cell was loaded with 55.03 g of this starting fluid. The pressure vessel was sealed and the reaction brought to initial condition (200 °C, 350 bar) rapidly. Several milliliters of fluid were bled during heat-up to prevent overpressurization, leaving an estimated 52.6 g of fluid in the cell at the beginning of the experiment (Table 2).

2.2. Analytical methods

To monitor the fluid composition and the extent of deuteration, sample aliquots of fluid were withdrawn through the capillary exit tube into gastight glass/PTFE syringes. Immediately prior to a sampling event, a small amount (~0.5 g) of fluid was removed and discarded in order to flush the exit tube of any residues. The concentration of molecular hydrogen (H₂) was determined after headspace extraction using a gas chromatograph supplied with nitrogen carrier gas, and equipped with a molecular sieve 5 Å column and thermal conductivity detector. Analytical reproducibility of H₂ data is ±10% or better (2σ). However, accuracy of reported concentrations is unknown, because the relative responses of H₂, HD, and D₂ (the latter likely to be the main form of molecular hydrogen) in the GC-TCD were not determined. Residual liquid after headspace extraction was diluted with MilliQ water and saved for analysis of major cations and anions, or stored with dichloromethane in the fridge in a screw capped vial for analysis of non-volatile organic compounds.

Concentrations of total dissolved inorganic carbon (ΣCO₂) and C₁ to C₆ alkanes and alkenes were determined using a purge-and-trap cryofocusing device coupled to a gas chromatograph equipped with a Porapak Q column and serially-connected thermal conductivity and flame ionization detectors. Analytical procedures were as described in Reeves et al. (2012). Analytical reproducibility on duplicate samples was ±5% or better (2σ). The C₅ and C₆ compounds could not be quantified accurately due to their semi-volatile nature; however, C₅ and C₆ were detected at all sampling points.

At each sampling, a separate ~1 to 2 ml aliquot was injected directly into a pre-weighed, evacuated serum vial capped with boiled blue butyl rubber stoppers, for analysis of the extent of deuteration of methane. A Hewlett-Packard (HP) 6890 gas chromatography-mass spectrometry (GC-MS) system equipped with a 5 Å molecular sieve column (HP-PLOT 30 m × 0.32 mm × 12.0 μm) and HP 5973 mass selective detector was used to determine the amount of deuteration in CH₄. Ion currents were monitored at integral masses between *m/z* 10 and 50. Extracted ion currents were quantified at *m/z* 14 through 20 for methane. Expected fragmentation patterns of the five methane-*d*

isotopologues (C^1H_4 , CH_3D , CH_2D_2 , CHD_3 , and CD_4) were determined by analysis of commercial synthetic standards (>98% purity, Cambridge Isotope Laboratories, Inc.). *Note to reader:* We will refer to the protiated methane isotopologue as C^1H_4 in the text when it is necessary to specifically distinguish it from bulk CH_4 .

3. RESULTS

3.1. Temperature and thermal maturity

Temperatures logged during the experiment are shown in **Fig. 1A**. A fluid sample was taken at the beginning and end of each temperature stage. One additional sample (#4) was drawn in the middle of the second temperature stage (300 °C).

Estimated thermal maturity as a function of time was calculated using EASY%Ro (Sweeney and Burnham, 1990), and is shown in **Fig. 1B**. Maturities encountered in the experiment spanned the entire range of the oil window (ca. 0.5% to 1.3% R_o -equivalent; Burnham, 2019).

3.2. Concentrations of aqueous species

3.2.1. Inorganic species

Measured concentrations of aqueous species are shown in **Fig. 2**. Concentrations of H_2 increased from undetectable (<10 $\mu\text{mol/kg}$) to up to 0.8 mmol/kg at the end of the experiment. Increasing concentrations of H_2 within temperature stages of the experiment suggests that generation of petroleum, as opposed to a mineral redox buffer, is influencing the H_2 concentration. H_2 increased much more slowly during the >300 °C stages compared to heating at 300 °C and below.

The concentration of ΣCO_2 increased during the early stages of the experiment, and leveled off at ~50 mmol/kg at 350 °C. The plateauing inorganic carbon concentration might indicate that carbonate reached saturation and began to precipitate (Seewald et al., 1998). Measurements of major cations may be used to validate this interpretation. Production of CO_2 as the most abundant product of hydrothermal alteration of kerogen is also consistent with prior experimental work (Seewald, 2003). Alternatively, carbonate could have been released from the rock as it had not been decalcified prior to heating.

3.2.2. Alkanes and alkenes

Concentrations of methane increased in every successive time step, as did concentrations of detected n -alkanes. Except for the beginning of the experiment, molar concentrations of C_1 and ΣC_{2-4} were very similar and increased in near lock step.

Alkenes (ethylene and propylene, **Fig. 2D–E**) rose in concentration with every increase in temperature, indicating generation of unsaturated hydrocarbons via thermolytic processes. While concentrations of n -alkanes increased monotonically from the beginning to end of each temperature stage, the concentrations of alkenes were flat—or in the 350 °C stage, trended downwards—with time during each stage. Concentrations of alkenes consistent with thermodynamic equilibrium at measured H_2 concentrations are on the order of $\sim 10^{-7.3}$ and $\sim 10^{-6.5}$ mol/kg for ethylene and propylene, respectively, at 350 °C (Reeves et al., 2012). These equilibrium concentrations are ~2 orders of magnitude lower than the observed alkene concentrations.

Evidence from hydrothermal experiments suggests that metastable alkane/alkene equilibrium should be attained under hydrothermal conditions with half-equilibration times of several hundred hours or less at temperatures of 325 to 350 °C (Seewald, 1994). Failure to achieve thermodynamic equilibrium within these timescales indicates that generation of thermogenic alkenes occurs concurrently with alkane/alkene hydrogen exchange. Various pyrolysis experiments have reported alkene production (Huizinga et al., 1987; Leif and Simoneit, 2000), lending further support to the hypothesis that continued production of alkenes competes with their conversion into alkanes via hydrogenation at these temperatures and timescales and under the redox conditions characterizing hydrothermal maturation of organic-rich mudrocks.

Unlike the C₂₊ alkanes, methane cannot dehydrogenate to form an alkene. Hence, hydrogen exchange of methane requires that the very stable C–H bond be broken. Under appropriate, generally water-absent conditions, methane exchanges hydrogen with certain catalytic materials such as γ -alumina at room temperature over hours to days (Sattler, 2018, and refs. therein) [or with organometallic catalysts under even colder conditions (Golden et al., 2001)]. However, such catalysts in their active forms are not known to occur naturally in aqueous environments. Experiments conducted by Reeves et al. (2012) with aqueous methane in the presence of iron-bearing minerals in flexible-cell Au-Ti reaction vessels revealed little-to-no observable exchange even over several months at temperatures as high as 323 °C (Reeves et al., 2012). Recently, Turner et al. (under review) conducted a set of experiments in flexible gold-cell hydrothermal reactors with CH₄ dissolved in supercritical water at 376 to 420 °C to constrain the rate of CH₄–H₂O hydrogen isotope exchange. Their results confirm that exchange occurs over timescales of hundreds of years at 300 °C and tens of years at 350 °C (half-exchange, $\tau_{1/2}$), much longer than the duration of our experiment. Therefore, hydrogen exchange of methane during our experiment was limited or insignificant, and the hydrogen isotopic composition of the generated CH₄ is controlled primarily by the processes of source rock-water hydrogen exchange and kinetic isotope fractionation during methane generation.

As discussed in the following section, this conclusion regarding the lack of significant CH₄–H₂O hydrogen exchange in our experiment is supported by the selective production of deuterated methane isotopologues.

3.3. Production of deuterated methane isotopologues

Mass spectra collected for standards are shown in Fig. 4. Fragment intensities were very similar to those determined by Dibeler and Mohler (1950). Mass spectra of samples are shown in Fig. 3. No methane peaks of usable size could be obtained for time point #1. All other time points yielded quantifiable extracted ion chromatogram peaks.

The mass spectra of commercial standards were used to fit the sample data using a constrained linear least-squares solver (LSQNONNEG) implemented in MATLAB. Estimated relative abundances of methane-*d* isotopologues are shown in Fig. 5A.¹ Relative abundances were converted into absolute abundances (Fig. 5B) by multiplying by the methane concentration. The proportion of D in methane-bound hydrogen, calculated from the isotopologue abundances, is shown in Fig. 6.

Methane formed during the early stages of the experiment at 200 °C was primarily C¹H₄ with some CH₃D, whereas at higher temperatures, the isotopologues produced consist almost exclusively of CD₄, CHD₃, and CH₃D (Fig. 5A and Fig. 6). These results suggest that at relatively lower temperatures of ~200 °C, the rate of methane generation

¹ Results of this experiment were first presented by one of us in the appendix of a Ph.D. thesis (Wang, 2017). That earlier analysis contained a mathematical error (neglected to divide by the relative peak areas of the pure isotopologue standards). As a result, Fig. B.3 of that thesis appears different than Fig. 5 in this paper.

approaches or exceeds the rate of D/H exchange between water and kerogen, whereas at higher temperatures, extensive D/H exchange between kerogen (or oils, if they are also precursors of methane) and water occurs prior to methane generation. CD₄ became the dominant methane species at temperatures of 300 °C and above, suggesting that more than 50% of all labile, methane-generating sites on kerogen were fully deuterated. Alternatively, the dominance of CD₄ might be explained by direct CH₄–H₂O isotopic exchange occurring after the generation of primarily non-deuterated methane. This is unlikely given the sluggish pace at which D/H exchange occurs for methane (Reeves et al., 2012; Wang et al., 2018). Experiments in which normal water is heated in the presence of CD₄ while the D/H of water is monitored may yield a more sensitive determination of the rate of CH₄–H₂O exchange.

Production of C¹H₄ in the first stage of the experiment (200 °C) indicates that the earliest “capping” hydrogen derives from kerogen or other H-containing species in the rock as opposed to from the H atoms of water. This can only be the case if kerogen has not yet undergone D/H exchange.² While constraints on timescales of D/H exchange at 200 °C are sparse, the available literature supports this assertion. Experiments conducted with model hydrocarbons indicate that D/H exchange of carbon-bound hydrogen takes at least several decades, much longer than the heating time in our experiment (Sessions et al., 2004; Schimmelmann et al., 2006; Sessions, 2016; and refs. therein).

Production of C¹H₄ and CH₃D appeared to cease by midway through the 300 °C stage (time point #4, 284 hours), or was overshadowed by the generation of much larger quantities of the higher isotopologues. Continued (though relatively minor) production of methane that was not fully-deuterated (CHD₃ and CH₃D, **Fig. 5B**) suggests that kerogen or oil from which methane was generated still did not fully exchange before methane formed.

If significant exchange were to occur, either between water and kerogen, or between water and generated methane after cracking, and this exchange occurs sequentially, the predominant isotopologue would be expected to follow the progression C¹H₄ → CH₃D → CH₂D₂ → CHD₃ → CD₄. Instead, CH₂D₂ represents a smaller fraction of the methane isotopologues than either CH₃D or CHD₃ at all times, and calculated proportions of CH₂D₂ do not exceed 10% at any point in the experiment (**Fig. 5A**). A possible explanation is that various CH_x moieties (e.g., aromatic C vs. methylene C vs. heteroatom-bound C) may have significantly different propensities to exchange and hydrogenation (cracking). Cracking that occurs much faster or slower than exchange may yield either fully-deuterated kerogen (e.g., –CD₃) or singly-deuterated methane, respectively, hence leading to an absence of CH₂D₂. Alternatively or additionally, D/H exchange of partially-deuterated longer-chain hydrocarbon molecules with water may be faster than cracking, such that the production of CH₂D₂ is “skipped”. The absence of CH₂D₂ is additional evidence that exchange between water and methane or methane and methane at temperatures of 200 to 350 °C is slow on the timescales relevant to these experiments, consistent with a prior set of experiments (Reeves et al., 2012). Rates of methane isotopic exchange could potentially be better constrained by heating normal water in the presence of CD₄ and monitoring the rate of increase in the δD value of water.

Fig. 6A shows the percentage of water-derived hydrogen in CH₄ vs. estimated maturity (EASY%Ro). Small symbols are from Wei et al. (2019). Thermal maturity for the Wei et al. data were calculated from a time-temperature curve reconstructed from their described experimental procedures. Both studies yielded methane with an increasing percentage of water-derived hydrogen as thermal maturity increased. The deuteration vs. maturity trends are sub-parallel to each other. The observed offset between the Wei et al. experimental results and ours is probably due to the different source rocks and experimental conditions, including the use of D₂O instead of normal water as the

² It is conceivable that the C¹H₄ observed at time point #2 may have been gas originally present but sorbed to a solid phase at the start of the experiment and later leached into the fluid, but we consider this unlikely because the concentration of methane tripled between time points #1 (19 h) and #2 (164 h). Release of sorbed gases was probably nearly complete by 19 h.

aqueous medium in our experiments. By the middle of the oil window (EASY%Ro = 0.75–0.9%), methane in both studies contained more than 50% of its hydrogen content derived from water.

Fig. 6B is a graph of cumulative CH₄ generated plotted against percentage of methane deuteration.³ Because approximately 100 μmol of CH₄ was generated in total, the x-axis of this panel can be read as % of cumulative methane generation. At 50% deuteration, only less than 10% of methane has been generated; stated another way, for 90% of the total methane generated in the experiment, more than half of the hydrogen content is derived from water. From **Fig. 5A**, the fully-deuterated isotopologue CD₄ predominates towards the end of the experiment (time points #7–9). These late time points mark the end of the oil window (EASY%Ro between 0.9 and 1.3%) (**Fig. 6A**), suggesting that the immediate precursors of methane have already fully-exchanged their hydrogens with water. The fourth (capping) H in methane may come directly from water or may be abstracted from deuterated kerogen (Dong et al., 2021).

4. DISCUSSION

4.1. Interpretation of D/H and clumped isotope signatures of CH₄

Efforts to understand the D/H ratios of natural gas hydrocarbons have generally been centered around determining the influence of thermal maturity, organic-inorganic interactions, catalysts, and/or biological processes on the fractionation of hydrogen isotopes in these molecules during their generation, alteration, and/or destruction in source rocks and reservoirs of sedimentary basins. Examples of quantitatively-based numerical models are those of (Sackett, 1978; Berner et al., 1995; Clayton, 2003; Tang et al., 2005; Lu et al., 2011, 2021; Ni et al., 2011, 2012).

Under hydrothermal conditions, water is known to provide hydrogen to methane via a free radical mechanism (He et al., 2019).

Correct interpretation of δD values and clumped isotope signatures of CH₄ depends on understanding the relative kinetics of (a) methane generation from kerogen maturation or cracking of high-molecular weight hydrocarbons; (b) hydrogen exchange of methane precursor molecules with other organic molecules and/or water; and (c) direct or indirect hydrogen exchange between CH₄ and H₂O. Timescales of all of these processes range between years to tens of millions of years at the peak petroleum-generating temperatures of 100 to 200 °C, hence the relative importance of these three processes broadly governs the amount of organic-derived and water-derived H in CH₄. These three processes are discussed separately here with respect to the experimental results and how they apply to the interpretation of isotope and isotopologue ratios of CH₄.

4.1.1. Methane generation

Methane is generated directly during catagenesis via cleavage of methyl groups from kerogen. It is also generated via cracking of high- and low-molecular weight hydrocarbons, low-molecular weight organic acids, and other organic molecules in source rocks and/or high-temperature reservoirs. Thermogenic methane production occurs over a very wide range of temperatures, with some reports of commercial volumes of thermogenic natural gas generated at temperatures lower than 86 °C (Laplante, 1974), perhaps even lower than 62 °C (Rowe and Muehlenbachs, 1999). Thermal maturities of corresponding source rocks of putative low-temperature hydrocarbon gases and condensates were estimated to be as low as ~0.25 to 0.4% R_o (Laplante, 1974; Stahl, 1977; Purcell et al., 1979; Connan and

³ Calculated as $[CH_4] \times V_{\text{remaining}} + \sum ([CH_4] \times V_{\text{withdrawn}})$.

Commented [WDT2]: file:///\\GYG-DAT01\dtwang\$\Sync\My%20Past%20Academic%20Life\HIPPIE-DIPPIE%20Revisited\matlab\concentration\TOTAL_GENERATED.xlsx

Cassou, 1980; Snowdon, 1980; Jenden et al., 1993; Muscio et al., 1994; Rowe and Muehlenbachs, 1999; Ramaswamy, 2002). Kerogen moieties will not have undergone much D/H exchange at these low thermal maturities (Dawson et al., 2005; Maslen et al., 2012; Vinnichenko et al., 2021), and thus CH₄ generated from immature or marginally-mature source rocks will partially inherit its hydrogen and their corresponding C–H linkages from the precursor organic matter. Since methyl groups of wood (and presumably other naturally-occurring organic matter) carry clumped isotope values that deviate from equilibrium (Lloyd et al., 2021a), and because equilibrium methyl group clumping values [$\Delta(^{13}\text{CH}_2\text{D}-\text{R})$ values] are quite similar to $\Delta^{13}\text{CH}_3\text{D}$ values of CH₄ at these temperatures (within several tenths of a permil; Wang et al., 2015a; Lloyd et al., 2021a), CH₄ generated from sedimentary organic matter at low levels of thermal stress will likely also carry non-equilibrated clumping values inherited from methane precursors. The process of terminating the CH₃· with a H· may be an additional source of disequilibrated clumped methane signatures (Dong et al., 2021).

(Sessions et al., 2004; Schimmelmann et al., 2006; Sessions, 2016).

at least certain timescales of direct hydrogen exchange between CH₄ and H₂O are likely in the tens of millions of years at temperatures below 200 °C

4.1.2. D/H exchange in precursor molecules

Since methyl groups of at least certain timescales of direct hydrogen exchange between CH₄ and ambient H₂O are likely in the tens of millions of years at temperatures below 200 °C

See (Wang et al., 2015b) for review on hydrogen exchange in precursors.

See (Lloyd et al., 2021a) and (Lloyd et al., 2021b; Turner et al., 2021)

4.1.3. D/H exchange between methane and water

The reader is referred to Reeves et al. (2012), Beaudry et al. (2021), and Turner et al. (under review) for further discussion on this.

The activation energy for methane generation in our experiment was 90.24 kJ/mol or 21.57 kcal/mol, with a pre-exponential factor of 2.19E-03/sec (*Supplementary Information*) Compare to (Seewald et al., 1998), it's different. Compare to (Lewan, 1985),

(Xie et al., 2021)

There are a number of features in the dataset that provide qualitative insight into the relative contribution of kerogen vs.

Four features in the dataset are notable: (i) the production of undeuterated C¹H₄ under incipient catagenic conditions; (ii) the predominance of CD₄ towards the end of the experiment, coinciding with the late oil window; (iii) the lack of direct methane-water isotopic exchange even at 350 °C; and (iv) the near-absence of CH₂D₂ during the experiment. These observations suggest that while some –CH₃ moieties in kerogen or longer-chain hydrocarbons undergo exchange more readily than cracking, some other moieties or compound classes are much less prone to exchange.

295 The

296 4.2. Generation potential of natural gas

297 Asdfasdasd

298

299 The volumetric significance of the water hydrogen reservoir hence may be more important than otherwise assumed.
300 **Is there HI vs. TOC vs. Ro data (or RockEval and TOC) data for global source rock databases? – ASK MIRELA,**
301 **KIRSTEN, GRADY**

302

303 No longer limited by HI can make as much as TOC allows (viz. Helgeson paper).

304 Other factors such as water pressure, fluid chemistry, fluid flow, the presence of oil or Type I organic matter and
305 methane or CO₂ partial pressures were found to play only minor roles in the development of vitrinite reflectance.
306 (Huang 1996, Exxon).

307 The literature is conflicted on the importance of water on vitrinite reflectance. For a review of the literature see
308 Hackley and Lewan, 2018.

309 We compared several different models of vitrinite reflectance: EASY%Ro (Sweeney & Burnham, 1990),
310 EASY%RoB (Burnham 2019), EASY%RoV (Burnham 2019), and Basin%Ro (Nielsen et al., 2016). We show
311 both F (fraction of oil generated as modeled) and modeled %Ro (Fig. S-1A, S-1B).

312 We assumed here that primary oil generation occurs principally within the range 0.5 to 1.3 %Ro-equivalent (Burn-
313 ham, 2019). Under conditions encountered in this experiment, the modeled conversion of oil is between 5% and
314 95%, respectively (Fig. S1).

315 Zhang et al

316 Oil cracking hydrous: Jin et al, He et al 2019

317 Kerogen maturation hydrous: Gao et al., 2014 (higher maturity)

318 French et al, Eagle Ford geochemistry for Sorg/TOC

319 While Lewan (1997) did not quantify the abundance of deuterated methane isotopologues, we understand that these
320 analyses have been conducted in follow-up. The data also show production of deuterated and perdeuterated isotop-
321 logues of methane (M.D. Lewan, personal correspondence).

322

323 While examining the total ion and extracted ion chromatograms to quantify the deuteration in CH₄, an unknown
324 and unexpected peak was found eluting immediately following the CH₄ and air peaks. This mystery peak appeared
325 to yield methyl fragments that were also progressively more deuterated with reaction time. Re-analysis of several
326 samples while scanning a higher mass range suggested that the mystery compound had stable fragments near m/z
327 45 to 50 (depending on degree of deuterium substitution). This was verified by GC-MS analysis of a commercial

isobutane standard (mostly isobutane-*d*₀) which yielded a base peak at *m/z* 43. No attempt to quantify the degree of deuteration in isobutane was made.

Comments in Wei et al., 2019, Org Geochem

Aromatic Hydrogen Exchange in Petroleum Source Rocks


R. Alexander, R. I. Kagi, A. V. Luchini and G. W. Woodburn

School of Applied Chemistry, Western Australian Institute of Technology, 801 Sturt Street, Bentley, 6102, Western Australia, Australia

5. CONCLUSIONS

Four features in the dataset are notable: (i) the production of undeuterated C¹H₄ under incipient catagenic conditions; (ii) the predominance of CD₄ towards the end of the experiment, coinciding with the late oil window; (iii) the lack of direct methane-water isotopic exchange even at 350 °C; and (iv) the near-absence of CH₂D₂ during the experiment. These observations suggest that while some –CH₃ moieties in kerogen or longer-chain hydrocarbons undergo exchange more readily than cracking, some other moieties or compound classes are much less prone to exchange.

Carefully-controlled, temperature-programmed hydrous deuteration (deuterous pyrolysis or deuterothermal pyrolysis) experiments on additional source rocks and kerogen types may reveal systematic differences in the kinetics of exchangeability vs. hydrocarbon generation. Such experiments have the potential of improving prediction of generative yields and oil compositions in basins where timing and quality of hydrocarbon charge are key uncertainties.



Oil window is 0.5 to 1.3 % Ro.

Thank you for giving me the link to your dissertation concerning clump isotopes of methane from various origins and settings. You did a fantastic job in interpreting, organizing and presenting the data on aqueous methane! I also enjoyed your pyrolysis work using the aqueous methane generated from the Eagle Ford sample.

The geochemistry that intrigues me is the reactions involving water dissolved in the bitumen of a source rock. Experiments indicate that the amount of water dissolved in the organic matter of a source rock determines the amount of oil expelled and retained in a maturing source rock. Understanding this aspect of dissolved water rather than aqueous organics is in its infancy, similar to that of the role of water dissolved in granite melts back in the 1930s. How does the amount of dissolved water and its chemistry influence the immiscibility of oil, hydrogen availability, and expulsion of oil from a source rock? Also, what reactions are occurring between dissolved water and bitumen to form H₂ and CO₂?

With respect to incorporating the role of water in numerical models there is a lot of work to be undertaken, but a Ph.D dissertation at Colorado School of Mines by Mohammed Al Duhaian (now with Saudi Aramco) has made

some preliminary quantitative attempts (attached). We have been doing some work on the effects of water on the mechanical properties of source rocks (attached). These studies are still rather qualitative but we are moving toward a more quantitative understanding. I put more emphasis on experimentation before numerical modeling is attempted.

6. ACKNOWLEDGMENTS

Financial support from the U.S. National Science Foundation (NSF awards EAR-1250394 to S.O., and XXX to J.S.S.), the Alfred P. Sloan Foundation via the Deep Carbon Observatory (to S.O. and J.S.S.), a Shell-MIT Energy Initiative Fellowship, and the Kerr-McGee Professorship at MIT (to S.O.) is acknowledged. We thank Aaron Sattler (ExxonMobil Research and Engineering) for advice on inverting mass spectral data, Keith F. M. Thompson for providing the Eagle Ford rock sample, and Chris Clayton for an email exchange that inspired this work. Comments by Michael Lewan on an earlier draft of this paper are gratefully acknowledged.

Potential reviewers: Katherine L. French, Justin Birdwell, Arndt Schimmelmann, Alex Sessions, Michael Lewan

Conflicted: John Eiler, Dan Stolper, Ed Young

7. REFERENCES

- Baskin, D.K., 1997. Atomic H/C ratio of kerogen as an estimate of thermal maturity and organic matter conversion. AAPG bulletin 81, 1437–1450.
- Beaudry, P., Stefánsson, A., Fiebig, J., Rhim, J.H., Ono, S., 2021. High temperature generation and equilibration of methane in terrestrial geothermal systems: Evidence from clumped isotopologues. *Geochimica et Cosmochimica Acta* 309, 209–234.
- Berner, U., Faber, E., Scheeder, G., Panten, D., 1995. Primary cracking of algal and landplant kerogens: kinetic models of isotope variations in methane, ethane and propane. *Chemical Geology* 126, 233–245.
- Burnham, A.K., 2019. Kinetic models of vitrinite, kerogen, and bitumen reflectance. *Organic Geochemistry* 131, 50–59.
- Cardneaux, A., Nunn, J.A., 2013. Estimates of maturation and TOC from log data in the Eagle Ford Shale, Maverick Basin of South Texas. *Gulf Coast Association of Geological Societies Transactions* 63, 111–124.
- Cardneaux, A.P., 2012. Mapping of the oil window in the Eagle Ford shale play of southwest Texas using thermal modeling and log overlay analysis (Masters Thesis). Louisiana State University.
- Clayton, C., 2003. Hydrogen isotope systematics of thermally generated natural gas. *International Meeting on Organic Geochemistry*, 21st, Kraków, Poland, Book Abstr. Part I 51–52.
- Connan, J., Cassou, A.M., 1980. Properties of gases and petroleum liquids derived from terrestrial kerogen at various maturation levels. *Geochimica et Cosmochimica Acta* 44, 1–23.
- Dawson, D., Grice, K., Alexander, R., 2005. Effect of maturation on the indigenous δD signatures of individual hydrocarbons in sediments and crude oils from the Perth Basin (Western Australia). *Organic Geochemistry* 36, 95–104.
- Dibeler, V.H., Mohler, F.L., 1950. Mass spectra of the deuteromethanes. *J. Research Nat. Bur. Standards* 45, 441–444.
- Dong, G., Xie, H., Formolo, M., Lawson, M., Sessions, A., Eiler, J., 2021. Clumped isotope effects of thermogenic methane formation: insights from pyrolysis of hydrocarbons. *Geochimica et Cosmochimica Acta*. doi:10.1016/j.gca.2021.03.009
- Douglas, P.M., Stolper, D.A., Eiler, J.M., Sessions, A.L., Lawson, M., Shuai, Y., Bishop, A., Podlaha, O.G., Ferreira, A.A., Neto, E.V.S., others, 2017. Methane clumped isotopes: Progress and potential for a new isotopic tracer. *Organic Geochemistry* 113, 262–282.
- Eldrett, J.S., Ma, C., Bergman, S.C., Lutz, B., Gregory, F.J., Dodsworth, P., Phipps, M., Hardas, P., Minisini, D., Ozkan, A., Ramezani, J., Bowring, S.A., Kamo, S.L., Ferguson, K., Macaulay, C., Kelly, A.E., 2015. An astronomically calibrated stratigraphy of the Cenomanian, Turonian and earliest Coniacian from the Cretaceous Western Interior Seaway, USA: Implications for global chronostratigraphy. *Cretaceous Research* 56, 316–344.
- Eldrett, J.S., Minisini, D., Bergman, S.C., 2014. Decoupling of the carbon cycle during Ocean Anoxic Event 2. *Geology* 42, 567–570.
- French, K.L., Birdwell, J.E., Lewan, M.D., 2020. Trends in thermal maturity indicators for the organic sulfur-rich Eagle Ford Shale. *Marine and Petroleum Geology* 118, 104459.
- Giunta, T., Young, E.D., Warr, O., Kohl, I., Ash, J.L., Martini, A., Mundle, S.O., Rumble, D., Pérez-Rodríguez, I., Wasley, M., others, 2019. Methane sources and sinks in continental sedimentary systems: New insights from paired clumped isotopologues 13CH3D and 12CH2D2. *Geochimica et Cosmochimica Acta* 245, 327–351.
- Glasoe, P.K., Long, F.A., 1960. Use of glass electrodes to measure acidities in deuterium oxide. *The Journal of Physical Chemistry* 64, 188–190.
- Golden, J.T., Andersen, R.A., Bergman, R.G., 2001. Exceptionally low-temperature carbon–hydrogen/carbon–deuterium exchange reactions of organic and organometallic compounds catalyzed by the $Cp^*(PMe_3)IrH(ClCH_2Cl)^+$ cation. *Journal of the American Chemical Society* 123, 5837–5838.
- Harbor, R.L., 2011. Facies characterization and stratigraphic architecture of organic-rich mudrocks, Upper Cretaceous Eagle Ford Formation, South Texas (Masters Thesis). University of Texas at Austin.

- He, K., Zhang, S., Mi, J., Fang, Y., Zhang, W., 2019. Carbon and hydrogen isotope fractionation for methane from non-isothermal pyrolysis of oil in anhydrous and hydrothermal conditions. *Energy Exploration & Exploitation* 37, 1558–1576.
- Hentz, T.F., Ruppel, S.C., 2010. Regional lithostratigraphy of the Eagle Ford Shale: Maverick Basin to East Texas Basin. *Gulf Coast Association of Geological Societies Transactions* 60, 325–337.
- Hoering, T., 1984. Thermal reactions of kerogen with added water, heavy water and pure organic substances. *Organic Geochemistry* 5, 267–278.
- Huizinga, B.J., Tannenbaum, E., Kaplan, I.R., 1987. The role of minerals in the thermal alteration of organic matter—IV. Generation of n-alkanes, acyclic isoprenoids, and alkenes in laboratory experiments. *Geochimica et Cosmochimica Acta* 51, 1083–1097.
- Hunt, J.M., 1996. *Petroleum geochemistry and geology*, 2nd ed. IHS Markit Well Database, 2019.
- Jenden, P.D., Drazan, D.J., Kaplan, I.R., 1993. Mixing of Thermogenic Natural Gases in Northern Appalachian Basin. *AAPG Bulletin* 77, 980–998.
- Labidi, J., Young, E.D., Giunta, T., Kohl, I.E., Seewald, J., Tang, H., Lilley, M.D., Früh-Green, G.L., 2020. Methane thermometry in deep-sea hydrothermal systems: Evidence for re-ordering of doubly-substituted isotopologues during fluid cooling. *Geochimica et Cosmochimica Acta* 288, 248–261.
- Laplante, R.E., 1974. Hydrocarbon Generation in Gulf Coast Tertiary Sediments. *AAPG Bulletin* 58, 1281–1289.
- Leif, R.N., Simoneit, B.R., 2000. The role of alkenes produced during hydrous pyrolysis of a shale. *Organic Geochemistry* 31, 1189–1208.
- Lewan, M., 1985. Evaluation of petroleum generation by hydrous pyrolysis experimentation [and discussion]. *Philosophical Transactions of the Royal Society of London. Series A, Mathematical and Physical Sciences* 315, 123–134.
- Lewan, M., 1997. Experiments on the role of water in petroleum formation. *Geochimica et Cosmochimica Acta* 61, 3691–3723.
- Lis, G.P., Schimmelmann, A., Mastalerz, M., 2006. D/H ratios and hydrogen exchangeability of type-II kerogens with increasing thermal maturity. *Organic Geochemistry* 37, 342–353.
- Lloyd, M.K., Eldridge, D.L., Stolper, D.A., 2021a. Clumped $^{13}\text{CH}_2\text{D}$ and $^{12}\text{CHD}_2$ compositions of methyl groups from wood and synthetic monomers: Methods, experimental and theoretical calibrations, and initial results. *Geochimica et Cosmochimica Acta* 297, 233–275.
- Lloyd, M.K., Trembath-Reichert, E., Dawson, K.S., Feakins, S.J., Mastalerz, M., Orphan, V.J., Sessions, A.L., Eiler, J.M., 2021b. Methoxyl stable isotopic constraints on the origins and limits of coal-bed methane. *Science* 374, 894–897.
- Lu, S., Wang, M., Xue, H., Li, J., Chen, F., Xu, Q., 2011. The Impact of Aqueous Medium on Gas Yields and Kinetic Behaviors of Hydrogen Isotope Fractionation during Organic Matter Thermal Degradation. *Acta Geologica Sinica - English Edition* 85, 1466–1477.
- Lu, S.-F., Feng, G.-Q., Shao, M.-L., Li, J.-J., Xue, H.-T., Wang, M., Chen, F.-W., Li, W.-B., Pang, X.-T., 2021. Kinetics and fractionation of hydrogen isotopes during gas formation from representative functional groups. *Petroleum Science* 18, 1021–1032.
- Maslen, E., Grice, K., Dawson, D., Wang, S., Horsfield, B., 2012. Stable Hydrogen Isotopes of Isoprenoids and N-Alkanes as a Proxy for Estimating the Thermal History of Sediments Through Geological Time., in: Harris, N.B., Peters, K.E. (Eds.), *Analyzing the Thermal History of Sedimentary Basins: Methods and Case Studies*, SEPM Society for Sedimentary Geology, p. 0.
- Muscio, G.P.A., Horsfield, B., Welte, D.H., 1994. Occurrence of thermogenic gas in the immature zone—implications from the Bakken in-source reservoir system. *Organic Geochemistry* 22, 461–476.
- Ni, Y., Liao, F., Dai, J., Zou, C., Zhu, G., Zhang, B., Liu, Q., 2012. Using Carbon and Hydrogen Isotopes to Quantify Gas Maturity, Formation Temperature, and Formation Age — Specific Applications for Gas Fields from the Tarim Basin, China. *Energy Exploration & Exploitation* 30, 273–293.
- Ni, Y., Ma, Q., Ellis, G.S., Dai, J., Katz, B., Zhang, S., Tang, Y., 2011. Fundamental studies on kinetic isotope effect (KIE) of hydrogen isotope fractionation in natural gas systems. *Geochimica et Cosmochimica Acta* 75, 2696–2707.

475 Purcell, L.P., Rashid, M.A., Hardy, I.A., 1979. Geochemical Characteristics of Sedimentary Rocks in Scotian Basin.
 476 AAPG Bulletin 63, 87–105.

477 Ramaswamy, G., 2002. A field evidence for mineral-catalyzed formation of gas during coal maturation. *Oil & gas*
 478 *journal* 100, 32–36.

479 Reeves, E.P., Seewald, J.S., Sylva, S.P., 2012. Hydrogen isotope exchange between \emph{n-alkanes and water
 480 under hydrothermal conditions. *Geochimica et Cosmochimica Acta* 77, 582–599.

481 Rowe, D., Muehlenbachs, K., 1999. Low-temperature thermal generation of hydrocarbon gases in shallow shales.
 482 *Nature* 398, 61–63.

483 Sackett, W.M., 1978. Carbon and hydrogen isotope effects during the thermocatalytic production of hydrocarbons
 484 in laboratory simulation experiments. *Geochimica et Cosmochimica Acta* 42, 571–580.

485 Sackett, W.M., Conkright, M., 1997. Summary and re-evaluation of the high-temperature isotope geochemistry of
 486 methane. *Geochimica et Cosmochimica Acta* 61, 1941–1952.

487 Sattler, A., 2018. Hydrogen/Deuterium (H/D) exchange catalysis in alkanes. *ACS Catalysis* 8, 2296–2312.

488 Schimmelmann, A., Boudou, J.-P., Lewan, M.D., Wintsch, R.P., 2001. Experimental controls on D/H and 13C/12C ratios of kerogen, bitumen and oil during hydrous pyrolysis. *Organic Geo-*
 489 *chemistry* 32, 1009–1018.

491 Schimmelmann, A., Lewan, M.D., Wintsch, R.P., 1999. D/H isotope ratios of kerogen, bitumen, oil, and water in
 492 hydrous pyrolysis of source rocks containing kerogen types I, II, IIS, and III. *Geochimica et Cosmochimica*
 493 *Acta* 63, 3751–3766.

494 Schimmelmann, A., Sessions, A.L., Mastalerz, M., 2006. Hydrogen isotopic (D/H) composition of organic matter
 495 during diagenesis and thermal maturation. *Annual Review of Earth and Planetary Sciences* 34, 501–533.

496 Seewald, J.S., 1994. Evidence for metastable equilibrium between hydrocarbons under hydrothermal conditions.
 497 *Nature* 370, 285–287.

498 Seewald, J.S., 2003. Organic–inorganic interactions in petroleum-producing sedimentary basins. *Nature* 426, 327–
 499 333.

500 Seewald, J.S., Benitez-Nelson, B.C., Whelan, J.K., 1998. Laboratory and theoretical constraints on the generation
 501 and composition of natural gas. *Geochimica et Cosmochimica Acta* 62, 1599–1617.

502 Sessions, A.L., 2016. Factors controlling the deuterium contents of sedimentary hydrocarbons. *Organic Geochem-*
 503 *istry* 96, 43–64.

504 Sessions, A.L., Sylva, S.P., Summons, R.E., Hayes, J.M., 2004. Isotopic exchange of carbon-bound hydrogen over
 505 geologic timescales. *Geochimica et Cosmochimica Acta* 68, 1545–1559.

506 Seyfried, W.E., Jr., Janecky, D.R., Berndt, M.E., 1987. Rocking autoclaves for hydrothermal experiments, II. The
 507 flexible reaction-cell system. *Hydrothermal Experimental Techniques* 9, 216–239.

508 Shuai, Y., Etiope, G., Zhang, S., Douglas, P.M., Huang, L., Eiler, J.M., 2018. Methane clumped isotopes in the
 509 Songliao Basin (China): New insights into abiotic vs. biotic hydrocarbon formation. *Earth and Planetary*
 510 *Science Letters* 482, 213–221.

511 Smith, J., Rigby, D., Gould, K., Hart, G., Hargraves, A., 1985. An isotopic study of hydrocarbon generation pro-
 512 cesses. *Organic Geochemistry* 8, 341–347.


513 Snowdon, L., 1980. Resinite—A potential petroleum source in the upper Cretaceous/Tertiary of the Beaufort-Mac-
 514 kenzie Basin, in: Miall, A.D. (Ed.), *Facts and Principles of World Petroleum Occurrence*, CSPG Special
 515 *Publications Memoir* 6, p. 509–521.

516 Stahl, W.J., 1977. Carbon and nitrogen isotopes in hydrocarbon research and exploration. *Chemical Geology* 20,
 517 121–149.

518 Stolper, D., Martini, A., Clog, M., Douglas, P., Shusta, S., Valentine, D., Sessions, A., Eiler, J., 2015. Distinguishing
 519 and understanding thermogenic and biogenic sources of methane using multiply substituted isotopologues.
 520 *Geochimica et Cosmochimica Acta* 161, 219–247.


521 Stolper, D.A., Lawson, M., Davis, C.L., Ferreira, A.A., Santos Neto, E.V., Ellis, G.S., Lewan, M.D., Martini, A.M.,
 522 Tang, Y., Schoell, M., Sessions, A.L., Eiler, J.M., 2014. Formation temperatures of thermogenic and bio-
 523 genic methane. *Science* 344, 1500–1503.

- Sun, X., Zhang, T., Sun, Y., Milliken, K.L., Sun, D., 2016. Geochemical evidence of organic matter source input and depositional environments in the lower and upper Eagle Ford Formation, south Texas. *Organic Geochemistry* 98, 66–81.
- Sweeney, J.J., Burnham, A.K., 1990. Evaluation of a Simple Model of Vitrinite Reflectance Based on Chemical Kinetics (1). *AAPG Bulletin* 74, 1559–1570.
- Tang, Y., Huang, Y., Ellis, G.S., Wang, Y., Kralert, P.G., Gillaizeau, B., Ma, Q., Hwang, R., 2005. A kinetic model for thermally induced hydrogen and carbon isotope fractionation of individual *n*-alkanes in crude oil. *Geochimica et Cosmochimica Acta* 69, 4505–4520.
- Thiagarajan, N., Xie, H., Ponton, C., Kitchen, N., Peterson, B., Lawson, M., Formolo, M., Xiao, Y., Eiler, J., 2020. Isotopic evidence for quasi-equilibrium chemistry in thermally mature natural gases. *Proceedings of the National Academy of Sciences* 117, 3989–3995.
- Tian, Y., Ayers, W.B., McCain Jr, D., 2013. The Eagle Ford Shale play, south Texas: regional variations in fluid types, hydrocarbon production and reservoir properties, in: *IPC 2013: International Petroleum Technology Conference, European Association of Geoscientists & Engineers*, p. IPTC 16808.
- Turner, A.C., Korol, R., Eldridge, D.L., Bill, M., Conrad, M.E., Miller, T.F., Stolper, D.A., 2021. Experimental and theoretical determinations of hydrogen isotopic equilibrium in the system $\text{CH}_4\text{--H}_2\text{--H}_2\text{O}$ from 3 to 200 °C. *Geochimica et Cosmochimica Acta* 314, 223–269.
- Turner, A.C., Pester, N.J., Bill, M., Conrad, M.E., Knauss, K.G., Stolper, D.A., under review. Experimental determination of hydrogen isotope exchange rates between methane and water under hydrothermal conditions. *Geochimica et Cosmochimica Acta*.
- Vinnichenko, G., Jarrett, A.J.M., van Maldegem, L.M., Brocks, J.J., 2021. Substantial maturity influence on carbon and hydrogen isotopic composition of *n*-alkanes in sedimentary rocks. *Organic Geochemistry* 152, 104171.
- Wang, D.T., 2017. The geochemistry of methane isotopologues (PhD Thesis). Massachusetts Institute of Technology and Woods Hole Oceanographic Institution. doi:10.1575/1912/9052
- Wang, D.T., Gruen, D.S., Lollar, B.S., Hinrichs, K.-U., Stewart, L.C., Holden, J.F., Hristov, A.N., Pohlman, J.W., Morrill, P.L., Könneke, M., Delwiche, K.B., Reeves, E.P., Sutcliffe, C.N., Ritter, D.J., Seewald, J.S., McIntosh, J.C., Hemond, H.F., Kubo, M.D., Cardace, D., Hoehler, T.M., Ono, S., 2015a. Nonequilibrium clumped isotope signals in microbial methane. *Science* 348, 428–431.
- Wang, D.T., Reeves, E.P., McDermott, J.M., Seewald, J.S., Ono, S., 2018. Clumped isotopologue constraints on the origin of methane at seafloor hot springs. *Geochimica et Cosmochimica Acta* 223, 141–158.
- Wang, X., Liu, W., Shi, B., Zhang, Z., Xu, Y., Zheng, J., 2015b. Hydrogen isotope characteristics of thermogenic methane in Chinese sedimentary basins. *Organic Geochemistry* 83, 178–189.
- Wei, L., Gao, Z., Mastalerz, M., Schimmelmann, A., Gao, L., Wang, X., Liu, X., Wang, Y., Qiu, Z., 2019. Influence of water hydrogen on the hydrogen stable isotope ratio of methane at low versus high temperatures of methanogenesis. *Organic Geochemistry* 128, 137–147.
- Whelan, J.K., Thompson-Rizer, C.L., 1993. Chemical Methods for Assessing Kerogen and Protokerogen Types and Maturity, in: Engel, M.H., Macko, S.A. (Eds.), *Organic Geochemistry: Principles and Applications*, Springer US, Boston, MA, p. 289–353.
- Xie, H., Dong, G., Formolo, M., Lawson, M., Liu, J., Cong, F., Mangenot, X., Shuai, Y., Ponton, C., Eiler, J., 2021. The evolution of intra- and inter-molecular isotope equilibria in natural gases with thermal maturation. *Geochimica et Cosmochimica Acta*. doi:10.1016/j.gca.2021.05.012
- Yeh, H.-W., Epstein, S., 1981. Hydrogen and carbon isotopes of petroleum and related organic matter. *Geochimica et Cosmochimica Acta* 45, 753–762.
- Young, E.D., Kohl, I.E., Lollar, B.S., Etiope, G., Rumble, D., Li, S., Haghnegahdar, M.A., Schauble, E.A., McCain, K.A., Foustoukos, D.I., Sutcliffe, C., Warr, O., Ballentine, C.J., Onstott, T.C., Hosgormez, H., Neubeck, A., Marques, J.M., Pérez-Rodríguez, I., Rowe, A.R., LaRowe, D.E., Magnabosco, C., Yeung, L.Y., Ash, J.L., Bryndzia, L.T., 2017. The relative abundances of resolved $^{12}\text{CH}_2\text{D}_2$ and $^{13}\text{CH}_3\text{D}$ and mechanisms controlling isotopic bond ordering in abiotic and biotic methane gases. *Geochimica et Cosmochimica Acta* 203, 235–264.



575


8. FIGURES



576


577 **Fig. 1.** Profiles of (A) temperature and (B) estimated maturity (as EASY%Ro) vs. time. Time zero ($t = 0$) is the
578 time at which the experiment was brought to initial conditions (200 °C and 350 bar). Numbers in circles represent
579 sampling points (Table 2).

580



581

582 **Fig. 2.** Concentrations of aqueous species over time during the experiment. (A) Hydrogen (measured as H_2); (B) methane; (C) total inorganic
 583 carbon (ΣCO_2); (D) ethane and ethylene; (E) propane and propylene; and (F) *n*-butane and isobutane. Note that injection of additional saline D_2O
 584 at 430 hours diluted the concentration of all aqueous species by ~50%. BDL, below detection limit ($<10 \mu\text{mol/kg}$ for H_2).



585

586 **Fig. 3.** Mass spectra of standards. Isotopologue is indicated in the upper left corner of each plot. Intensities were
 587 normalized such that the m/z 14 to 20 signals sum to unity.






Fig. 4. Mass spectra of samples. Time point is indicated in the upper left corner of each plot. Intensities were normalized such that the m/z 14 to 20 signals sum to unity. Two samples were taken for time point #6, hence there are two plots. No GC-MS data was obtained for time point #1.



593

594 **Fig. 5.** Calculated (A) relative and (B) absolute abundances of methane isotopologues.



595

596 **Fig. 6.** Extent of methane deuteration [methane-bound D/(D+H)] vs. **(A)** estimated maturity (EASY%Ro) and **(B)**
 597 cumulative methane generated. The data shown for time point #6 is the average of the two replicate samples. Small
 598 symbols in (A) are data from Wei et al. (2019) representing percentage of water-derived H in CH₄ (see text). Pie
 599 charts above (B) represent relative abundances of isotopologues before, during, and after peak oil generation (time
 600 points #2, 4, and 9, respectively).

601




Fig. 7. Cartoon showing process of sequential deuteration of kerogen and oil along with generation of deuterated methane. Snapshots shown represent stages of (A) mild heating (incipient catagenesis); (B) moderate heating (oil generation); and (C) extensive heating (gas window).




Fig. 8. Schematic yields of oil and natural gas when generation occurs from source rock in the absence (A) and presence (B) of water as a source of hydrogen. (A) Traditional model of the amount and timing of organic alteration products generated during progressive burial in sedimentary basins that assumes oxygen and hydrogen in organic alteration products are derived only from kerogen. The form of this figure is constrained by the maturation trends shown in the Van Krevelen diagram. (B) Schematic illustration of the amount and timing of organic alteration products generated if water and minerals are allowed to contribute the requisite hydrogen and oxygen for the formation of hydrocarbons. Illustration is after Seewald (2003) and Hunt (1996).

614 **TABLES**

615

616 **Table 1**

617 Elemental analysis of Eagle Ford shale powder that was either: dried but otherwise untreated (UNEX), Soxhlet-extracted (EX),
618 or extracted + decarbonated (DECA). Data from C. Johnson, WHOI, 1996.

(wt%)	UNEX	EX*	DECA
C	12.1	11.0	6.23
H	0.38	0.25	1.24
N	0.18	0.17	0.74
S	0.37	<0.2	2.3

619 *Used in the experiment.

Table 2

Concentration of aqueous species during heating of Soxhlet-extracted Eagle Ford shale at 200 to 350 °C and 350 bar in the presence of saline D₂O fluid.

Time Pt #	Time (h)	H ₂ (μmol/kg) ^a	CH ₄ (μmol/kg)	ΣCO ₂ (mmol/kg)	CH ₄ /ΣC ₂₋₄ ^b	ΣH ₂ S (mmol/kg)	pD (25 °C) ^c
<i>Experiment begun with 52.6 g of fluid at temperature of 200 °C</i>							
1	19	BDL (<10)	1.2	4.8	0.78		
2	164	BDL (<10)	3.8	10.8	1.06		
<i>Temperature raised to 300 °C</i>							
3	191	773	8.7	21.9	1.68		
4	284	183	235	45.8	1.30		
5	427	290	396	65.5	1.09		
<i>Injected ~18.3 g starting fluid and raised temperature to 325 °C</i>							
6	451	353	319	44.7	1.06		
7	598	586	825	45.3	0.96		
<i>Raised temperature to 350 °C</i>							
8	617	718	1.32 × 10 ³	54.4	1.01		
9	836	821	3.47 × 10 ³	51.2	1.06	18.0	5.90

Analytical uncertainties (2s) are ±2 °C for T; ±5% for H₂, ΣCO₂, CH₄, and C₂ to C₄ hydrocarbons, ±2% for ΣH₂S; and ±0.05 units for pD. Concentrations are molar quantities per kg fluid.

^a Determined from thermal conductivity response calibrated against a known H₂ standard. As discussed in the text, the listed hydrogen concentration was estimated as if all hydrogen were in the form of H₂. In reality, differences in the TCD response of H₂ and D₂ mean that the actual concentration of molecular hydrogen (likely primarily D₂) are somewhat different than those listed.

^b Calculated as the molar ratio of methane to the sum of ethane, propane, isobutane, and *n*-butane.

^c The listed pD value was calculated from pH measured with a glass electrode: pD = pH_{measured} + 0.41 (Glasoe and Long, 1960).