## Selected Heats of Formation, Entropies, and Free Energies

substance	$\Delta H_{\mathrm{f}}^{\circ}\left(\mathrm{kJ/mol}_{\mathrm{rxn}}\right)$	$S^{\circ}(J/K^{\bullet}mol)$	$\Delta G^{\circ}_{f} (kJ/mol_{rxn})$
H(g)	218.2	114.6	203.2
C(g)	716.682	158.096	671.257
N(g)	470.4	153.3	455.5
O(g)	249.4	160.95	230.1
C(s, graphite)	0	5.69	0
C(s, diamond)	1.90	2.4	2.87
$H^+(aq)$	0	0	0
$OH^-(aq)$	-229.94	-10.5	-157.3
$H_2(g)$	0	131.0	0
$N_2(g)$	0	191.5	0
$O_2(g)$	0	205.0	0
$CH_4(g)$	-74.85	186.2	-50.8
$CO_2(g)$	-393.5	213.6	-394.4
$H_2O(g)$	-241.8	188.7	-228.6
$H_2O(l)$	-285.5	69.9	-237.2
HCl(g)	-92.307	186.908	-95.299
$NH_3(g)$	-46.11	192.45	-16.45
$NH_4Cl(s)$	-314.43	94.6	-203.87
$NO_2(g)$	33.85	240.6	51.8
$N_2O_4(g)$	9.66	304.3	98.29