

quantum reverse-bridge ablation (held-out sample)

category	description	target	generated
parameterised	This is a parameterised quantum circuit wi	q[1]; ry(2.7483749808876965) q[0]; sxdg q[1];	818) q[9];recrx(0.04413440483075049) q[6]
parameterised	This is a parameterised quantum circuit wi	ccz q[6],q[4],q[7]; [4],q[0]; sdg q[1];	2];ry(4.0733927803011656) q[2];tdg q[6];h
highly-entangled	This is a highly-entangled quantum circuit	tdg q[2]; iswap q[1],q[4];	q[6];u(0.744546648287117,2.50457647329
highly-entangled	This is a highly-entangled quantum circuit	h q1; cx q0,q1; rz(param0) q1; cx q0,q1; h gate xx_mi	q1; ry(param0) q1; sdg q0; sdg q0; sx q1; s
measurement-heavy	This is a measurement-heavy quantum circ	3) q[0],q[1]; xx_plus_yy(0.2812769118258048,5.56196 31087) q[1],q[2];	7),q[2];rcccx q[0],q[7],q[1],q[3];c3sqrtx q[0],
measurement-heavy	This is a measurement-heavy quantum circ	crz(4.821809512175018) q[0],q[1]; ccx q[2];	4];ccz q[8],q[7],q[5];cp(2.918380537368832
entangling	This is a entangling quantum circuit windo	x q2,q3; tdg q3; h q3; } gate xx_plus_yy(param0,param1) q0,q	ry((param0) q0; sdg q1; sdg q0; sx q1; sdg c
entangling	This is a entangling quantum circuit windo	0524109945011) q[8],q[4], rcccx q[1],q[3],q[2],q[6]; ccx q[0];	q[1];id q[5];z q[1];rz(3.07403834373802041
single-qubit	This is a single-qubit quantum circuit windo	.5000566246675264,3.163042162042089 u2(2.382597760033 q[0];	1];crz(0.6230123252646397) q[2],q[4];c3sqr
single-qubit	This is a single-qubit quantum circuit windo	id q[1]; ryy(2.351954592710642) q[1],q[0]; rzx(4.0107	4];crzz(2.3284230663465185,0.1762537784