

Lecture 18: The Reactor as a Chemical Plant (Xenon and Samarium Poisoning)

CBE 30235: Introduction to Nuclear Engineering — D. T. Leighton

February 23, 2026

Introduction: The "Ash" is Alive

So far, we have treated fission products as generic "waste" or "precursors." However, the split fragments of Uranium are rarely stable isotopes. They are highly radioactive and undergo their own decay chains, often transmuting into elements with vastly different nuclear properties.

From a neutronics perspective, most fission products are irrelevant—they have small absorption cross-sections. But a select few have **massive** hunger for neutrons. We call these **Neutron Poisons**.

The most significant of these is **Xenon-135**.


- σ_a for U-235: ≈ 600 barns.
- σ_a for Boron-10 (Control Rods): $\approx 3,800$ barns.
- σ_a for Xenon-135: $\approx 2,600,000$ **barns**.

A tiny trace of Xenon has the same stopping power as a bank of control rods.


1 The Chemical Engineering Perspective: Series Reactions

For the Chemical Engineers in the room, the behavior of Xenon-135 is a classic example of **Series Reaction Kinetics** with a competing consumption path.

Consider a chemical reaction series:



where B is also consumed by a side reaction with a reactant R :



In our Nuclear Reactor:

1. **Species A (Iodine-135):** The "Reactant" or Precursor. It is produced by fission and decays into B.
2. **Species B (Xenon-135):** The "Intermediate." This is the poison we care about.
3. **Reaction k_2 (Decay):** Xenon naturally decays into Cesium (Species C).

4. **Reaction k_3 (Burnout):** Xenon captures a neutron (Reactant R) and transmutes into Xenon-136 (Species D), which is harmless.

The concentration of the Poison (B) depends on the balance between production from A , natural decay to C , and "burnout" to D .

2 The Iodine-Xenon Chain

Let's formalize the physics.

- **Iodine-135 (I):**

- Produced directly from fission ($\gamma_I \approx 0.061$).
- Half-life: 6.6 hours.
- Decays into Xenon-135.

- **Xenon-135 (X):**

- Produced from Iodine decay AND directly from fission ($\gamma_X \approx 0.003$).
- Half-life: 9.1 hours.
- Removed by decay OR by neutron capture ($\sigma_a \approx 2.6 \times 10^6$ b).

2.1 The Balance Equations

We write the mass balances (differential equations) for Iodine (I) and Xenon (X):

$$\frac{dI}{dt} = \gamma_I \Sigma_f \phi - \lambda_I I \quad (1)$$

(Rate of Change = Production from Fission - Decay of Iodine)

$$\frac{dX}{dt} = \underbrace{\lambda_I I}_{\text{Source from I}} + \underbrace{\gamma_X \Sigma_f \phi}_{\text{Direct Fission}} - \underbrace{\lambda_X X}_{\text{Natural Decay}} - \underbrace{\sigma_a \phi X}_{\text{"Burnout"}} \quad (2)$$

3 Equilibrium Xenon (Steady State)

At steady state (constant power for > 50 hours), the derivatives are zero ($\frac{d}{dt} = 0$). Solving for equilibrium Xenon X_∞ :

$$X_\infty = \frac{(\gamma_I + \gamma_X) \Sigma_f \phi}{\lambda_X + \sigma_a \phi} \quad (3)$$

The key insight here is the **Burnout Term** ($\sigma_a \phi X$). At high power (high ϕ), neutrons destroy Xenon almost as fast as it is made, keeping the level manageable.

4 From Atoms to Dollars: The Reactivity Worth

Knowing the number of Xenon atoms (X) is academic; operators need to know the **Reactivity Worth** (ρ_{Xe}). Since Xenon acts as a parasitic absorber, it increases the macroscopic absorption cross-section of the core, decreasing the thermal utilization factor (f).

The reactivity penalty is approximately:

$$\rho_{Xe} \approx -\frac{\Sigma_a^{Xe}}{\Sigma_a^{Fuel}} = -\frac{X\sigma_a^{Xe}}{\Sigma_a^{Fuel}} \quad (4)$$

At High Flux (typical power reactor), the equilibrium Xenon worth saturates at approximately:

$$\rho_{Xe,\infty} \approx -0.05 \quad (\text{or } -5\% \Delta k/k)$$

This is huge! It means we must load 5% extra fuel just to overcome the Xenon that will exist during steady-state operation.

Historical Note: The Hanford Mystery

This effect was discovered during the Manhattan Project. On Sept 26, 1944, the B-Reactor (the first plutonium production reactor) was started up. After reaching full power, it mysteriously died a few hours later. Enrico Fermi and John Wheeler correctly identified Xenon-135 as the culprit. Fortunately, the engineers had over-designed the core, allowing them to add extra fuel to override the poison. (See Rhodes, [Hanford and History: B Reactor's 60th Anniversary](#)).

5 Shutdown Behavior: The "Iodine Pit"

This is the most dangerous aspect of Xenon. To understand it, we must look at the "reservoirs" of Iodine and Xenon that exist at the moment of shutdown ($t = 0$).

5.1 Step 1: The Initial Inventories ($t = 0$)

Assume the reactor has been operating at a constant flux ϕ_0 for a long time (equilibrium).

- **Equilibrium Iodine (I_0):**

$$I_0 = \frac{\gamma_I \Sigma_f \phi_0}{\lambda_I}$$

Note that I_0 is directly proportional to flux (ϕ_0).

- **Equilibrium Xenon (X_0):**

$$X_0 = \frac{(\gamma_I + \gamma_X) \Sigma_f \phi_0}{\lambda_X + \sigma_a \phi_0}$$

At high flux ($\sigma_a \phi_0 \gg \lambda_X$), the ϕ_0 terms in the numerator and denominator cancel out, meaning X_0 **saturates** to a constant value.

Key Insight: As flux increases, the Iodine reservoir (I_0) grows linearly, but the Xenon reservoir (X_0) stays roughly constant. This creates a massive imbalance: a huge "tank" of Iodine ready to pour into a small "bucket" of Xenon the moment the reactor shuts down.

5.2 Step 2: The Shutdown Equations ($t > 0$)

At $t = 0$, we SCRAM the reactor ($\phi \rightarrow 0$). The differential equations simplify because the production terms (fission) and burnout terms ($\sigma_a \phi X$) vanish.

Iodine Evolution:

$$\frac{dI}{dt} = -\lambda_I I \implies I(t) = I_0 e^{-\lambda_I t}$$

The Iodine simply decays away, feeding the Xenon.

Xenon Evolution:

$$\frac{dX}{dt} = \lambda_I I(t) - \lambda_X X(t)$$

Substituting $I(t)$:

$$\frac{dX}{dt} + \lambda_X X = \lambda_I I_0 e^{-\lambda_I t}$$

This is a first-order linear ODE. Solving for $X(t)$ with the initial condition $X(0) = X_0$:

$$X(t) = X_0 e^{-\lambda_X t} + \frac{\lambda_I I_0}{\lambda_I - \lambda_X} \left(e^{-\lambda_X t} - e^{-\lambda_I t} \right) \quad (5)$$

5.3 Step 3: Analyzing the Terms

The equation tells a story of two competing effects:

1. **The Decay Term** ($X_0 e^{-\lambda_X t}$): The initial Xenon tries to decay away. This term is strictly decreasing.
2. **The Source Term (The "Pit")**: The second term represents the new Xenon being born from the decay of the massive Iodine inventory (I_0). Because $\lambda_I > \lambda_X$ initially (production is faster than decay), this term causes the concentration to **rise**.

The sum of these two terms creates the characteristic "hump" shape. The Xenon concentration rises, reaches a maximum (the "Peak"), and only then begins to fall.

6 The Magnitude of the Problem: Flux Dependence

Why is this a problem for power reactors but not for low-power research reactors?

Let's look at the ratio of the peak Xenon (X_{max}) to the equilibrium value (X_0). Substituting high-flux approximations into our equation, we find that the peak depends heavily on the pre-shutdown flux ϕ_0 :

Flux (ϕ_0)	Reactor Type	Peak Ratio (X_{max}/X_0)
10^{12} n/cm ² s	University Research	≈ 1.05 (Barely noticeable)
10^{13} n/cm ² s	Early Submarines	≈ 1.50 (Significant)
2×10^{14} n/cm ² s	Modern PWR	≈ 12.0 (Massive)

Table 1: The "Penalty" of High Flux. In a modern power reactor, the Xenon concentration after shutdown can jump to **12 times** its steady-state value.

Because the negative reactivity is proportional to X , a factor of 12 increase means a massive insertion of negative reactivity—often far more than the control rods can overcome to restart the reactor. This forces the "Dead Time" of 24-30 hours where startup is physically impossible.

7 Operational Strategy: The "Dead Time" and Restart

Handling Xenon is one of the primary challenges for a reactor operator.

7.1 1. The "Rod Dance" (Restarting with Xenon)

If you restart the reactor while Xenon is present (e.g., 4 hours after shutdown), you must perform a counter-intuitive control rod maneuver:

1. **The Pull:** To achieve criticality, you must withdraw rods significantly *further* than normal to compensate for the poison.
2. **The Burn:** As power rises, the neutron flux immediately starts "burning" the Xenon away ($Xe + n \rightarrow Xe_{136}$).
3. **The Push:** As the poison vanishes, positive reactivity is added to the core. If the operator does nothing, power will run away. Therefore, as power rises, the operator must **insert** control rods to maintain stability.

7.2 2. The Naval Scenario: "Towed Home"

For naval reactors, the Iodine Pit poses a tactical risk. If a ship SCRAMs (e.g., due to a cooling issue or other fault), the "Xenon Clock" starts ticking.

- If the crew fixes the issue and restarts quickly (within $\approx 1 - 2$ hours), they can burn the Xenon out.
- If the repair takes too long, the Xenon builds up to a peak where its negative reactivity may exceed the reactor's total excess reactivity (especially near the end of core life).
- **Result:** The reactor is "**Xenon Precluded.**" It cannot be restarted, even with all rods removed. The ship loses propulsion for ≈ 24 hours until the Xenon naturally decays. In a combat scenario, this leaves the vessel "Dead in the Water".

8 Case Study: The Role of Xenon in the Chernobyl Disaster

While the RBMK reactor's positive void coefficient (Lecture 17) was the "gun," **Xenon poisoning** was the reason the operators pulled the trigger.

1. **The Power Dip:** In preparation for a test, the operators reduced power. However, a control system error caused the power to plummet to near-zero (< 30 MW).
2. **The Poisoning:** The Iodine inventory from the previous high-power operation immediately began decaying into Xenon. The Xenon concentration spiked (The Iodine Pit), driving reactivity deep into the negative.
3. **The Fatal Decision:** The operators tried to raise power back to the test level (700 MW), but the Xenon was "holding it down." To compensate, they withdrew virtually every control rod in the core.
4. **The Trap:** This left the reactor in a precarious state:

- The rods were physically out (no immediate "brakes").
 - The Xenon was the only thing holding the reactor sub-critical.
5. **The Burnout:** When they finally managed to raise power, the neutron flux began to burn the Xenon away ($Xe + n \rightarrow Xe_{136}$). This added positive reactivity *just* as the steam voids began to form (Positive Void Coefficient). The combination led to the prompt critical excursion.


Safety Lesson: Never fight a Xenon transient with control rod withdrawal beyond your safety margin. If you are "poisoned out," you must accept the delay and wait for the decay.

9 Samarium-149: The Stable Poison

If Xenon behaves like a dynamic wave, Samarium-149 (^{149}Sm) behaves like an **Integrator**. It represents a long-term, cumulative reactivity penalty that fundamentally differs from Xenon because Samarium is stable.

9.1 Production and Burnout Mechanisms

Samarium-149 is a strong, stable neutron absorber produced through the decay chain:



- **Precursor:** Promethium-149 (^{149}Pm), which has a half-life of ≈ 53 hours.
- **The Poison:** Samarium-149 is a stable isotope ($\lambda_{Sm} = 0$). It possesses a very large thermal neutron absorption cross-section ($\sigma_a \approx 41,000$ barns).
- **Burnout:** The only way ^{149}Sm is removed from the core is by absorbing a neutron to become Samarium-150 (^{150}Sm). Because ^{150}Sm has a comparatively tiny cross-section (≈ 100 barns), this absorption effectively eliminates the poison.

Unlike xenon, Samarium-149 does not decay radioactively. Its behavior is governed by the balance between production from Promethium decay and neutron absorption ("burnout"). This leads to two important and somewhat counterintuitive results:

- The *equilibrium* samarium concentration is independent of reactor power.
- The *shutdown increase* in samarium is modest, even after long outages.

9.2 Equilibrium Promethium Concentration

Promethium-149 is produced directly from fission with cumulative yield γ_{Pm} and decays to Samarium with decay constant λ_{Pm} . At steady state, production balances decay:

$$\gamma_{Pm}\Sigma_f\phi = \lambda_{Pm}Pm_0 \quad (6)$$

yielding the equilibrium Promethium concentration:

$$Pm_0 = \frac{\gamma_{Pm}\Sigma_f\phi}{\lambda_{Pm}} \quad (7)$$

Key point: The Promethium inventory scales *linearly* with neutron flux.

9.3 Equilibrium Samarium Concentration

Samarium-149 is produced solely by Promethium decay and destroyed solely by neutron absorption. At equilibrium:

$$\lambda_{Pm} Pm_0 = \sigma_{a,Sm} \phi Sm_0 \quad (8)$$

Substituting for Pm_0 and canceling the flux gives:

$$Sm_0 = \frac{\gamma_{Pm} \Sigma_f}{\sigma_{a,Sm}} \quad (9)$$

Critical Insight: The equilibrium Samarium concentration is **independent of neutron flux**. Whether the reactor operates at 10% or 100% power, it will asymptotically reach the same Samarium worth.

Typical values:

- $\sigma_{a,Sm} \approx 4 \times 10^4$ barns (thermal)
- Equilibrium reactivity penalty: -0.6% to -1.0% $\Delta k/k$

9.4 Shutdown Behavior: Samarium Growth

Upon shutdown, the neutron flux collapses:

$$\phi \rightarrow 0$$

and Samarium burnout ceases entirely. However, the existing Promethium inventory continues to decay into Samarium with half-life:

$$t_{1/2}^{Pm} = 53 \text{ hours}, \quad \lambda_{Pm} = \frac{\ln 2}{53 \times 3600} \approx 3.6 \times 10^{-6} \text{ s}^{-1}$$

The time-dependent Samarium concentration after shutdown is:

$$Sm(t) = Sm_0 + Pm_0 \left(1 - e^{-\lambda_{Pm} t}\right) \quad (10)$$

After a sufficiently long shutdown:

$$Sm_{\max} = Sm_0 + Pm_0 \quad (11)$$

9.5 Why the Shutdown Increase is Modest

The magnitude of the shutdown increase depends on the ratio:

$$\frac{Pm_0}{Sm_0} = \frac{\sigma_{a,Sm} \phi}{\lambda_{Pm}} \quad (12)$$

Using representative LWR values:

$$\begin{aligned} \sigma_{a,Sm} &= 4 \times 10^4 \times 10^{-24} = 4 \times 10^{-20} \text{ cm}^2 \\ \phi &\approx 1 \times 10^{14} \text{ n/cm}^2 \cdot \text{s} \end{aligned}$$

$$\sigma_{a,Sm} \phi \approx 4 \times 10^{-6} \text{ s}^{-1} \quad (13)$$

Comparing with λ_{Pm} :

$$\frac{Pm_0}{Sm_0} \approx \frac{4 \times 10^{-6}}{3.6 \times 10^{-6}} \approx 1.1 \quad (14)$$

At full power, the Promethium and Samarium inventories are comparable.

Thus, after a long shutdown:

$$Sm_{\max} \approx 2 Sm_0$$

The additional reactivity penalty is typically only:

$$-0.3\% \text{ to } -0.6\% \Delta k/k$$

9.6 Flux Dependence of the Shutdown Effect

Because $Pm_0 \propto \phi$:

- **High flux reactors:** noticeable but limited Samarium growth
- **Low flux reactors:** negligible post-shutdown increase

At low flux ($\phi \sim 10^{13} \text{ n/cm}^2\cdot\text{s}$), $Pm_0 \ll Sm_0$ and shutdown effects are barely observable.

9.7 Comparison with Xenon

Samarium differs fundamentally from Xenon-135:

- Samarium is stable and permanent
- Xenon decays away
- Xenon shutdown peaks can be several times equilibrium
- Samarium shutdown peaks are typically $\lesssim 2\times$ equilibrium

Design Implication: Reactor cores are loaded with sufficient excess reactivity at beginning of life to permanently compensate for equilibrium Samarium and its modest shutdown increase. This unavoidable reactivity penalty is known as the *Samarium Tax*.




Figure 1: **The Consequence of High Flux.** In a low-power reactor (Blue), the Xenon peak is a gentle bump ($\approx 1.2\times$) that never exceeds the control rod capacity. In a high-power commercial reactor (Red), the peak is a massive spike ($\approx 12\times$) because the accumulated Iodine inventory is enormous. This renders the reactor inoperable for nearly two days.