

# Penguin Species Classification Memo

Tran Quoc Bao Dang

2026-06-13

## Contents

|          |                                 |          |
|----------|---------------------------------|----------|
| <b>1</b> | <b>Executive Summary</b>        | <b>2</b> |
| <b>2</b> | <b>Background and Objective</b> | <b>2</b> |
| <b>3</b> | <b>Data Description</b>         | <b>2</b> |
| <b>4</b> | <b>Exploratory Analysis</b>     | <b>2</b> |
| <b>5</b> | <b>Analytical Approach</b>      | <b>2</b> |
| <b>6</b> | <b>Results</b>                  | <b>4</b> |
| <b>7</b> | <b>Limitations</b>              | <b>4</b> |
| <b>8</b> | <b>Recommendation</b>           | <b>4</b> |

# 1 Executive Summary

Dr. Sharma provided measurements from an unidentified penguin specimen with a bill length of 45 mm and a flipper length of 220 mm. Using the Palmer Penguins dataset, I compared these measurements to known Adelie, Chinstrap, and Gentoo penguins. Both visual comparison and statistical analysis suggest that the specimen is most likely a Gentoo penguin. The specimen's flipper length is particularly characteristic of Gentoo penguins. While the classification appears strong, confidence could be improved with additional measurements such as body mass or bill depth.

## 2 Background and Objective

The objective of this project is to identify the most likely species of an unlabeled penguin specimen discovered during a previous expedition. The available measurements are bill length (45 mm) and flipper length (220 mm). Dr. Sharma would like to determine:

1. The most likely species.
2. The confidence in that classification.
3. How the specimen compares visually to known penguin species.

## 3 Data Description

The Palmer Penguins dataset contains measurements from three penguin species: Adelie, Chinstrap, and Gentoo. For this analysis, bill length and flipper length were used because these measurements were available for the unknown specimen. Records with missing values were removed prior to analysis.

After removing records with missing values, the dataset contains 342 penguins across three species: 151 Adelie, 68 Chinstrap, and 123 Gentoo.

## 4 Exploratory Analysis

Figure 1 shows the relationship between bill length and flipper length for the three species. Gentoo penguins generally have longer flippers than Adelie and Chinstrap penguins. The unknown specimen falls near the Gentoo cluster due to its unusually long flipper length of 220 mm.

Table 1: Comparison of the Unknown Specimen with Typical Measurements by Species (mm)

| species          | Avg Bill Length | Avg Flipper Length |
|------------------|-----------------|--------------------|
| Adelie           | 38.8            | 190.0              |
| Chinstrap        | 48.8            | 195.8              |
| Gentoo           | 47.5            | 217.2              |
| Unknown Specimen | 45.0            | 220.0              |

## 5 Analytical Approach

The unknown specimen was compared to known penguin measurements using a statistical classification procedure designed to estimate the most likely species.




Figure 1: Bill length vs. flipper length for known penguin species. The red star marks the unknown specimen (bill length = 45mm, flipper length = 220mm).

## 6 Results

The estimated probabilities for each species are shown in Table 2. Gentoo received the highest probability and was therefore identified as the most likely species. This result is consistent with the visual evidence presented in Figure 1 and the comparison shown in Table 1.

Table 2: Estimated Probability of Species Membership

| Species   | Probability |
|-----------|-------------|
| Adelie    | 0.03        |
| Chinstrap | 0.00        |
| Gentoo    | 99.96       |

The model assigned a 99.96% probability to Gentoo, compared to 0% for Chinstrap and 0.03% for Adelie, making Gentoo the clear frontrunner.

The unknown specimen has a bill length of 45 mm and a flipper length of 220 mm. While its bill length falls between the Chinstrap and Gentoo averages, its flipper length is most consistent with Gentoo penguins, providing strong evidence for that classification.

## 7 Limitations

This classification is based only on two measurements: bill length and flipper length. Including body mass, bill depth, sex, island, or other biological information could improve confidence. Because the specimen is preserved and from a prior expedition, measurement error or preservation effects may also affect the result.

## 8 Recommendation

Based on both visual inspection and statistical classification, the specimen should be provisionally classified as a Gentoo penguin. The evidence supporting this classification is strong because the specimen’s flipper length is highly characteristic of Gentoo penguins. If additional measurements such as body mass or bill depth become available, the classification should be re-evaluated to further increase confidence.