11 November 2026 Mathematical comparison for revision

The common web and its individual coefficients

Goertzen–Lacabanne–Tubbenhauer–Vaz (A) and
Bowman–de la Fuente–Libedinsky–Plaza (B)

Verdict

The constructions agree before a specific canonical correction. Start with B's uncorrected three-parabolic product $\mathcal{BS}_m$. Replace each of B's two strings of simple reflections by the normalized symmetrizer of the parabolic it generates. The resulting character equals A's entire six-stage web, in every rank and in every standard coordinate. The proof compares two finite flag counts after a common annular merge and split. One counts hyperplanes containing the sum of two subspaces. The other counts planes in their intersection containing a fixed line. Their projective spaces have the same dimension.

B's actual proof corrects its seed to $\mathbf{C}_{X_0}$, then performs the simple-reflection multiplications with further corrections. Its endpoint is exactly the multiplicity-one canonical summand of A indexed by $A_{m;m}$, after inversion. The ith canonical constituent of B's raw product supplies exactly the ith distinguished summand of A after averaging. In particular B's corrected seed supplies the diagonal. The shared gallery identifies where the two arguments begin to calculate different quantities.

A computes the sum $\binom{m+1}{2}$. B computes its final entry m, on the same highest-weight module after scalar extension. B also identifies the first nonzero rational and quantum cohomology in a sequence of degrees, and obtains finite-group lower bounds in every fixed positive degree. The comparison also resolves A's conjecture about the interior of the row. At fixed m, complete the first j canonical constituents of B's raw product. At the full-spherical standards their sum has the same coefficients through degree one as A's web in ambient parameter j. This gives $\sum_{i=1}^{j} \mu_{i;m} = \binom{j+1}{2}$ for every $j \le m$. Taking consecutive differences proves $\mu_{j;m} = j$. Neither v1 contains this partial-sum calculation.

For the revision, *concurrent* is justified. B's submitted bibliography contains the exact title, four authors and project URL. GitHub records public pushes containing the construction and conjectural diagonal value before either arXiv submission. The order in which anyone encountered these materials and found the endpoint or proof remains undocumented. The flag identity makes a precise question about the shared construction possible. It cannot date its discovery.

Versions and scope

A is arXiv:2611.09114v1, `main.tex` with its embedded bibliography.

B is arXiv:2611.11677v1, `Mu_parabolic_complete.tex` and `bibliography.bib`. Source references use the numbering in these two PDFs. The proof of the individual formula is in Section 4, on pages 8–9. The records and calculations accompany this report. No source for the third proof was supplied.

1 Conventions and the common canonical element

Put $r = m,\ N = 2m+1,\ I = \{s_1, \dots, s_{N-1}\},\ w_I = w_0$, and $V = I \setminus \{s_m, s_{m+1}\}$. Both papers compose periodic permutations by $(xy)(i) = x(y(i))$. Right multiplication by s_i swaps positions $i, i+1$. For s_0 the new first and last entries are $w(N) - N, w(1) + N$. The increasing window selects the shortest representative in W/W_I; the increasing inverse window selects the shortest representative in $W_I\backslash W$. A allows the extended group; its construction has degree zero.

Parameters. Both use $H_s^2 = 1 + (v^{-1} - v)H_s,\ C_s = H_s + v$. With $T_w = v^{-\ell(w)}H_w$, the coefficient of H_y in C_w is $h_{y,w}(v) = v^{\ell(w)-\ell(y)}P_{y,w}(v^{-2})$. A uses $q = v^{-2}$ in its flag counts. B sets $q = v^2$ for its Gaussian polynomials in §3.1. Gaussian reciprocity contributes the factor $v^{-2b(a-b)}$ when changing $\begin{bmatrix} a \\ b \end{bmatrix}_{v^{-2}}$ to $\begin{bmatrix} a \\ b \end{bmatrix}_{v^2}$. For $L = \ell(w_P)$, A's C_P, h_P are B's C_{w_P}, π_P, respectively:

$$C_P = \sum_{z \in W_P} v^{L-\ell(z)} H_z, \qquad h_P = \sum_{z \in W_P} v^{L-2\ell(z)}, \qquad C_P^2 = h_P C_P.$$

Sides and bases. A's web maps the right ideal $C_I\mathcal{H}$ to $C_V\mathcal{H}$. Evaluate on C_I, then apply the linear anti-involution $H_w^* = H_{w^{-1}}$, which fixes v and reverses products. The result lies in B's right ideal $C_I\mathcal{H}$, with right V-spherical character. For an A double coset $D = W_V d W_I$ with longest element z, its standard map goes to $v^{\ell(z)} \sum_{w \in D^{-1}} T_w$, and its canonical map to $C_{z^{-1}}$. B's bases are $M_b = C_I H_b$ and $\mathbf{C}_b = C_{w_I b}$, for $b \in {}^I W$, and $[M_a]\mathbf{C}_b = h_{w_I a, w_I b}$. At A's splitter the inverted standard is $M_e = C_I$. Thus for *all* j, with $b = u_{j;m}^{-1}$,

$$\big[[B_m]_v\big]\theta_{A_{j;m}} = [M_e]\mathbf{C}_b = v^{D_{j;m}} P_{w_0, C_{j;m}}(v^{-2}), \qquad D_{j;m} = 2j(N-j) - 2(m-j) - 3.$$

Here $C_{j;m} = u_{j;m} w_0$ is a permutation, while C_w denotes a basis vector. See A Definitions 2D.1–3, 2E.1–2, Lemma 2E.7, Proposition 2E.9 and §3B; B §2.1, Lemma 2.1.

The diagonal. Write $\nu = (1^m, 0, (-1)^m)$, $\sigma = (m, m+2)$, and $t_\nu(i) = i + N\nu_i$ on $1 \le i \le N$. In A's increasing window at $j = m$ the ordered residues are $1, \dots, m-1, m+2;\ m+1;\ m, m+3, \dots, N$ with shifts $-1, 0, 1$ on the respective groups. Therefore $u_{m;m} = \sigma t_{-\nu}$. Its inverse $t_\nu \sigma$ has window $[N+1, \dots, N+m-1, 1-m, m+1, N+m, 2-m, \dots, 0]$, precisely B's β_m. As $w_I \beta_m = C_{m;m}^{-1}$, inversion identifies the ordinary KL polynomials and the full Hecke vectors satisfy $(\theta_{A_{m;m}}(C_I))^* = \mathbf{C}_{\beta_m}$. The dot action gives $\beta_m \cdot_p 0 = p\nu + \sigma\rho - \rho = p\nu - 2(\alpha_m + \alpha_{m+1})$, hence

$$\lambda_{m;m}(p) = 2\varpi_{m-1} + (p-2)(\varpi_m + \varpi_{m+1}) + 2\varpi_{m+2}.$$

B includes the factor p in its translation label. See A Lemma 6A.2 and B Lemma 2.16.

2 The two galleries in one Schur algebroid

Write $C_P = C_{w_P}$, $\pi_P = h_P$, $\Pi_n = v^{-\binom{n}{2}}[n]_{v^2}!$, $[n] = \Pi_n/\Pi_{n-1}$, and $\eta = [2] = v + v^{-1}$. For a consecutive interval $[a, b]$ in a parabolic set we mean $s_a, \dots, s_b$. In the right spherical module the sequence of merge targets $P_1, \dots, P_d$ represents

$$D(P_1, \dots, P_d) = \frac{C_I C_{P_1} \cdots C_{P_d}}{\prod_{i=1}^{d} \pi_{P_{i-1} \cap P_i}}, \qquad P_0 = I.$$

Between consecutive targets one splits to their intersection and merges to the next target. The splits are inclusions. Before inversion the merge from X to $Y \supset X$ is left multiplication by C_Y/π_X on the appropriate image. Inversion reverses the product. Thus every quotient below is integral on its specified domain (A, Lemma 3B.1; B, Definition 3.2 and Lemma 3.3).

Symbol	Generators	Position blocks or symmetric-group type
I	$[1, N-1]$	(N)
J	$S \setminus \{s_m, s_{m+1}\}$	$(2m, 1)$ across the cut
V	$I \cap J$	$(m, 1, m)$
K	$[2, N-2]$	$(1, 2m-1, 1)$
L	$K \cap J = K \cap V$	$(1, m-1, 1, m-1, 1)$
K'	$I \setminus \{s_1, s_{m+1}\}$	$(1, m, m)$
E	$[3, N-1]$	$(1, 1, 2m-1)$
T	$J \cap K' = V \cap K'$	$(1, m-1, 1, m)$
D'	$E \cap K'$	$(1, 1, m-1, m)$

In particular $\mathcal{R}_m = D(J, K, V)$ and $\mathcal{BS}_m = D(J, K', E)$. Both start with $F = C_I C_J/\pi_V$; this is itself canonical by B, Lemmas 3.1 and 3.19. The first two stages of A's six-stage web are exactly $I \supset V \subset J$. For the general $\mathcal{BS}_j$ replace J, K' by $J_j = S \setminus \{s_j, s_{j+1}\}$ and $K_j = I \setminus \{s_1, s_{j+1}\}$. The cut is crossed by merging the final $N-j-1$ packet with the first j packet; the singleton stays separate. The intermediate splittings have blocks $(j, 1, N-j-1)$, $(1, j-1, 1, N-j-1)$, and $(1, 1, j-1, N-j-1)$, apart from the degeneracy at $j = 1$, where the last two overlaps both equal $K_1 = E$. For $\mathcal{CBS}_s = D(P_s, E)$, $P_s = S \setminus \{s_s\}$; one joins the packets $(s, N-s)$ across the cut, then splits to $(1, 1, s-2, N-s)$ and merges the last two packets, omitting zero packets. These rotations and their inverse undo one another in determinant degree. Explicitly $C_{J_j} = \tau^{-(N-j-1)} C_{[1,N-2]} \tau^{N-j-1}$ and $C_{P_s} = \tau^{-(N-s)} C_I \tau^{N-s}$ (A, Lemma 2C.9).

For $j \geq 2$ the three overlaps for $\mathcal{BS}_j$ have factors $\Pi_j \Pi_{N-j-1}$, $\Pi_{j-1} \Pi_{N-j-1}$, $\Pi_{j-1} \Pi_{N-j-1}$; for $j = 1$ these same scalar formulas hold. For $\mathcal{CBS}_s$ they are $\Pi_s \Pi_{N-s}$ and $\Pi_{s-2} \Pi_{N-s}$. A's three factors are $\Pi_m^2, \Pi_{m-1}^2, \Pi_{m-1}^2$. The length contribution at an individual merge is $\ell(w_{P_i}) - \ell(w_{P_{i-1} \cap P_i})$.

An exact identity before canonical correction

Set $Q_1 = [2, m]$, $Q_2 = [1, m-1]$ and let T_1, T_2 denote right multiplication by $C_{Q_1}/\Pi_{m-1}, C_{Q_2}/\Pi_{m-1}$, respectively, on the indicated spherical subspaces. Then

$$\boxed{\mathcal{R}_m = D(J, K', E, K', V) = \mathcal{BS}_m T_1 T_2.} \qquad (*)$$


The maximal moving words in these two normalized parabolic factors are exactly $s_2 \cdots s_m$ and $s_1 \cdots s_{m-1}$ from B, Lemma 2.4. The equality holds for the full standard and canonical expansions.

Proof. Refine J to V in both galleries. The first overlaps are $L \subset V$ and $T \subset V$, so this factors an inclusion. Everything after this refinement is a finite flag calculation. Fix the two end flags $U \subset U'$ and $W \subset W'$ of dimensions $m, m+1$ in an N-space over $\mathbb{F}_q$. In the V, K, V gallery the K-flag is a line $a \subset U \cap W$ and a hyperplane $H \supset U' + W'$. In the V, K', E, K', V gallery the two K'-flags are $(a, U'), (a, W')$, and the E-flag is (a, D_2) with $a \subset D_2 \subset U' \cap W'$, $\dim D_2 = 2$. All refinements are uniquely determined by these data. If $d = \dim(U' \cap W')$, then

$$\dim((U' \cap W')/a) = d - 1 = N - \dim(U' + W').$$

For each a, the numbers of two-planes and hyperplanes are therefore equal. Each gallery counts $[\dim(U \cap W)]_q[d-1]_q$ points, where $[k]_q = 1 + \cdots + q^{k-1}$ and $[0]_q = 0$.

The normalization shifts agree. Writing $b_n = \binom{n}{2}$, the first tail contributes $b_{2m-1} + 2b_m - 4b_{m-1}$, and the second contributes $b_{2m-1} + 6b_m - 4(b_m + b_{m-1})$. The two galleries have the same input and output flags, so the orbit-dependent standard shifts also agree. Equality over every finite field gives the generic Schur identity. Finally $K' = Q_1 \times [m+2, N-1]$ and $V = Q_2 \times [m+2, N-1]$. The common factor lies in E, commutes with Q_1, Q_2, and cancels one Π_m from each of $\pi_{D'} = \pi_T = \Pi_m \Pi_{m-1}$. This gives $(*)$. □



Common affine part | Line and hyperplane | Line and plane

Cut-open webs at $m = 3$, read upwards. Stack the common part below either tail and identify each pair of red dashed sides. Blue strands have label one. The label-six strand in the common part straddles the cut. The vertices are the normalized splits and merges of A 2D.4.

Multiplying the $2m-2$ individual simple KL elements instead of the two normalized parabolic factors gives a different test. At the identity coordinate, multiplication by $T_1 T_2$ acts by $[m]^2$, whereas the individual factors act by η^{2m-2} (B, (3.4)). Since $\varepsilon(\mathcal{BS}_m) \neq 0$, evaluation at $v = 1$ distinguishes the two products by $m^2 \neq 4^{m-1}$ for $m \geq 3$. Averaging further by C_V/π_V preserves this coordinate. At $m = 2$ each parabolic factor is already one simple KL element, so the raw products agree.

What the corrected seed contains

Write $b_j = u_{j;m}^{-1}$ and let $[\,\mathbf{C}_b\,]$ denote a coefficient in the spherical canonical basis. For every $1 \le j \le m$,

$$\boxed{[\,\mathbf{C}_{b_j}\,](\mathbf{C}_{X_0}T_1T_2) = \delta_{jm}.} \tag{$**$}$$

At the other end of $(*)$, B, Proposition 3.21, also gives $\mathcal{BS}_m = \sum_{j=1}^{m} \mathbf{C}_{z(j,1,0)}$. In fact its standards are in $\mathbb{Z}[v]$, with constant term 1 precisely at the m indicated orbit maxima and zero elsewhere. Subtracting those canonical elements leaves a bar-invariant vector in $v\mathbb{Z}[v]\{M_c\}$, hence zero. Thus $\mathcal{R}_m = \sum_{j=1}^{m} \mathbf{C}_{z(j,1,0)}T_1T_2$. In particular $(**)$ and A, Lemma 4I.1, place each off-diagonal family-F term, once, in

$$(\mathcal{BS}_m - \mathbf{C}_{X_0})T_1T_2 = \Big(\mathcal{BS}_{m-1} - \eta \sum_{s=2}^{m-1} \mathcal{CBS}_s\Big)T_1T_2.$$

Here $\mathcal{BS}_{m-1}$ is taken in the same period $2m+1$.

Proof of $(**)$. For consecutive generators starting at s_a let $Z_0 = Z$, $Z_i = ZC_{[a,a+i-1]}/\Pi_i$; assume $ZH_s = v^{-1}Z$ for $s \in [a+1, a+i-1]$. The left coset representatives are $g_k = s_a \cdots s_{a+k-1}$, $0 \le k \le i$, so that

$$Z_i = \sum_{k=0}^{i} v^{i-k} Z H_{g_k}, \quad Z_1 = ZC_{s_a}, \quad Z_i = Z_{i-1}C_{s_{a+i-1}} - Z_{i-2} \quad (i \ge 2).$$

The last generator commutes with g_k for $k \le i-2$, giving the last recurrence. In B, Proposition 2.8, this subtracts the preceding cover at each step. The remaining terms $\mathbf{C}_e, \mathbf{C}_{\gamma_k}$ are eigenvectors for every later generator, since the word for γ_k is $BC^kB^{N-2k-1}A^k$. Using $[i+1] = \eta[i] - [i-1]$ gives

$$\mathbf{C}_{X_0}T_1 = \mathbf{C}_{x_{m-1}} + [m-1]\mathbf{C}_e + \sum_{k=1}^{m-2} [m-1-k]\mathbf{C}_{\gamma_k}.$$

The terms after $\mathbf{C}_{x_{m-1}}$ retain the right descent s_{m+1} upon multiplication by T_2, as s_{m+1} commutes with Q_2. Applying the same recurrence to B, Proposition 2.11, yields

$$\mathbf{C}_{x_{m-1}}T_2 = \mathbf{C}_{\beta_m} + \sum_{t=0}^{m-2} [t+1]\mathcal{E}_t.$$

Indeed each canonical term of $\mathcal{E}_t$ is sorted on its first $m+1$ positions by B, Lemma 4.12(ii) and (4.17)–(4.18). Every subsequent generator acts by η, and each such term has right descent s_m. The canonical basis of the right s-spherical subspace consists of the canonical elements with right descent s. The word of b_j is $C^{j-1}B^{m-j}A \,|\, B \,|\, CB^{m-j}A^{j-1}$. The adjacent ascents AB, BC give both missing right descents s_m, s_{m+1} (B, Lemma 4.9). Therefore none of the remainders contains any $\mathbf{C}_{b_j}$. □

The full vector $\mathbf{C}_{X_0}T_1T_2$ still differs from $\mathbf{C}_{\beta_m}$: its identity coordinate has valuation $3-2m$ by B, Proposition 2.6, whereas that of $\mathbf{C}_{\beta_m}$ has valuation one. Formula $(**)$ isolates the diagonal *within family F*. It does not determine the identity coordinates of the other family-F constituents.

3 Where the two coefficient calculations diverge

Both arguments use the same refinement identity. If z is the longest index in a right-parabolic orbit, the coefficient of M_c in a vector invariant under that parabolic is $v^{\ell(z)-\ell(c)}$ times its coefficient at z. One proves this along a single edge by the quadratic Hecke relation. A uses the identity to refine output flags (Lemma 4D.1). B uses it to compute once per orbit (Lemmas 3.11, 3.17 and Proposition 3.21).

Two multiplicity-one expansions. There is a parallel already at the *uncorrected* stage. Write $z(t,a,b)$ for the longest element of B's orbit of type t with indicators a,b for the first two letters. Then

$$\mathcal{BS}_m = \sum_{t=1}^{m} \mathbf{C}_{z(t,1,0)}, \qquad \mathcal{BS}_{m-1} - \eta \sum_{s=2}^{m-1} \mathcal{CBS}_s = \sum_{t=1}^{m-1} \mathbf{C}_{z(t,1,0)}, \qquad \eta = v + v^{-1}.$$

Indeed, B Proposition 3.21 gives least exponents $1,1,0,1$ at feasible maxima with prefixes BB, BA, AB, AA, respectively, for $\mathcal{BS}_m$. The AB maxima occur exactly for $1 \le t \le m$, and their constant terms are one. All other standard coordinates have positive valuation by refinement. Subtract the canonical vectors displayed in the first sum. The difference is bar invariant and every standard coordinate lies in $v\mathbb{Z}[v]$, so it vanishes (B Lemma 3.1). B Proposition 3.5, with $X_0 = z(m,1,0)$, gives the second equality. In its proof the AB constants below the top cancel against $\mathcal{BS}_{m-1}$; in column AA the $-v^{-1}$ from that subtraction is canceled by $+\eta\mathcal{CBS}_t$ at each type $2 \le t < m$. This isolates a single seed.

A establishes multiplicity one *after* completing its web. Lemma 3C.3 fixes flags $I_0 \subset K_0$ of dimensions $r, N-r$ and $F \subset F'$ of dimensions $m, m+1$. For a line $L \subset F$ and a hyperplane $H \supset F'$, put $e = \mathbf{1}_{L \not\subset I_0}$, $d = \mathbf{1}_{H \not\supset K_0}$. The remaining choices of $I' \subset K'$ of dimensions $m, m+1$, between $I_0 + L$ and $K_0 \cap H$, number $\begin{bmatrix} N-2r-e-d \\ m-r-e \end{bmatrix}_q (m+1-r-d)_q$ over a field of size q. Sum against the corresponding line and hyperplane counts (A (3C.2)), substitute $q = v^{-2}$ and multiply by v^{D-d_A}, $D = 2m^2 + 2m - 3$ (A Proposition 3C.4). If $F_A(q)$ is the resulting counting polynomial before normalization, the exponents of a nonzero canonical multiplicity lie between $-E$ and E, $E = 2\deg F_A - (D - d_A)$. A applies positivity after clearing the overlap denominator. With $l = m - r$, the bounds for the six allowed families A–F are respectively $2l-3, l-2, l-1, l-2, l-1, 0$ (Lemmas 4B.1 and 4C.2). At an F-standard $A_{j;m}$, a full-spherical, strong one-incidence or weak one-incidence contribution of rank $j+h$ starts in degree at least $2h+6, 2h+2, 2h+4$, respectively. Larger F-terms contribute only positive powers there. Thus the constant one belongs to $A_{j;m}$ itself (Lemma 4I.1). At a full-spherical standard, in contrast, one-incidence terms with their multiplicities start in degree at least three (Lemma 4H.1), which suffices for the row sum.

The same Gaussian and different subtractions

Let $t = v^2$, $G = \begin{bmatrix} 2m-1 \\ m-1 \end{bmatrix}_t$, and use $(k)_t = 1 + \cdots + t^{k-1}$, $[k] = v^{1-k}(k)_t$. At the identity the raw values are

$$\epsilon(\mathcal{R}_m) = v^{3-2m}(m)_t^3 G, \qquad \epsilon(\mathcal{BS}_m) = v(m)_t G.$$

These instances of the same rectangle follow from the operator identity and the finite eigenvalue, with ratio $[m]^2$ (A Prop. 3C.4, B proof of Prop. 2.6 in §3.5). Even some auxiliary operators agree after merging all outputs. Define $U_j = C_I C_{J_j} C_I / \pi^2_{I \cap J_j}$ and let $V_s = C_I C_{P_s} C_I / \pi^2_{I \cap P_s}$ be A's spherical web (Definition 4E.1). With $p_E(F) = F C_I / \pi_E$,

$$p_E(\mathcal{BS}_j) = [j]^2 U_j, \qquad p_E(\mathcal{CBS}_s) = [s][s-1] V_s.$$

Indeed $C_Q C_I = \pi_Q C_I$ for $Q \subset I$. The two factorial ratios remaining are $\pi_{I \cap J_j} \pi_{K_j} / (\pi_{J_j \cap K_j} \pi_{K_j \cap E}) = [j]^2$ and $\pi_{I \cap P_s} / \pi_{P_s \cap E} = [s][s-1]$. Thus B's seed correction includes scalar multiples of A's cumulative spherical webs after the total merge. If instead one merges first to $Q = I \backslash \{s_1\}$ and appends the move to $I \backslash \{s_f\}$ for $f = m, m+1$, the result for $\mathcal{CBS}_s$ is $[s-1]$ times A's one-line gallery $\mathcal{E}_{s,f}$. This follows from $\pi_{P_s \cap Q} / \pi_{P_s \cap E} = [s-1]$ and A Definition 4F.1.

A takes the differences $C_{Z_s} = V_s - V_{s-1}$, whose standard coefficients are differences of consecutive Gaussians (Lemma 4E.4). It subtracts $g_{m-s} C_{Z_s}$, where $g_l = \sum_{a=0}^{l-2} \binom{a+3}{3} (v^{2l-3-2a} + v^{-(2l-3-2a)})$. To determine these multiplicities, A reverses a convolution of Gaussian polynomials and truncates modulo q^m; its common series $1/((1-q)^3 (q;q)_\infty)$ cancels, leaving $\binom{m+1}{2} q^{m-1}$ (Lemmas 4J.1–4K.1). The surviving linear coefficient at the origin is the sum over family F (Theorem 4L.1).

B applies ϵ to its corrected seed to obtain $h_{X_0} = vG + O(v^{2m+1})$. The first m coefficients are ordinary partition numbers, since partitions of size below m fit the Gaussian rectangle. The first reflection chain satisfies $h_{x_1} = \eta h_{X_0} - 1$ and

$$h_{x_{k+1}} = \eta h_{x_k} - h_{x_{k-1}} - v^k \begin{bmatrix} N-1 \\ k \end{bmatrix}_t \quad (1 \le k \le m-2).$$

The last term is h_{γ_k} (Lemma 3.25). Bruhat covers supply the predecessor; positivity excludes the lower γ_j, paired deletion reduces the surviving multiplicity to one, and a nonidentity seed bound excludes the κ_j (Proposition 2.8, Lemmas 3.24, 4.20, 5.1). If $d_{a,b}$ counts partitions of $a+b$ with at least a parts, the coefficient of v^{a+1+2b} in h_{x_a} is $\sum_{i=0}^{a} d_{i,b}$ for $0 \le b \le m-1-a$ (Lemmas 2.9, 5.2). In particular $h_{x_{m-1}}$ starts with mv^m.

On the second chain the predecessor and remainder have identity-coordinate valuation at least $t'+3$ in the step to $y_{t'}$ (Proposition 2.11 and Lemma 2.12). Of its noncover candidates the ordered terms vanish after rank reduction, the pure terms have sufficiently high valuation by finite ascent bounds, and mixed terms vanish after a rank-two star operation and a descent test (Lemmas 6.1, 6.2, 6.5). This transports m down one degree at each step. The sample individual remainders for $m = 5$ in B are computational. The uniform proof only needs the degree bound. Neither this partition recurrence nor A's Gaussian convolution is a translation of the other.

What was already known. Scott–Xi, arXiv:0909.3394v4, Theorem 3.3, obtained $n+2$ for $\mu(xw_0, vxw_0)$ with a regularly translated lower endpoint. Their Lemma 3.4 gives 2 at fixed source for their particular word in every rank $n \ge 4$; its $n = 4$ instance is the common $m = 2$ example. Section 4 explicitly asks the fixed-source question. Their suggested weight for $n = 4k$ has coefficient $p-2$ at every internal node, so for $n > 4$ it differs from the common diagonal weight considered here. The low-rank coincidence, the fixed-source question and the use of degree bounds in a KL recursion all have this earlier precedent.

4 Every entry of the row

In fact the two calculations together give

$$\boxed{\mu_{j;m} = j \quad (1 \le j \le m).}$$

The additional step is to stop B's *uncorrected parabolic family* at j, while keeping the output boundary and ambient rank of A fixed. We give the argument since the diagonal theorem and the row sum by themselves do not imply this conclusion.

1. Nested canonical sums before completing the web. Hereafter val denotes the lowest exponent of v, with $\operatorname{val} 0 = +\infty$. Let $T = T_1 T_2$ be the normalized finite tail which gives $\mathcal{BS}_m T = \mathcal{R}_m$. Use B's notation for its longest orbit representatives, and put

$$X_i = \mathbf{C}_{z(i,1,0)}, \qquad D_s = \mathbf{C}_{z(s,1,1)}, \qquad \Phi_j = \sum_{i=1}^{j} X_i, \qquad Y_j = \Phi_j T, \qquad Y_0 = 0.$$

Here X_i denotes a vector, not one of B's reflection-chain permutations x_i. We show

$$\mathcal{BS}_j = \Phi_j + \sum_{s=2}^{\min(j,m-1)} a_{js}(v) D_s, \qquad D_s = \mathcal{CBS}_s, \qquad \operatorname{val} a_{js} \ge 2 - N + 2j \quad (a_{js} \ne 0). \tag{1}$$

Only $s \le m-1$ is needed: at $j = m$ every $a_{js} = 0$. For $s \le m-1$ B Proposition 3.21 gives $\mathcal{CBS}_s$ a unique coordinate with nonpositive valuation: at the AA maximum of type s, its constant is one. All others are positive, including nonmaxima by refinement. This proves its canonicality. The $\mathcal{BS}_j$ are right E-invariant, so only maxima of E-orbits can be canonical indices; this is the parabolic canonical basis, or follows by descending triangularity in the invariant subspace. Outside the support listed in B Proposition 3.21 no canonical term can occur: clearing the Poincaré denominator gives a positive product in the ordinary KL basis. Apply the same valuation argument as A Lemma 4B.1 at the maxima. Columns BB, BA have positive valuation. Column AB has valuation zero, so its canonical multiplicity is constant. Column AA gives the bound $2 - N + 2j$ in (1). At an AB maximum of type i the coordinate of $D_s = \mathcal{CBS}_s$, when present, has valuation $N - 2i$ by the same table. Multiplied by a_{js} it starts no lower than $2 + 2(j - i)$. Other AB canonical vectors with constant multiplicities contribute only positive powers there. Thus the constant one belongs to X_i for every feasible $1 \le i \le j$. This proves (1) and in particular $Y_m = \mathcal{R}_m$.

2. What survives from A's canonical expansion. Each Y_j is integral and bar invariant. The domain of T is the right E-invariant module; both normalized factors are integral sums over finite parabolic cosets there (B Lemma 3.3). If $p = (\Pi_{m-1})^2$ is their common denominator, $p(Y_j - Y_{j-1}) = X_j C_{Q_1} C_{Q_2}$ has nonnegative ordinary canonical coefficients. Consequently the cleared coefficient of *each* Y_j is bounded coefficientwise by that of $p\mathcal{R}_m$. Thus Y_j has only A's six families. The valuation of a nonzero multiplicity is at least the corresponding valuation for $\mathcal{R}_m$, by subtracting $\operatorname{val} p$ from the inequality for cleared coefficients. This proves that the degree margins in A Lemmas 4H.1 and 4I.1 apply to Y_j as well. In particular, for each family-F index its multiplicity in Y_j is a constant zero or one, nondecreasing in j, since its multiplicity in $\mathcal{R}_m$ is one. All standard and canonical indices of Y_j have type at most j: (1) and B Proposition 3.21 bound the initial standard support, finite right generators preserve word type, and positivity after clearing excludes any additional canonical support.

For completeness we compute the constant at $A_{j;m}$. A flag for $\mathcal{BS}_j T$ chooses intermediate $U \subset U'$ of dimensions $j, j+1$, with $I_0 \subset U \subset U' \subset K_0$, in the finite quotient of A Definition 3C.1. To check the input conditions directly, B Lemma 3.19 restricts the first parabolic product to words whose A letters are in the first j positions, whose singleton is neutral and whose C letters are after position $j+1$. On the rank orbit this says precisely $I_0 \subset U \subset U' \subset K_0$. The one-cut product similarly imposes just $I_0 \subset U \subset K_0$. Equivalently, divide the nested periodic lattices by the final lattice as in A's proof of Proposition 3C.4. For the output $F \subset F'$ of dimensions $m, m+1$, it also chooses a line $L \subset U \cap F$ and a two-plane D with $L \subset D \subset U' \cap F'$. Indeed the flags between the first affine parabolic and the output successively retain (L, U'), (L, D), (L, F'); this works also when $j = 1$ with the repeated cut suppressed. For $\mathcal{CBS}_s T$, the first affine parabolic retains just an s-space U with $I_0 \subset U \subset K_0$, and the final conditions become $L \subset F$, $L \subset D \subset U \cap F'$.

At $A_{j;m}$ the rank is j, so $U = I_0$ in the first gallery. A Lemma 4C.2 gives $\dim(I_0 \cap F) = \dim(I_0 \cap F') = 1$ and $\dim(K_0 \cap F') = m - j + 2$. Thus L is forced. The line U'/I_0 must belong to the image of $K_0 \cap F'$ modulo I_0; then $U' \cap F'$ is the unique possible D. The count is $(m-j+1)_q$ where q is the flag parameter. Its normalization shift is $2(m-j)$: the gallery normalization constant is

$$d_j^{\mathrm{BS}} = 4m - 3 + 2(j-1)(N-j-1),$$

as follows either by summing the parabolic length shifts or by the origin count below, and A Lemma 4C.2 gives $d_{A_{j;m}} = 2j(N-j) - 2(m-j) - 3$. Thus the normalized coordinate has constant one. For $\mathcal{CBS}_s T$, $s < j$ is impossible and $s = j$ would require $D \subset I_0 \cap F'$, also impossible. The constant therefore belongs to Y_j. By the margins of A Lemma 4I.1 it is the multiplicity of $C_{A_{j;m}}$. Monotonicity and the type bound now show that the family-F indices occurring in Y_j are exactly $A_{1;m}, \dots, A_{j;m}$, once each.

3. Coordinates at the full-spherical standards. At Z_r we have $I_0 \cap F' = 0$, $\dim(F \cap K_0) = m - r$, and $\dim(F' \cap K_0) = m - r + 1$, by A Definition 3C.1 and Lemma 4C.2. The choices of L, D in K_0 number $(m-r)_q^2$, and D is disjoint from I_0. For the BS gallery choose U'/I_0 containing the image of D, then a hyperplane U/I_0 containing the image of L. For the CBS gallery choose U/I_0 containing the image of D. The two flag counts are, respectively,

$$(m-r)_q^2 (j-r)_q \begin{bmatrix} N-2r-2 \\ j-r-1 \end{bmatrix}_q, \qquad (m-r)_q^2 \begin{bmatrix} N-2r-2 \\ s-r-2 \end{bmatrix}_q.$$

The first vanishes for $r \geq j$, the second for $r > s - 2$; these cases are omitted before evaluating Gaussians. The standard normalization is $v^{d - d_{Z_r}}$ times the count at $q = v^{-2}$, where $d_{Z_r} = 2r(N-r)$ as in A. To verify the absolute shifts without a diagram, use B's origin formulas in the proof of Proposition 2.6, multiplied by $[m]^2$ for T. Reversing the displayed counting polynomials at $r = 0$ gives $d = d_j^{\mathrm{BS}}$ as above and $d = d_s^{\mathrm{CBS}} = 6m - 2s + 2(s-2)(N-s)$. The flag normalization depends on the endpoint standard only through d_{Z_r} (A Lemmas 2D.4, 2E.3). Subtracting $2r(N-r)$ in either formula amounts to replacing (m, j, s, N) by $(m-r, j-r, s-r, N-2r)$.

Set $h = j - r > 0$, $t = v^2$. On its support the BS coordinate is therefore

$$v^{3-2h} (m-r)_t^2 (h)_t \begin{bmatrix} 2(m-r)-1 \\ h-1 \end{bmatrix}_t.$$

The corresponding CBS coordinate has valuation $2m + 4 - 2s$; by (1) its contribution to Y_j starts no lower than $5 + 2(j-s)$. Consequently, through degree one inclusive, the Z_r-coordinate of Y_j agrees with the Z_r-coordinate of $\mathcal{R}_j$ in *ambient parameter* j. Indeed $(m-r)_t \equiv (h)_t \bmod t^h$; both Gaussian numerators in the rectangle product formula start contributing only in degree h or higher. Multiplication by v^{3-2h} puts the error in degree at least three. For $r \geq j$ these coordinates vanish.

4. The same spherical corrections in the smaller parameter. For $r \leq s \leq j$, the Z_r-coordinate of the canonical C_{Z_s} is

$$\begin{bmatrix} 2(m-r)+1 \\ s-r \end{bmatrix}_t - \begin{bmatrix} 2(m-r)+1 \\ s-r-1 \end{bmatrix}_t$$

(A Lemma 4E.4). It agrees modulo t^{j-r+1} with the same coordinate in ambient parameter j: in both rectangle formulas the first numerator exponent is at least $j - r + 2$, unless the lower entry is negative, in which case the Gaussian is zero. In particular, for $s \leq j - 2$ this agreement persists through degree one after multiplication by A's correction g_{j-s}, whose valuation is $3 - 2(j-s)$.

Descend on r from j. At Z_r all one-incidence terms in Y_j, including their multiplicities, have valuation at least three (step 2 and A Lemma 4H.1). Every family-F coefficient there is proper, hence has positive valuation, and its multiplicity is constant. Full-spherical indices of rank less than r have zero coordinate. Thus after subtracting the already known higher full-spherical terms, the nonpositive powers at Z_r determine its bar-symmetric multiplicity completely. By step 3 and the preceding Gaussian comparison these powers agree with A's smaller-parameter computation (Lemmas 4J.2, 4K.1). It follows that the multiplicity at Z_r in Y_j is g_{j-r} for $0 \leq r \leq j-2$, and zero for $r \geq j-1$. At $r = 0$, after all these subtractions the coefficient of v is also the one in A's smaller calculation, namely $\binom{j+1}{2}$. For $j = 1$, the raw coordinate begins with v, and there is no spherical subtraction; this gives the same formula directly.

The only remaining contributors in degree one at Z_0 are the family-F terms of step 2. We have proved $\sum_{i=1}^{j} \mu_{i;m} = \binom{j+1}{2}$ while *holding m fixed*. Taking consecutive differences gives the individual formula. Neither v1 states this partial-sum calculation. It uses B's parabolic orbit estimates and A's multiplicity and Gaussian calculations, together with the finite flag identity for T.

Equivalently, before reading the distinguished coefficient, $X_i T = Y_i - Y_{i-1}$ contains $C_{A_{i;m}}$ once and none of the other family-F vectors. Its full-spherical coefficient at Z_r is $g_{i-r} - g_{i-1-r}$, with $g_a = 0$ for $a \leq 1$. The residual one-incidence terms can persist as operators. Their origin coordinates, including multiplicities, start in degree at least three by the same cleared-positivity bound. This identifies a separate canonical summand on B's side for each entry in A's row.

5 An independent edge calculation and rank reduction

Put $b_{j;m} = u_{j;m}^{-1}$ and $h_x = [M_e]\mathbf{C}_x$. In B's word convention the negative, zero and positive shifts in A's sorted window give the letters $\mathsf{C}, \mathsf{B}, \mathsf{A}$, respectively. Thus the word of $b_{j;m}$ is

$$\mathsf{C}^{j-1}\mathsf{B}^{m-j}\mathsf{ABCB}^{m-j}\mathsf{A}^{j-1}.$$

B's first and second chains keep m letters of each wound type throughout. The off-diagonal endpoints have $j < m$ of each.

The first entry admits a finite Hecke calculation

Deduction. For every $m \geq 2$,

$$P_{w_0,C_{1;m}}(q) = 1 + q + \cdots + q^{m-1}, \qquad \mu_{1;m} = 1.$$

Indeed, the window of s_0 has 0 at position 1 and $N+1$ at position N. Moving them to positions $m, m+2$ gives

$$b = b_{1;m} = s_0(s_1 \cdots s_{m-1})(s_{N-1} \cdots s_{m+2}) \in W_J.$$

Use the same $V = I \cap J$ and $L = V \setminus \{s_1, s_{N-1}\}$ as in A's gallery. Thus $W_V = S_m \times S_m$. Write $[a] = (v^a - v^{-a})/(v - v^{-1})$. In the spherical right module consider the bar-invariant vector $G = M_e C_{s_0} C_V/\pi_L$, where $C_V = C_{w_V}$. The right W_V-orbit of the left coset indexed by s_0 has minimal index s_0 and maximal index b. Its elements move the one A within the first block of m positions and the one C within the last block, keeping the middle B fixed. The stabilizer is W_L. Expanding the parabolic symmetrizer into minimal coset representatives gives

$$G = \sum_{x \in \mathcal{O}_V(s_0)} v^{\ell(b)-\ell(x)} M_x + v[m]^2 M_e.$$

Here $M_e C_{s_0} = M_{s_0} + vM_e$, and $\pi_V/\pi_L = [m]^2$. For the orbit term, each shortest shuffle moves the two marked letters away from their initial positions by length-increasing steps. Thus it acts by $M_{s_0} H_z = M_{s_0 z}$, with exponent $\ell(w_V) - \ell(w_L) - \ell(z) = \ell(b) - \ell(s_0 z)$. This is also the orbit formula underlying B §3.3. Now $v[m]^2 - [m][m-1] = v^m[m]$. The vector

$$\mathbf{C}_b = G - [m][m-1]M_e$$

is characterized by bar invariance, leading standard M_b with coefficient one, and strictly positive powers of v at all other standards. This proves the displayed KL formula since $h_b = v^m[m]$ and $\ell(b) = 2m - 1$.

This finite calculation gives the entire first-column KL polynomial. For example, at $m = 3$ it gives $1 + q + q^2$. Section 4 determines the leading coefficient of every column.

The first seed in B's uncorrected family

The same entry can be extracted from B's product in the ambient period $2m + 1$. With T_1, T_2 as in $(*)$ and A's strong one-line canonical $\mathcal{E}_{1,m}$ refined to V, one has

$$\mathcal{BS}_1 T_1 T_2 = \mathbf{C}_{b_{1;m}} + [m-1]\mathcal{E}_{1,m}.$$

Here $\mathcal{BS}_1 = D(J_1)$ has coefficient v^{2m-1} at e and v^{c-3} at the rank-one words with A in position one and C in position $3 \leq c \leq N$ (B Lemma 3.19). For the three output orbit maxima H, U, F, the A is in position m and the C in positions $1, m+1, m+2$, respectively. The four standard coordinates are

	H	U	F	e
$\mathcal{BS}_1 T_1 T_2$	$[m-1]$	$v^{m-1}[m-1]$	$v^{m-1}[m]$	$v^{2m-1}[m]^2$
$\mathcal{E}_{1,m}$	1	v^{m-1}	v^m	$v^{2m}[m]$.

For the first row, T_1 sends the terms $3 \leq c \leq m+1$ to their orbit sum with top coefficient $\sum_{c=3}^{m+1} v^{c-3+c-m-1} = [m-1]$, and multiplies the others by $[m]$. For H, the second averaging contributes the sum $\sum_{c=2}^{m} v^{c-2+c-m} = [m-1]$ divided by $[m-1]$; for U, F it moves a single A to the end of the first block. The second row follows from A Lemmas 4F.2 and 4D.1. These are all occupied orbits. Subtract the rows with the stated factor, using $v^{m-1}[m] - v^m[m-1] = 1$.

What rank reduction does compute

Let Θ_k have word $\mathsf{C}^k\mathsf{B}^{N-2k}\mathsf{A}^k$. It is the finite-orbit maximum of type k (B Lemma 3.11), hence represents A's full-spherical standard Z_k after output refinement. The coefficient at its longest fine representative has shift zero (A Lemma 4D.1). For $0 \le k < j \le m$ we obtain the all-degree identity

$$[\Delta_{Z_k}]C_{A_{j;m}} = [M_{\Theta_k}]\mathbf{C}_{b_{j;m}} = h^{(m-k)}_{b_{j-k;m-k}},$$

where the superscript gives the smaller rank parameter. For $k > 0$, both ordinary endpoints $w_I\Theta_k, w_I b_{j;m}$ take the values $2N, 1-N$ at positions $1, N$, respectively. Every fundamental-window entry lies between those bounds. Deleting the common first and last letters removes one C and one A: the relabelling in B Definition 4.17 gives $w'_I\Theta_{k-1}, w'_I b_{j-1;m-1}$, with period $N-2$. To check Bruhat comparability use the rank functions $\#\{i \le a : w(i) \ge b\}$. At each cut the deleted periodic strands contribute equally to the two functions. On the remaining strands their inequalities are exactly the smaller-period inequalities. Iterating reaches the comparable pair $w'_I, w'_I b_{j-k;m-k}$. B Proposition 4.18 now applies at each deletion and preserves the normalized coefficient. For $j \le k$ the original coordinate vanishes since $\ell(\Theta_k) = 2k(N-k) \ge 2j(N-j) > D_{j;m}$.

In particular B gives $[v][\Delta_{Z_k}]C_{A_{m;m}} = m-k$ for $0 \le k < m$. The final case reduces to s_0 in period three. The independent KL recursion checked the full polynomial identity for every k, j in ranks $m = 2, 3, 4$. For example, at $m = 4$ the rows of linear coefficients, with columns $j = 1, 2, 3, 4$, are $(1,2,3,4), (0,1,2,3), (0,0,1,2), (0,0,0,1)$. Together with A's one-line degree bound at each Z_k (Lemma 4H.1), these tests at full-spherical standards reproduce the row sums in parameters $m-k$. They do not provide another independent linear equation for the interior of a fixed row.

Two obstructions for $2 \le j < m$

At the ordinary pair $w_I, w_I b_{j;m}$ the only common window position is the central B at $m+1$. The other two B-blocks have $b(i) = i+1$ and $i-1$, respectively. Deleting the common position and image leaves type j in period $N-1$. In B Lemma 3.9, $\ell = 2j(N-j) - \mathrm{inv}$ with $\mathsf{C} < \mathsf{B} < \mathsf{A}$. Deletion drops two inversions, one with the preceding A, one with the following C. The ordinary length gap therefore drops by $2j-2 > 0$, preventing Henderson cancellation at this strand. Across fixed j the full gap difference is $D_{j;m} - D_{j;j} = (4j-2)(m-j)$.

There is also a concrete failure at the seed. B Proposition 3.21 gives lowest exponent $2-(N-2j) = 1-2(m-j) < 0$ for $\mathcal{BS}_j$ at the orbit maximum $z(j,1,1)$. The terms $\mathcal{BS}_{j-1}$ and $\mathcal{CBS}_s$ for $s \le j-1$ miss this type-j orbit. Replacing m by j in the correction while retaining period $2m+1$ therefore leaves a negative power. A proof of stability by this route would require a different correction or a different comparison.

6 What follows for coefficients and cohomology

Write $k = \overline{\mathbb{F}}_p$, $N = 2m + 1$, and $G = \mathrm{SL}_N(k)$. For degree one, both papers assume Lusztig's character formula in the given rank and $p \geq 5(N-1)$ for finite group comparison. These assumptions hold for every sufficiently large p at fixed m.

Setting and source	Consequence
A, Theorem 4L.1 and Corollary 6C.2	$\sum_{j=1}^{m} \mu_{j;m} = m(m+1)/2$. Each coefficient equals the rational H^1-dimension of $L(\lambda_{j;m}(p))$ and the finite H^1-dimension for its restriction to $\mathrm{SL}_N(\mathbb{F}_p)$ and its descendant $M_{j;m}(p)$ on $\mathrm{PSL}_N(\mathbb{F}_p)$, over k. A obtains $\max_j \mu_{j;m} \geq \lceil (m+1)/2 \rceil$.
B, Theorems A and B	$\mu_{m;m} = m$, and $\dim_{\mathbb{F}_p} H^1(\mathrm{PSL}_N(\mathbb{F}_p), V_{m,p}) = m$. All $V_{m,p}$ are faithful and absolutely irreducible.
The two stated KL theorems	$\sum_{j<m} \mu_{j;m} = m(m-1)/2$ and $\max_j \mu_{j;m} \geq m$. Those two numerical statements alone leave the maximizer undetermined.
Joint calculation, §4	$\mu_{j;m} = j$ throughout. A's m modules therefore realize every first cohomology dimension $1, \ldots, m$ in this single finite group for each sufficiently large p.
B, Theorem B, (2.17), Corollary 2.20	For $0 \leq s < m$, the normalized polynomial for y_s starts with mv^{s+1}. For large p, $\dim_k H^{s+1}(G, L(y_s \cdot_p 0)) = m$, and its rational cohomology vanishes in all lower degrees. The identified diagonal is $s = 0$.
B, Theorem C, Lemma 2.18	Fix $1 \leq n \leq m$. For sufficiently large p, then sufficiently large a, a faithful absolutely irreducible k-module for $\mathrm{PSL}_N(\mathbb{F}_{p^a})$ has $\dim_k H^n \geq m$. B proves the lower bound by injective restriction.
B, Theorem D, (2.18)	For a primitive complex root of odd order $\ell > N$, $\dim_{\mathbb{C}} H^{s+1}(U_\zeta, L_\zeta(y_s \cdot_\ell 0)) = m$, with vanishing in lower degrees. This is Ext in finite-dimensional integrable type-1 modules of Lusztig's divided-power quantum group.
A with B's (2.18)	The same standard quantum KL formula transports A's row sum to quantum H^1 for the m modules $L_\zeta(u_{j;m}^{-1} \cdot_\ell 0)$.

The coefficient fields. On the diagonal the highest weights agree. The split group gives an $\mathbb{F}_p$-form of the simple highest-weight module. Its root-lattice weight makes the finite center act trivially. That center has prime-to-p order, so inflation across it is an isomorphism in every degree. Thus, as modules for the *same* finite group,

$$M_{m;m}(p) \cong V_{m,p} \otimes_{\mathbb{F}_p} k.$$

Extending the finite group bar complex shows that the scalar extension preserves the cohomology dimension. The same descent applies to A's other restricted highest weights. See A Lemmas 6A.2, 6B.1 and 6C.1; B Proposition 2.15 and the proof of Theorem A.

An additional finite-group consequence. A stronger comparison applies to B's particular weights. Parshall–Scott–Stewart, *Shifted generic cohomology*, Theorem 5.12 and its proof on p. 1780 (Bendel–Nakano–Pillen, Theorem 7.5), take $H^d(G(\mathbb{F}_{p^a}), L(\lambda))$ to $H^d(G, L(\lambda))$ when λ is p^a-restricted, lies in the affine dot orbit of zero, and $p \geq (4d+1)(N-1)$. Take $\lambda = y_{n-1} \cdot_p 0$; B Lemma 2.17 gives restrictedness for every $a \geq 1$. For p also large enough for B's rational formula, applying this comparison for $1 \leq d \leq n$ and inflating across the center gives $\dim_k H^n = m$ and vanishing below n on $\mathrm{PSL}_N(\mathbb{F}_{p^a})$, including $a = 1$. Degree zero vanishes by nontrivial irreducibility. This strengthens the stated Theorem C. In all settings n can be fixed and the rank can grow; neither paper proves unboundedness at fixed p. doi:10.1112/S0010437X13007331.

7 Exact calculations in the spherical module

The accompanying Python program implements two calculations over $\mathbb{Z}[v, v^{-1}]$. In the first, represent a standard vector by the levels $a_i = \lfloor (x(i) - 1)/N \rfloor$ of the left-minimal window. To apply C_P/π_T, where T is the previous intersection, cut the circle at a missing reflection and add one to the levels that pass the cut. In each position block sort the lifted levels in increasing order. Let L be the stabilizer of this minimum in W_P, $d(x)$ the within-block inversion number and $D = \ell(w_P) - \ell(w_L)$. The new coefficient at y in its orbit O is

$$v^{D-d(y)} \frac{\pi_L}{\pi_T} \sum_{x \in O} v^{-d(x)} [M_x] F.$$

Indeed $M_x C_P = v^{-d(x)} M_{\min O} C_P$ and splitting W_P into minimal left W_L-coset representatives evaluates the latter expression. All Laurent divisions in the program check for zero remainder.

The second calculation uses only right simple reflections. For a right descent s of x, recursively calculate $\mathbf{C}_{xs} C_s$. At each strict lower z subtract the constant term of its standard coordinate times $\mathbf{C}_z$. The resulting monic bar-invariant vector satisfies the canonical triangularity condition. This independently computes B's seed, every endpoint, and the canonical transition matrix used to decompose the products. The numerical inputs are the parabolic sets, the chain words and the endpoint windows.

Let $L_m = C_{s_2} \cdots C_{s_m} C_{s_1} \cdots C_{s_{m-1}}$ and let D_m denote the two normalized parabolic averages. Each cell below gives the degree-one identity coefficient followed by the number of distinct canonical indices with nonzero graded multiplicity.

m	$\mathcal{R}_m$	$\mathcal{BS}_m L_m$	$\mathbf{C}_{X_0} L_m$	$\mathbf{C}_{X_0} D_m$	$\mathbf{C}_{\beta_m}$
2	4, 5	4, 5	3, 3	3, 3	2, 1
3	11, 11	18, 23	12, 13	7, 6	3, 1
4	25, 17	81, 70	50, 38	14, 9	4, 1

In each rank all five columns have identical standard support, respectively 38, 368, 3098 indices, and coefficient one at M_{β_m}. Every coordinate of $\mathcal{BS}_m D_m$ agrees with $\mathcal{R}_m$. Of the family-F summands, the first two columns contain the entire row with multiplicity one, and the last three contain just its diagonal. The independent KL recursion gives rows $(1, 2), (1, 2, 3), (1, 2, 3, 4)$. For instance

$$[M_e]\mathbf{C}_{\beta_2} = 2v + 4v^3 + 5v^5 + 4v^7 + v^9, \qquad [M_e]\mathbf{C}_{X_0} = v + v^3 + 2v^5 + v^7 \quad (m = 2).$$

Executing `python3 compare.py -full 2 3 4` in the directory containing `compare.py` reproduces the tables and full identity polynomials in `results.txt` without external packages. A second script, `python3 partial.py 2 3 4 5`, checks the mixed full-spherical coordinates and the discarded degrees directly. In rank five its successive corrected degree-one coefficients are $1, 3, 6, 10, 15$; the independent KL recursion gives $(1, 2, 3, 4, 5)$. Outside the rows printed in A, a separate run gives $\mu_{2;6} = \mu_{2;7} = \mu_{2;8} = 2$, $\mu_{3;6} = \mu_{3;7} = 3$ and $\mu_{4;6} = 4$. The last expansion has 120 862 standard terms. The script `check_off_diagonal.py` and its output accompany the calculation.

A boundary projection still detects the difference. At $m = 2$, $L_m = D_m$. For $m \geq 3$, even after the projector C_V/π_V the vector $\mathcal{BS}_m L_m$ is not proportional to $\mathcal{R}_m$ over $\mathbb{Q}(v)$. It suffices to compare two coordinates at $v = 1$. Write $t = 2^{m-1}/m$. The ratio at M_e is t^2 since each simple factor multiplies this coordinate by 2, whereas each divided parabolic factor multiplies it by m. The ratio at M_{β_m} is t, as follows.

In type m the first affine product has just the word $A^m BC^m$. The next K_m-average fixes the initial A and puts the B in any one of positions $2, \ldots, m+1$. The E-average then gives coefficient one to every type-m word with prefix AB (the stabilizer of the remaining letters has size $(m-1)!m!$, exactly the overlap denominator). The last m letters stay fixed during L_m. In the V-orbit of β_m they contain one C and $m-1$ letters A, so only the initial prefix ABC^{m-1} of length $m+1$ can contribute: the leftmost letter is A and the second letter cannot be C. All first-chain swaps must be chosen to put B at position $m+1$. Each of the 2^{m-1} choices in the second chain then contributes. There are m possible suffixes and m^2 words in the target orbit. The normalized V-average gives $m2^{m-1}/m^2 = t$ at M_{β_m}, where $\mathcal{R}_m$ has coefficient one. The denominators and the coordinate used to compare scalars are nonzero at 1. Since $t > 1$, the two ratios differ.

Partial galleries. For $T_j = \mathcal{BS}_j D_m$ with $1 \leq j \leq m \leq 4$, exact elimination gives family-F multiplicity one exactly at indices $1, \ldots, j$. A truncated gallery can also contain canonical indices outside the six families for $\mathcal{R}_m$. At $(m, j) = (3, 2)$ its top word is $CCABBBA$, and the desired $CBABCBA$ is a lower summand. This is why identifying the positive summand supported on the relevant seed indices matters in the all-rank argument.

8 What the public record establishes

Both supplied source archives and both PDFs match the version-specific arXiv downloads byte for byte. All times below are UTC.

Record	Finding
Discovery accounts	A, Sections 7A–C, describes starting from planar webs and modifying a graph on an annulus. B's AI statement (pp. 3–4) dates the project's start to the 2025 ICERM semester. The authors describe identifying β_r in data supplied by Astra, then requesting a nearby X_0 and a path of finite reflections. They give no dates for these two steps.
2 November 2026	The manuscript commit records 21:39:44. A public push of this exact commit to `main` is recorded at 21:39:52 (event 26157801356). The pinned TeX, lines 3186–3246, already contains the endpoints, the row-sum theorem, four computed rows and the conjectural value $\mu_{j;m} = j$.
3 November 2026	The AI guide and Markdown manuscript commit records 02:54:46. The matching public push is recorded at 02:54:56 (event 26176959605). The pinned Markdown gives the same family and suggested formula. Both events were returned by the GitHub events API. The supplement preserves the relevant fields.
arXiv	A v1 was submitted on 6 November at 07:48:34 and B v1 on 8 November at 23:04:40. The daily listings placed A on 9 November and B on 10 November. B submitted before A's scheduled announcement at 01:00 on 9 November.
B's v1 bibliography	In the submitted source, `bibliography.bib`, lines 504–511, records A's four authors, title, year, project URL and the note *Work in progress*. Of 54 entries, 41 distinct keys are cited and 13 are unused, including `GLTV`. The TeX contains no `\nocite`. A is absent from the 41 printed references.
Inference	The public push records establish that the repository contained a checkable construction before either submission. The bibliography entry establishes that information about it reached B's source by submission.
Unresolved	Which person or tool supplied the entry, which materials anyone actually read, and when that happened relative to discovery of the family and proof.

Interpretation. The common diagonal beyond $N = 5$, the identical first affine parabolic and the equality of the completed raw products give precise features to trace. The flag identity and the partial-row argument are deductions of this comparison. The different seed, reflection chains and higher-degree theorem identify what B adds at the diagonal. None of these features determines what anyone read. The public manuscript supplied the expected diagonal value as a conjecture. A later literature search after discovery, earlier exposure to the family followed by development of a proof, and earlier exposure to the proof followed by reworking or extension all fit those facts. An entry imported by an agent or contributor without every coauthor seeing it could accompany any of these histories.

Questions for the authors. Did the reflection strings arise by refining a parabolic average, and did you consider the products $\mathcal{BS}_j$ in a fixed larger rank? When do β_r and, separately, X_0, the two chains and their corrections first appear in a dated calculation, draft or conversation? Which revision introduced `GLTV`, and does a surviving search result or retrieval record identify the page or file opened and its date? A first formula, a first proof and a first recorded visit answer different questions.

9 Text for the revision

The introduction below assumes the revision includes the partial-row proof in §4. For a shorter revision covering only the raw operator identity, omit the final sentence of the introduction passage and the final two sentences of the technical remark.

Introduction

In concurrent work, Bowman, de la Fuente, Libedinsky and Plaza [B+26, Theorems A–D] determine the last coefficient in our row, $\mu_{m;m} = m$, and obtain unbounded cohomology in every fixed positive degree. Their degree-one example and ours at $j = m$ have the same highest weight. Our web computes the sum of the whole row. Comparing it with their parabolic construction gives the partial sums $\sum_{i=1}^{j} \mu_{i;m} = \binom{j+1}{2}$ and hence $\mu_{j;m} = j$ for every $1 \le j \le m$.

Technical remark

Remark 1 *Use the inverse-permutation convention of Lemma 3B.1, and write $\mathcal{BS}_m$ for the uncorrected product of [B+26, Definition 3.2]. Put $Q_1 = \langle s_2, \dots, s_m \rangle$ and $Q_2 = \langle s_1, \dots, s_{m-1} \rangle$. Then*

$$\mathcal{R}_m = \frac{\mathcal{BS}_m C_{Q_1} C_{Q_2}}{([m-1]!)^2}.$$

The two parabolic factors have relative longest words $s_2 \cdots s_m$ and $s_1 \cdots s_{m-1}$, the two strings in [B+26, Lemma 2.4]. To see the identity, split the common annular merge back to $(m, 1, m)$ and fix the input and output flags $U \subset U'$ and $W \subset W'$ of dimensions $m, m+1$. Both galleries choose a line $a \subset U \cap W$. Our gallery then chooses a hyperplane containing $U' + W'$, while the other chooses a plane between a and $U' \cap W'$. The two choices have the same count since $\operatorname{codim}(U' + W') = \dim((U' \cap W')/a)$. Both finite tails have normalization shift $m^2 + 2m - 3$. In [B+26, Proposition 3.5] the product $\mathcal{BS}_m$ is first corrected to a single canonical element. Following the two strings with the corrections of [B+26, Propositions 2.8 and 2.11] then gives the canonical term indexed here by $A_{m;m}$. More generally, completing the canonical element indexed by $z(i, 1, 0)$ in [B+26, Section 3.2] with these two parabolic factors gives the term indexed by $A_{i;m}$ with multiplicity one. All other terms from our distinguished family have multiplicity zero, as the partial-sum calculation shows.

Methods section, evidence-only version

Our manuscript and accompanying repository were publicly available before the two arXiv submissions. The source archive of [B+26, version 1] includes an uncited bibliography entry for our project webpage. The authors describe their search for the family and the subsequent proof in their AI statement.

Methods section, conditional version

Our manuscript was publicly available while the two projects were in progress. The source archive of [B+26, version 1] includes an uncited entry for our project webpage. If the page was encountered during that project, the public record leaves open whether this happened before or after the family was found, or during a later literature search. The evidence-only version is my choice for publication. The conditional version is supportable if the authors want to record the timing question explicitly. A statement that an agent found the construction, that it transformed a proof, or that either discovery was independent would need evidence beyond the material audited here.

References

[B+26] C. Bowman, D. de la Fuente, N. Libedinsky and D. Plaza. The cohomology of finite simple groups is unbounded. 2026. https://arxiv.org/abs/2611.11677.