

# Economic Time Series Modeling and Forecasting

David Ubilava

July 2021

# Contents

<b>Preamble</b>	<b>3</b>
<b>Preliminaries</b>	<b>4</b>
<b>1 Introduction to Forecasting</b>	<b>5</b>
<b>2 Stochastic Process and Time Series</b>	<b>7</b>
<b>Deterministic Time Series Models</b>	<b>12</b>
<b>3 Trends and Seasonality</b>	<b>13</b>
3.1 Spurious Relationship . . . . .	13
3.2 Trends . . . . .	16
3.3 Seasonality . . . . .	18
3.4 Exercise . . . . .	19
<b>Dynamic Time Series Models</b>	<b>22</b>
<b>4 Linear Autoregression</b>	<b>23</b>
4.1 Modeling . . . . .	24
4.2 Forecasting . . . . .	27
<b>5 Vector Autoregression</b>	<b>31</b>
<b>6 Threshold Autoregression</b>	<b>32</b>
<b>Forecast Assessment</b>	<b>33</b>
<b>7 Forecast Evaluation</b>	<b>34</b>
<b>8 Forecast Combination</b>	<b>35</b>
<b>Appendix: Introduction to R</b>	<b>36</b>
8.1 Data Management . . . . .	36
8.2 Data Visualisation . . . . .	38

<i>CONTENTS</i>	2
8.3 Regression Analysis . . . . .	42

# Preamble

These notes are prepared to teach an undergraduate-level course on economic forecasting. The content is presented in four parts. The first part, *Preliminaries*, introduces the concept of forecasting in time series context. The second part, *Deterministic Time Series Models*, covers trends and seasonal models. The third part, *Dynamic Time Series Models*, covers linear, multivariate, and nonlinear autoregressive processes. The fourth part, *Forecast Assessment*, goes over forecast evaluation and combination routines, and forecast accuracy tests.

# Preliminaries

# Chapter 1

## Introduction to Forecasting

Economic events tend to co-occur, precede, or succeed one another. Understanding the essence of such relationships – that is, identifying causal mechanisms that facilitate correlation among economic variables – is at the core of econometric analysis. Throughout a relatively brief history of the study of econometrics, numerous methods and techniques have been proposed and developed – all aimed to give an empirical content to economic models. These methods and techniques allow us to test theories, evaluate policy outcomes, etc.

Econometric models rely on correct (and accurate) identification of the causal mechanism in the underlying process. But they are also predictive by nature – they help us make economic forecasts even when the causal mechanism may not be well identified. In other words, while correlation does not necessarily imply causality, if the goal is to make a forecast, a mere correlation might as well suffice.

Roots of forecasting extend very much to the beginning of human history. In their desire to predict the future, people have attempted to make forecasts of their own, or have used services of others. Fortunetellers, for example, have been forecast experts of some sort, basing their predictions on magic. They are less common in the current age. Astrologers, who rely on astronomical phenomena to foresee the future, maintain their relevance to this date. Over time, and particularly with the development of the study of econometrics, more rigorous forecasting methods have been introduced and developed. All methods – primitive or complex, spurious or scientifically substantiated – have one thing in common: they all rely (or, at least, pretend to rely) on *information*.

Information is key in forecasting. It comes in many forms, but after it is organized and stored, what we end up with is data. A diverse set of forecasting methods typically rely on insights from econometric analysis of time series – chronologically stored data, collected at regular intervals over a period of time. In time series analysis, the implied assumption is that the past tends to repeat

itself, at least to some extent. So, if we well study the past, we *may* be able to forecast an event with some degree of accuracy.

Accurate forecasting is difficult. No matter how rich the available data are, or how well the econometric model fits the data, there still is a surprise element concerning the forecast – something that has never happened in past, and is only specific to the future. And because of this, there is no such thing as precise forecast, even if by fluke we were to exactly predict an outcome of an event. But some forecasts are better than others. And in search of such forecasts the study of time series econometrics has evolved.

## Chapter 2

# Stochastic Process and Time Series

Time series are realizations of a chronologically stored sequence of random variables. This sequence of random variables is referred to as the *stochastic process*. Thus, a time series is a realisation of the stochastic process. We index time periods as  $1, 2, \dots, T$ , and denote the set of observations as  $\{y_1, \dots, y_T\}$ . One can view a time series as a finite sample from an underlying doubly-infinite sequence:  $\{\dots, y_{-1}, y_0, y_1, y_2, \dots, y_T, y_{T+1}, y_{T+2}, \dots\}$ .

If all random variables, from where the time series are drawn, have the same distribution, then we refer to such time series as *stationary*. Stationarity is an important feature, and the assumption, on which the time series analysis heavily relies.

Before diving any further into the concepts and methods of time series econometrics, consider the simplest kind of time series comprised of realizations from independent and identically distributed random variable with zero mean and constant variance:  $\varepsilon_t \sim iid(0, \sigma^2)$ . The following graph plots this time series against time.

Such time series are referred to as *white noise*. That is, a time series,  $y$ , is a white noise process if:

$$\begin{aligned} E(y_t) &= 0, \quad \forall t \\ Var(y_t) &= \sigma^2, \quad \forall t \\ Cov(y_t, y_{t-k}) &= 0, \quad \forall k \neq 0 \end{aligned}$$

Because each observation is drawn from the same distribution, white noise is a stationary time series. Indeed, it is a special type of stationary time series insofar as its mean, variance, and covariance are time-invariant. Note, for a time






Figure 2.1: White noise: an illustration

series to be stationary, the mean doesn't need to be zero (as long as it is constant over time), nor covariances need to be equal to zero (as long as they are constant over time, though they may vary with  $k$ ). Thus,  $\{y_t\}$  is stationary if the mean and variance are independent of  $t$ , and the autocovariances are independent of  $t$  for all  $k$ .

In fact, it is more of a norm rather than an exception for a time series to be correlated over time. Indeed, because of the sequential nature of time series, we commonly observe dependence among the temporally adjacent time series. That is, for most economic time series, we would expect  $y_t$  and  $y_{t-1}$  to be correlated. Such correlations are referred to as *autocorrelations*, and are given by:

$$Cor(y_t, y_{t-k}) = \frac{Cov(y_t, y_{t-k})}{Var(y_t)}, \quad k = 1, 2, \dots$$

Autocorrelations are commonly illustrated via the so-called *autocorrelogram*, which plots the sequence of autocorrelation coefficients against the lags at which these coefficients are obtained. For example, an autocorrelogram of the previously illustrated white noise process is as follows:

For each  $k$ , the vertical line extending from zero represents the autocorrelation coefficient at that lag. The red dashed lines denote the 90% confidence interval, given by  $\pm 1.96/\sqrt{T}$ , where  $T$  is the length of the time series.







Figure 2.2: Autocorrelogram

It is common to transform time series by taking logarithms, differences, or differences of logarithms (growth rates). Such transformations usually are done to work with the suitable variable for the desired econometric analysis. For example, if an economic time series is characterized by an apparent exponential growth (e.g., real GDP), by taking natural logarithms the time series “flatten” and the fluctuations become proportionate. The difference operator is denoted by  $\Delta$ , so that  $\Delta y_t = y_t - y_{t-1}$ . The following three graphs illustrate (i) a time series with an apparent exponential growth, (ii) the natural logarithm of this time series, and (iii) their differences (i.e., the log-differences of the original series).





# Deterministic Time Series Models

## Chapter 3

# Trends and Seasonality

Economic time series usually are characterized by trending behavior, and often present a seasonal pattern as well. Trend is a unidirectional change of time series over an extended period of time that arises from the accumulation of information over time. Seasonality is a repeating pattern *within a calendar year* that arises from the links of technologies, preferences, and institutions to the calendar. Modeling and forecasting these time series features is a fairly straightforward task. But before we get to it, let's discuss what may happen if we were to ignore the presence of trends and/or seasonality when analyzing the time series data.

### 3.1 Spurious Relationship

Nothing about trending time series necessarily violates the classical linear regression model assumptions. The issue may arise, however, if an unobserved trending variable is simultaneously correlated with the dependent variable as well as one of the independent variables in a time series regression. In such case, we may find a (statistically significant) relationship between two or more unrelated economic variables simply because they are all trending. Such relationship is referred to a *spurious relationship*.


To illustrate, consider two trending variables:

$$y_t = \gamma t + \nu_t, \quad \nu \sim N(0, \sigma_\nu^2),$$

and

$$x_t = \delta t + v_t, \quad v \sim N(0, \sigma_v^2),$$


where  $Cov(\nu_t, v_t) = 0$ . For simplicity, we can assume  $\sigma_\nu^2 = \sigma_v^2 = 1$ . Suppose,  $\gamma$  and  $\delta$  are some positive scalars, say, 0.3 and 0.5, respectively. That is,  $y$  and  $x$  are trending in the same direction. Below is an example of such time series:



If we were to estimate

$$y_t = \alpha + \beta x_t + \varepsilon_t,$$

we are likely to find the relationship between the two – in this case  $\beta > 0$  – even though, we know, the two are not related. To illustrate this, we will generate 1000 samples of size 120 for  $y$  and  $x$ , and in each case we will estimate the parameter  $\beta$ . The following graph illustrates the empirical distribution of these parameter estimates:




Luckily, we can easily “fix” the issue, by incorporating a trend in the regression:

$$y_t = \alpha + \beta x_t + \eta t + \varepsilon_t.$$

Once the trend is accounted for, the previously illustrated “bias” disappears. Using a similar simulation exercise as before, the following graph illustrates the empirical distribution of these parameter estimates:





In fact, this “fix” is equivalent to regressing a de-trended  $y$  on a de-trended  $x$ . To de-trend a variable, we first run a regression:  $y_t = \gamma_0 + \gamma_1 t + \nu_t$ , and then obtain the fitted values for some fixed trend (typically zero), that is:  $\tilde{y}_t = \hat{\gamma}_0 + \hat{\nu}_t$ , where  $\hat{\gamma}_0$  and  $\hat{\nu}_t$  are the parameter estimate and the residuals from the foregoing regression.

## 3.2 Trends

### 3.2.1 Modeling

As seen, accounting for trends in a time series can help us resolve some regression issues. But a trend in and of itself can be an inherent feature of a times series. To that end, we can apply deterministic trends to forecast time series.

The simplest (and perhaps most frequently applied) model to account for the trending time series is a *linear* trend model:

$$y_t = \alpha + \beta t$$

Other likely candidate trend specifications are *polynomial* (e.g. quadratic, cubic, etc.), *exponential*, and *shifting* (or *switching*) trend models, respectively given by:

$$\begin{aligned} y_t &= \alpha + \beta_1 t + \beta_2 t^2 + \dots + \beta_p t^p \\ y_t &= e^{\alpha + \beta t} \quad \text{or} \quad \ln y_t = \alpha + \beta t \\ y_t &= \alpha + \beta_1 t + \beta_2 (t - \tau) I(t > \tau), \quad \tau \in \mathbb{T} \end{aligned}$$

Of these, here we will primarily consider linear and quadratic trends. An exponential trend, from the standpoint of modeling and forecasting, is equivalent to a linear trend fitted to natural logarithm of the series. For a time series  $\{y_t : t = 1, \dots, T\}$ , the natural logarithm is:  $z_t = \ln y_t$ . Some of the benefits of such a transformation are that:

- they are easier to interpret (relative/percentage change).
- they homogenizes the variance of the time series.
- they may result in improved forecasting accuracy.

Exponential trends are suitable when a time series is characterized by a stable relative change over time (e.g., when economic time series grow by 2% every year).

We will cover the shifting/switching trend models in another chapter.

### 3.2.2 Forecasting

Trends are (relatively) easy to model and forecast. Caution is needed, however, with (higher order) polynomial trends, as they may fit well in-sample, but cause major problems out-of-sample.

Consider a linear trend model with an additive error term:

$$y_t = \alpha + \beta t + \varepsilon_t$$

We estimate the model parameters,  $\theta = \{\alpha, \beta\}$ , by fitting the trend model to a time series using the least-squares regression:

$$\hat{\theta} = \underset{\theta}{\operatorname{argmin}} \sum_{t=1}^T (y_t - \alpha - \beta t)^2.$$

Fitted values are then given by:

$$\hat{y}_t = \hat{\alpha} + \hat{\beta}t$$

Any future realization of a random variable is assumed to follow a linear trend model:

$$y_{t+h} = \alpha + \beta(t+h) + \varepsilon_{t+h}.$$

An optimal forecast of  $y_{t+h}$ , therefore, is given by:

$$y_{t+h|t} = E(y_{t+h}|\Omega_t) = E[\alpha + \beta(t+h) + \varepsilon_{t+h}] = \alpha + \beta(t+h).$$

The forecast error is:

$$e_{t+h|t} = y_{t+h} - y_{t+h|t} = \varepsilon_{t+h}$$

The forecast variance, then, is:

$$\sigma_{t+h|t}^2 = E(e_{t+h|t}^2) = E(\varepsilon_{t+h}^2) = \hat{\sigma}^2, \quad \forall h$$

From this, we can obtain interval forecast at any horizon, which is:

$$y_{t+h|t} \pm 1.96\hat{\sigma}.$$

A few features of trend forecasts to note:

- they tend to understate uncertainty (at long horizons as the forecast interval doesn't widen with the horizon);
- short-term trend forecasts can perform poorly; long-term trend forecasts typically perform poorly;
- sometimes it may be beneficial, from the standpoint of achieving better accuracy, to forecast growth rates, and then reconstruct level forecasts.

### 3.3 Seasonality

#### 3.3.1 Modeling

Seasonality is typically modeled as monthly or quarterly pattern, but can also be modeled as a higher frequency pattern (e.g. weekly). Some examples of time series with apparent seasonal patterns are:

- Agricultural production.
- Sales of energy products.
- Airfare (in non-pandemic times).

One way to deal with the seasonality in data is to “remove” it prior to the use of the series (i.e., work with a seasonally adjusted time series). Indeed, some economic time series are only/also available in a seasonally-adjusted form.

Otherwise, and perhaps more interestingly, we can directly model seasonality in a regression setting by incorporating seasonal dummy variables.

A seasonal model is given by:

$$y_t = \sum_{i=1}^s \gamma_i d_{it} + \varepsilon_t,$$

where  $s$  denotes the frequency of the data, and  $d_{it}$  takes the value of 1 repeatedly after every  $s$  periods, and such that  $\sum_i d_{it} = 1, \forall t$ .

Alternatively the seasonal model can be rewritten as:

$$y_t = \alpha + \sum_{i=1}^{s-1} \delta_i d_{it} + \varepsilon_t,$$

in which case  $\alpha$  is an intercept of an omitted season, and  $\delta_i$  represents a deviation from it during the  $i^{th}$  season.

Both variants of the seasonal model result in an identical fit and forecasts.

### 3.3.2 Forecasting

Any future realization of a random variable that is assumed to follow a seasonal model is:

$$y_{t+h} = \alpha + \sum_{i=1}^{s-1} \delta_i d_{i,t+h} + \varepsilon_{t+h}.$$

The optimal forecast of  $y_{t+h}$  is:

$$y_{t+h|t} = E(y_{t+h}|\Omega_t) = \alpha + \sum_{i=1}^{s-1} \delta_i d_{i,t+h}$$

The forecast error is:

$$e_{t+h|t} = y_{t+h} - y_{t+h|t} = \varepsilon_{t+h}$$

The forecast variance is given by:

$$\sigma_{t+h|t}^2 = E(e_{t+h|t}^2) = E(\varepsilon_{t+h}^2) = \hat{\sigma}^2, \quad \forall h$$

The interval forecast at any horizon is:

$$y_{t+h|t} \pm 1.96\hat{\sigma}.$$

## 3.4 Exercise

```
# set the length of a time series to 180 (e.g., 15 years of monthly data)
n <- 180

# set seed to 2 and generate a sequence of standard normal error terms
set.seed(2)
e <- rnorm(n)

# suppose the data generating process of a time series is:
# y_{t} = 0.7*y_{t-1}+0.01*t+e_{t}
# simulate these series, assume y_{0} = 0.
y <- e

y[1] <- 0.01+e[1]
```


```

for(i in 2:n){
  y[i] <- 0.7*y[i-1]+0.05*i-0.0002*i^2+e[i]
}

# store the simulated series in a data.table, where the first column is a column of dates
dt <- data.table(year=seq(as.Date("1980-01-01"),by="quarter",along.with=y),y=y,trend=c(1:n))

# plot the series (using ggplot)
ggplot(dt,aes(x=year,y=y))+
  geom_line(size=1)+
  theme_classic()+
  theme(axis.title=element_text(size=14),axis.text=element_text(size=12))

```



```

# suppose we think the linear trend model is appropriate; use the first 120 observations
# to estimate trend, and using this estimate generate forecasts for the remaining series
R <- round(.75*n)

reg <- lm(y[1:R]~trend[1:R],data=dt)

# store the vector of coefficients
theta <- reg$coefficients

# calculate the residual standard deviation
sigma <- sqrt(crossprod(reg$residuals)/(R-2))

```

```


# generated and include in the dataset the fitted values and 90% CI bounds
dt[, `:=` (yhat=(theta[1]+theta[2]*trend))]
dt[, `:=` (y.lo=(yhat-1.96*as.vector(sigma)), y.hi=(yhat+1.96*as.vector(sigma)))]

# set the fitted values of the in-sample range to NA
dt[trend <= R, c("yhat", "y.lo", "y.hi")] <- NA

# re-arrange the dataset into the "long" format (helpful for plotting)
dt_long <- melt(dt[, .(year, y, yhat, y.lo, y.hi)], id.vars="year")

# plot the out-of-sample forecasts along with the observed time series
ggplot(dt_long, aes(x=year, y=value, color=variable, linetype=variable)) +
  geom_line(na.rm=T, size=1) +
  scale_color_manual(breaks = c("y", "yhat", "y.lo", "y.hi"), labels = c("time series", "point forecast", "lower bound", "upper bound")) +
  scale_linetype_manual(breaks = c("y", "yhat", "y.lo", "y.hi"), labels = c("time series", "point forecast", "lower bound", "upper bound")) +
  theme_classic() +
  theme(axis.title=element_text(size=14), axis.text=element_text(size=12), legend.position = "top")

```



# Dynamic Time Series Models

## Chapter 4

# Linear Autoregression

Economic time series are often characterized by stochastic cycles. A cycle is a pattern of periodic fluctuations, not contained within a calendar year. A stochastic cycle is one generated by random variables. In general terms, the process is given by:

$$Y_t = f(Y_{t-1}, Y_{t-2}, \dots; \theta) + \varepsilon_t. \quad t = 1, \dots, T$$

An autoregressive process (or, simply, an autoregression) is a regression in which the dependent variable and the regressors belong to the same stochastic process.

Autoregression of order  $p$ , referred to as  $AR(p)$ , has the following functional form:

$$y_t = \alpha + \beta_1 y_{t-1} + \beta_2 y_{t-2} + \dots + \beta_p y_{t-p} + \varepsilon_t$$

The sum of the autoregressive parameters,  $\beta_1, \dots, \beta_p$ , depicts the persistence of the series. The larger is the persistence (i.e., closer it is to one), the longer it takes for the effect of a shock to dissolve.

The autocorrelation,  $\rho$ , and partial autocorrelation,  $\pi$ , functions of the covariance-stationary  $AR(p)$  process have the following distinctive features:

- $\rho_1 = \pi_1$ , and  $\pi_p = \beta_p$ .
- The values of  $\beta_1, \dots, \beta_p$  determine the shape of the autocorrelation function (ACF); in any case, the smaller (in absolute terms) is the persistence measure, the faster the ACF decays toward zero.
- The partial autocorrelation function (PACF) is characterized by “statistically significant” first  $p$  spikes  $\pi_1 \neq 0, \dots, \pi_p \neq 0$ , and the remaining  $\pi_k = 0, \forall k > p$ .



## 4.1 Modeling

### 4.1.1 AR(1)

Consider the first-order autoregression:

$$y_t = \alpha + \beta_1 y_{t-1} + \varepsilon_t,$$

where  $\alpha$  is a constant term;  $\beta_1$  is the *persistence* parameter; and  $\varepsilon_t$  is a white noise process.

Note that a necessary and sufficient condition for an  $AR(1)$  process to be covariance stationary is that  $|\beta_1| < 1$ . We can see this by substituting recursively the lagged equations into the lagged dependent variables:

$$\begin{aligned} y_t &= \alpha + \beta_1 y_{t-1} + \varepsilon_t \\ y_t &= \alpha + \beta_1(\alpha + \beta_1 y_{t-2} + \varepsilon_{t-1}) + \varepsilon_t \\ &= \alpha(1 + \beta_1) + \beta_1^2(\alpha + \beta_1 y_{t-3} + \varepsilon_{t-2}) + \beta_1 \varepsilon_{t-1} + \varepsilon_t \\ &\vdots \\ &= \alpha \sum_{i=0}^{k-1} \beta_1^i + \beta_1^k y_{t-k} + \sum_{i=0}^{k-1} \beta_1^i \varepsilon_{t-i} \end{aligned}$$

The end-result is a general linear process with geometrically declining coefficients. The  $|\beta_1| < 1$  is required for convergence.

Assuming  $|\beta_1| < 1$ , as  $k \rightarrow \infty$  the process converges to:

$$y_t = \frac{\alpha}{1 - \beta_1} + \sum_{i=0}^{\infty} \beta_1^i \varepsilon_{t-i}$$

The *unconditional mean* of this process is:

$$\mu = E(y_t) = E\left(\frac{\alpha}{1 - \beta_1} + \sum_{i=0}^{\infty} \beta_1^i \varepsilon_{t-i}\right) = \frac{\alpha}{1 - \beta_1}$$

And its *unconditional variance* is:

$$\gamma_0 = Var(y_t) = Var\left(\frac{\alpha}{1 - \beta_1} + \sum_{i=0}^{\infty} \beta_1^i \varepsilon_{t-i}\right) = \frac{\sigma_\varepsilon^2}{1 - \beta_1^2}$$

*Autocovariance* is simply the covariance between  $y_t$  and  $y_{t-k}$ , that is:

$$\gamma_k = Cov(y_t, y_{t-k}) = E[(y_t - \mu)(y_{t-k} - \mu)] = E(y_t y_{t-k}) - \mu^2$$

Some algebraic manipulation can help us show that:

$$\gamma_k = \beta_1 \gamma_{k-1},$$

and that:

$$\rho_k = \beta_1 \rho_{k-1}$$

(recall, *autocorrelation*  $\rho_k = \gamma_k / \gamma_0$ ).

In fact, autocorrelation of any order can be given as the autoregression parameter (which in this instance is equivalent to the persistence measure) to that power:

$$\begin{aligned}\rho_1 &= \beta_1 \rho_0 = \beta_1 \\ \rho_2 &= \beta_1 \rho_1 = \beta_1^2 \\ &\vdots \\ \rho_k &= \beta_1 \rho_{k-1} = \beta_1^k\end{aligned}$$

The autocorrelation of AR(1), given that  $|\beta_1| < 1$ , is a geometric decay; the smaller is  $|\beta_1|$  the more rapid is the decay.

By imposing certain restrictions, the AR(1) will reduce to other already known models:


- If  $\beta_1 = 0$ ,  $y_t$  is equivalent to a white noise.
- If  $\beta_1 = 1$  and  $\alpha = 0$ ,  $y_t$  is a random walk.
- If  $\beta_1 = 1$  and  $\alpha \neq 0$ ,  $y_t$  is a random walk with drift.

In general, a smaller persistence parameter results in a quicker adjustment to the *unconditional mean* of the process.

The autocorrelation and partial autocorrelation functions of the AR(1) process have three distinctive features:

- $\rho_1 = \pi_1 = \beta_1$ . That is, the persistence parameter is also the autocorrelation and the partial autocorrelation coefficient.
- The autocorrelation function decreases exponentially toward zero, and the decay is faster when the persistence parameter is smaller.
- The partial autocorrelation function is characterized by only one spike  $\pi_1 \neq 0$ , and the remaining  $\pi_k = 0, \forall k > 1$ .

To illustrate, let's generate a series of 120 observations that follow the process:  $y_t = 0.8y_{t-1} + \varepsilon_t$ , where  $y_0 = 0$  and  $\varepsilon \sim N(0, 1)$ , and plot the ACF and PACF of this series.



#### 4.1.2 AR(2)

Consider the second-order autoregression:

$$y_t = \alpha + \beta_1 y_{t-1} + \beta_2 y_{t-2} + \varepsilon_t$$

where  $\alpha$  is a constant term;  $\beta_1 + \beta_2$  is the persistence measure; and  $\varepsilon_t$  is a white noise process.

In what follows, the necessary (1 and 2) and sufficient (3 and 4) conditions for an  $AR(2)$  process to be covariance stationary are:

1.  $|\beta_2| < 1$
2.  $|\beta_1| < 2$
3.  $\beta_1 + \beta_2 < 1$
4.  $\beta_2 - \beta_1 < 1$

The autocorrelation functions of the  $AR(2)$  process have the following distinctive features:

- $\rho_1 = \pi_1$  (which is true for any  $AR(p)$  process), and  $\pi_2 = \beta_2$ .
- The autocorrelation function decreases toward zero. The path, however, varies depending on the values of  $\beta_1$  and  $\beta_2$ . Nonetheless, the decay is faster when the persistence measure is smaller.
- The partial autocorrelation function is characterized by only two spikes  $\pi_1 \neq 0$  and  $\pi_2 \neq 0$ , and the remaining  $\pi_k = 0, \forall k > 2$ .

## 4.2 Forecasting

The optimal one-step-ahead forecast:

$$y_{t+1|t} = E(y_{t+1}|\Omega_t) = E(\alpha + \beta_1 y_t + \varepsilon_{t+1}) = \alpha + \beta_1 y_t$$

The one-step-ahead forecast error:

$$e_{t+1|t} = y_{t+1} - y_{t+1|t} = \alpha + \beta_1 y_t + \varepsilon_{t+1} - (\alpha + \beta_1 y_t) = \varepsilon_{t+1}$$

The one-step-ahead forecast variance:

$$\sigma_{t+1|t}^2 = Var(y_{t+1}|\Omega_t) = E(e_{t+1|t}^2) = E(\varepsilon_{t+1}^2) = \sigma_\varepsilon^2$$

The one-step-ahead (95%) interval forecast:

$$y_{t+1|t} \pm z_{.025} \sigma_{t+1|t} = y_{t+1|t} \pm 1.96 \sigma_\varepsilon$$

The optimal two-step-ahead forecast:

$$y_{t+2|t} = E(y_{t+2}|\Omega_t) = E(\alpha + \beta_1 y_{t+1} + \varepsilon_{t+2}) = \alpha(1 + \beta_1) + \beta_1^2 y_t$$

The two-step-ahead forecast error:

$$\begin{aligned} e_{t+2|t} &= y_{t+2} - y_{t+2|t} \\ &= \alpha(1 + \beta_1) + \beta_1^2 y_t + \beta_1 \varepsilon_{t+1} + \varepsilon_{t+2} - [\alpha(1 + \beta_1) + \beta_1^2 y_t] \\ &= \beta_1 \varepsilon_{t+1} + \varepsilon_{t+2} \end{aligned}$$

The two-step-ahead forecast variance:

$$\begin{aligned} \sigma_{t+2|t}^2 &= Var(y_{t+2}|\Omega_t) \\ &= E(e_{t+2|t}^2) = E(\beta_1 \varepsilon_{t+1} + \varepsilon_{t+2})^2 = \sigma_\varepsilon^2(1 + \beta_1^2) \end{aligned}$$

The two-step-ahead (95%) interval forecast:

$$y_{t+2|t} \pm z_{.025} \sigma_{t+2|t} = y_{t+2|t} \pm 1.96 \sigma_\varepsilon \sqrt{1 + \beta_1^2}$$

The optimal h-step-ahead forecast:

$$y_{t+h|t} = E(y_{t+h}|\Omega_t) = E(\alpha + \beta_1 y_{t+h-1} + \varepsilon_{t+1}) = \alpha \sum_{j=0}^{h-1} \beta_1^j + \beta_1^h y_t$$

The h-step-ahead forecast error:

$$e_{t+h|t} = y_{t+h} - y_{t+h|t} = \sum_{j=0}^{h-1} \beta_1^j \varepsilon_{t+h-j}$$

The h-step-ahead forecast variance:

$$\sigma_{t+h|t}^2 = \text{Var}(y_{t+h}|\Omega_t) = E(e_{t+h|t}^2) = \sigma_\varepsilon^2 \sum_{j=0}^{h-1} \beta_1^{2j}$$

The h-step-ahead (95%) interval forecast:

$$y_{t+h|t} \pm z_{.025} \sigma_{t+h|t} = y_{t+1|t} \pm 1.96 \sigma_\varepsilon \sqrt{\sum_{j=0}^{h-1} \beta_1^{2j}}$$

In a covariance-stationary process, i.e. when  $|\beta_1| < 1$ , as  $h \rightarrow \infty$ :

The optimal point forecast:

$$y_{t+h|t} = \frac{\alpha}{1 - \beta_1}$$

The forecast variance:

$$\sigma_{t+h|t}^2 = \frac{\sigma_\varepsilon^2}{1 - \beta_1^2}$$

The (95%) interval forecast:

$$y_{t+h|t} \pm z_{.025} \sigma_{t+h|t} = \frac{\alpha}{1 - \beta_1} \pm 1.96 \frac{\sigma_\varepsilon}{\sqrt{1 - \beta_1^2}}$$

The optimal one-step-ahead forecast:

$$\begin{aligned} y_{t+1|t} &= E(y_{t+1}|\Omega_t) \\ &= E(\alpha + \beta_1 y_t + \beta_2 y_{t-1} + \varepsilon_{t+1}) = \alpha + \beta_1 y_t + \beta_2 y_{t-1} \end{aligned}$$

The one-step-ahead forecast error:

$$\begin{aligned} e_{t+1|t} &= y_{t+1} - y_{t+1|t} \\ &= \alpha + \beta_1 y_t + \beta_2 y_{t-1} + \varepsilon_{t+1} - (\alpha + \beta_1 y_t + \beta_2 y_{t-1}) = \varepsilon_{t+1} \end{aligned}$$

The one-step-ahead forecast variance:

$$\sigma_{t+1|t}^2 = \text{Var}(y_{t+1}|\Omega_t) = E(e_{t+1|t}^2) = E(\varepsilon_{t+1}^2) = \sigma_\varepsilon^2$$

The one-step-ahead (95%) interval forecast:

$$y_{t+1|t} \pm z_{.025} \sigma_{t+1|t} = y_{t+1|t} \pm 1.96 \sigma_\varepsilon$$

The optimal two-step-ahead forecast:

$$\begin{aligned} y_{t+2|t} &= E(y_{t+2}|\Omega_t) = E(\alpha + \beta_1 y_{t+1} + \beta_2 y_t + \varepsilon_{t+1}) \\ &= \alpha(1 + \beta_1) + (\beta_1^2 + \beta_2) y_t + \beta_1 \beta_2 y_{t-1} \end{aligned}$$

The two-step-ahead forecast error:

$$\begin{aligned} e_{t+2|t} &= y_{t+2} - y_{t+2|t} = \alpha + \beta_1 y_{t+1} + \beta_2 y_t + \varepsilon_{t+2} \\ &\quad - (\alpha + \beta_1 y_{t+1|t} + \beta_2 y_t) = \beta_1 \varepsilon_{t+1} + \varepsilon_{t+2} \end{aligned}$$

The two-step-ahead forecast variance:

$$\sigma_{t+2|t}^2 = \text{Var}(y_{t+2}|\Omega_t) = E(e_{t+2|t}^2) = E(\beta_1 \varepsilon_{t+1} + \varepsilon_{t+2})^2 = \sigma_\varepsilon^2(1 + \beta_1^2)$$

The two-step-ahead (95%) interval forecast:

$$y_{t+2|t} \pm z_{.025} \sigma_{t+2|t} = y_{t+2|t} \pm 1.96 \sigma_\varepsilon \sqrt{1 + \beta_1^2}$$

The optimal h-step-ahead forecast (iterated method):

$$\begin{aligned} y_{t+1|t} &= \alpha + \beta_1 y_t + \beta_2 y_{t-1} \\ y_{t+2|t} &= \alpha + \beta_1 y_{t+1|t} + \beta_2 y_t \\ y_{t+3|t} &= \alpha + \beta_1 y_{t+2|t} + \beta_2 y_{t+1|t} \\ &\vdots \\ y_{t+h|t} &= \alpha + \beta_1 y_{t+h-1|t} + \beta_2 y_{t+h-2|t} \end{aligned}$$

The h-step-ahead forecast error:

$$e_{t+h|t} = y_{t+h} - y_{t+h|t} = \varepsilon_{t+h} + \beta_1 e_{t+h-1|t} + \beta_2 e_{t+h-2|t}$$

The h-step-ahead forecast variance:

$$\begin{aligned} \sigma_{t+h|t}^2 &= \text{Var}(y_{t+h}|\Omega_t) = E(e_{t+h|t}^2) \\ &= \sigma_\varepsilon^2 + \beta_1^2 \text{Var}(e_{t+h-1|t}) + \beta_2^2 \text{Var}(e_{t+h-2|t}) \\ &\quad + 2\beta_1\beta_2 \text{Cov}(e_{t+h-1|t}, e_{t+h-2|t}) \end{aligned}$$

Note that, in general, formulas for  $\sigma_{t+1|t}^2, \sigma_{t+2|t}^2, \dots, \sigma_{t+h|t}^2$  are the same for any  $AR(p)$ ,  $p \geq h-1$ .

The h-step-ahead (95%) interval forecast:

$$y_{t+h|t} \pm z_{.025} \sigma_{t+h|t} = y_{t+1|t} \pm 1.96 \sigma_{t+h|t}$$

The optimal h-step-ahead forecast:

$$y_{t+h|t} = E(y_{t+h}|\Omega_t) = \alpha + \beta_1 y_{t+h-1|t} + \beta_2 y_{t+h-2|t} + \dots + \beta_p y_{t+h-p|t}$$

The h-step-ahead forecast error:

$$e_{t+h|t} = \varepsilon_{t+h} + \beta_1 e_{t+h-1|t} + \beta_2 e_{t+h-2|t} + \cdots + \beta_p e_{t+h-p|t}$$

The h-step-ahead forecast variance:

$$\begin{aligned} \sigma_{t+h|t}^2 &= \text{Var}(y_{t+h}|\Omega_t) = E(e_{t+h|t}^2) \\ &= \sigma_\varepsilon^2 + \sum_{i=1}^p \beta_i^2 \text{Var}(e_{t+h-i|t}) + 2 \sum_{i \neq j} \beta_i \beta_j \text{Cov}(e_{t+h-i|t}, e_{t+h-j|t}) \end{aligned}$$

The h-step-ahead (95%) interval forecast:

$$y_{t+h|t} \pm z_{.025} \sigma_{t+h|t} = y_{t+h|t} \pm 1.96 \sigma_{t+h|t}$$

## Chapter 5

# Vector Autoregression



## Chapter 6

# Threshold Autoregression

# Forecast Assessment

## Chapter 7

# Forecast Evaluation

## Chapter 8

# Forecast Combination

# Appendix: Introduction to R

R is a programming language for data analysis and visualisation. Here I introduce basic commands that should facilitate your understanding of R. You can further enhance your skillset using numerous online resources, as well as your own trial-and-error. To the extent that new features are added to R on a daily basis, there are virtually no limits to how far you can advance your knowledge of this programming language.

We will work in RStudio—the go-to interface for R (as R itself is not an overly user-friendly platform). Thus, you will need to have installed both, R and RStudio on your device (the latter will ‘find’ and connect with the former on its own). R is available from CRAN, and RStudio is available from RStudio.

## 8.1 Data Management

There are a number of ways in which we can work with data in R. Let’s begin with matrices.

Consider a string of observations:

```
a <- c(1,0,4,3,2,6)
a
```

```
## [1] 1 0 4 3 2 6
```

A *string*, unlike a *vector*, has no dimensions. But we can transform it to a  $n \times 1$  vector using the `as.matrix()` function:

```
b <- as.matrix(a)
b
```

```
##      [,1]
## [1,]    1
## [2,]    0
## [3,]    4
```

```
## [4,]    3
## [5,]    2
## [6,]    6
```

The result is a  $6 \times 1$  vector, or a column matrix. To obtain a  $1 \times 6$  vector, or a row matrix, we *transpose* the foregoing vector using the `t()` function:

```
bt <- t(b)
bt

##      [,1] [,2] [,3] [,4] [,5] [,6]
## [1,]    1    0    4    3    2    6

dim(bt)

## [1] 1 6
```

We can create any  $n \times k$  matrix, using the `matrix()` function. For example, consider a  $3 \times 2$  matrix:

```
B <- matrix(a,nrow=3,ncol=2)
B

##      [,1] [,2]
## [1,]    1    3
## [2,]    0    2
## [3,]    4    6
```

We can add column names and row names to this matrix:

```
colnames(B) <- c("c1","c2")
rownames(B) <- c("r1","r2","r3")
B

##      c1 c2
## r1   1  3
## r2   0  2
## r3   4  6
```

If, at this point, we would like to only work with, say, the first column of the matrix, we can call it using its column number, (1), or the column name, ("c1"), as follows:

```
B[, "c1"]

## r1 r2 r3
##  1  0  4
```

Similarly, if we want to refer to a matrix element, say  $b_{3,2}$ , we can do this as follows:

```
B[3,2]
```

```
## [1] 6
```

Matrix multiplication is done using `%*%` command, granted that the two matrices are compatible. For example, we obtain a product of matrix  $B$  and a new  $2 \times 1$  vector,  $d$ , as follows:

```
d <- as.matrix(c(5,-2))
Bd <- B%*%d
Bd
```

```
##      [,1]
## r1    -1
## r2    -4
## r3     8
```

We can add columns (and rows) to the existing matrix using a `cbind()` function:

```
c3 <- c(0,1,0)
D <- cbind(B,c3)
D
```

```
##      c1 c2 c3
## r1    1  3  0
## r2    0  2  1
## r3    4  6  0
```

We can invert a(n invertible) matrix using the `solve()` function:

```
Di <- solve(D)
Di
```

```
##              r1 r2      r3
## c1 -1.0000000  0  0.5000000
## c2  0.6666667  0 -0.1666667
## c3 -1.3333333  1  0.3333333
```

## 8.2 Data Visualisation

One of the comparative advantages of R is in its graphing aesthetics. Currently, the best graphs are plotted via the **ggplot2** package. Notably, this package requires that the data are maintained in the `data.frame` or the `data.table` format (for the latter, you need to load the **data.table** package). Let's create a `data.table` object and observe its few lines:

```
set.seed(1)
x <- runif(120,0,2)
y <- 0.2+0.7*x+rnorm(120)

library(data.table)
```


```
dt <- data.table(y=y,x=x)
dt

##           y           x
##  1:  2.9733299 0.53101733
##  2:  0.6817335 0.74424780
##  3:  1.6917341 1.14570673
##  4:  1.4994931 1.81641558
##  5: -0.2609185 0.40336386
## ---
## 116: 0.1835826 0.02615515
## 117: 1.9894321 1.43113213
## 118: 2.4197029 0.20636847
## 119: 1.8521905 0.89256870
## 120: 2.3040499 1.28020209
```

Now, let's load **ggplot2** and generate a simple scatter plot:

```
library(ggplot2)


ggplot(dt,aes(x=x,y=y))+
  geom_point()
```



We can augment this plot in a number of different ways. Here we change the point color to red, add the fitted regression line to the plot, add labels to the figure, and apply a 'classic' background theme:




```
ggplot(dt,aes(x=x,y=y))+
  geom_point(color="red")+
  geom_smooth(method="lm",formula=y~x,se=F,color="gray")+
  labs(title="A scatterplot with fitted regression line",
       x="Treatment Variable",
       y="Outcome Variable",
       caption="Caption: in case if needed.")+
  theme_classic()
```



Caption: in case if needed.

As another example, let's generate a histogram (of the dependent variable):


```
ggplot(dt,aes(x=y))+
  geom_histogram(color="white",fill="red",binwidth=.5)+
  labs(title="A basic histogram")+
  theme_classic()
```



We typically apply a line plot to illustrate a time series (that are ordered by date). In what follows, we add a date column to our data frame and then plot the dependent variable in the chronological order:

```
dt$date <- seq(from=as.Date("2000-01-01"),by="month",along.with=y)

ggplot(dt,aes(x=date,y=y))+
  geom_line(color="red")+
  labs(title="A basic time series plot",
       x="Year",
       y="Outcome Variable")+
  theme_classic()
```



### 8.3 Regression Analysis

To illustrate the OLS regression in R, we apply the previously generated  $x$  and  $y$  as independent and dependent variables. To begin, we obtain the least squares estimator “by hand” as follows:

```
X <- cbind(1,x)
b <- solve(t(X)%*%X)%*%t(X)%*%y
b
```

```
##      [,1]
## 0.3577680
## x 0.5781188
```

This can be easily done using the `lm()` function:

```
ols <- lm(y~x)
ols
```

```
##
## Call:
## lm(formula = y ~ x)
##
## Coefficients:
## (Intercept)          x
##      0.3578      0.5781
```

We can apply the `summary()` function to see the complete set of regression results:

```
summary(ols)

##
## Call:
## lm(formula = y ~ x)
##
## Residuals:
##      Min       1Q   Median       3Q      Max
## -1.9662 -0.5983 -0.1127  0.5639  2.3882
##
## Coefficients:
##              Estimate Std. Error t value Pr(>|t|)
## (Intercept)   0.3578     0.1904   1.879 0.062717 .
## x             0.5781     0.1641   3.522 0.000609 ***
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Residual standard error: 0.9716 on 118 degrees of freedom
## Multiple R-squared:  0.09514,    Adjusted R-squared:  0.08748
## F-statistic: 12.41 on 1 and 118 DF,  p-value: 0.0006091
```