a Direct paths between hubs (15 largest)
Pooled
Birds
Squamates
Cartilaginous fishes
GC −> FreeRate CV
rate per branch −> informative sites/branch
rate per branch −> FreeRate CV
informative sites/branch −> CID vs species tree
rate per branch −> QIRP (P correct)
GC −> informative sites/branch
informative sites/branch −> QIRP (P correct)
QIRP (P correct) −> CID vs species tree
GC −> rate per branch
GC −> QIRP (P correct)
GC −> CID vs species tree
FreeRate CV −> CID vs species tree
FreeRate CV −> QIRP (P correct)
FreeRate CV −> informative sites/branch
rate per branch −> CID vs species tree
−0.5
0.0
0.5
direct path coefficient (SD/SD)
b Genomic predictors: total (points) = direct + indirect
direct (pooled; Birds for chr. length)
indirect via hubs
soft−masked fraction −> GC
soft−masked fraction −> rate per branch
soft−masked fraction −> FreeRate CV
soft−masked fraction −> informative sites/branch
soft−masked fraction −> QIRP (P correct)
soft−masked fraction −> CID vs species tree
distance to contig end −> GC
distance to contig end −> rate per branch
distance to contig end −> FreeRate CV
distance to contig end −> informative sites/branch
distance to contig end −> QIRP (P correct)
distance to contig end −> CID vs species tree
contig span −> GC
contig span −> rate per branch
contig span −> FreeRate CV
contig span −> informative sites/branch
contig span −> QIRP (P correct)
contig span −> CID vs species tree
chicken chr. length −> GC
chicken chr. length −> rate per branch
chicken chr. length −> FreeRate CV
chicken chr. length −> informative sites/branch
chicken chr. length −> QIRP (P correct)
chicken chr. length −> CID vs species tree
−0.4
−0.2
0.0
0.2
0.4
effect of the genomic predictor (SD/SD)