
a Best substitution model
b Rate classes (k) of the best model
k=1: no +R. +H (GHOST) best on 6 loci, counted under its k
c Mean k by data size
BirdsAnc0 (47,126)
SquamataAnc0 (28,908)
CartilaginousFishesAnc0 (14,853)
AnuraAnc0 (6,500)
VertebratesAnc5 (4,691)
RayFinnedFishesAnc05 (2,297)
AnuraAnc1 (1,677)
GymnophionaAnc0 (1,326)
AmphibiansAnc0 (1,223)
RayFinnedFishesAnc66 (818)
VertebratesAnc4 (654)
RayFinnedFishesAnc64 (418)
VertebratesAnc3 (343)
RayFinnedFishesAnc03 (322)
RayFinnedFishesAnc02 (308)
RayFinnedFishesAnc04 (247)
VertebratesAnc1 (212)
RayFinnedFishesAnc00 (190)
VertebratesAnc2 (62)
0%
25%
50%
75%
100%
GTR
TN
HKY
K2P
k=1
k=2
k=3
k=4
k=5
k=6
k=7
sites
2400–2500
2100–2400
1800–2100
1500–1800
1199–1500
6.0
1.0
0–25
25–50
50–100
100–150
150–200
200–300
300–400
400–600
tips (after collapsing)
blank: < 20 loci