

December 22, 2022  
**Delone lattice studies in  $\mathbf{C}^3$ , the space of three complex  
 variables**

LAWRENCE C. ANDREWS<sup>a\*</sup> AND HERBERT J. BERNSTEIN<sup>b</sup>

<sup>a</sup>*Ronin Institute, 9515 NE 137th St, Kirkland, WA, 98034-1820 USA, and* <sup>b</sup>*Ronin  
 Institute, c/o NSLS-II, Brookhaven National Laboratory, Upton, NY, 11973 USA.*

*E-mail: lawrence.andrews@ronininstitute.org*

**Delaunay; Delone; Selling;  $\mathbf{C}^3$**

### Abstract

The Delone scalars are studied in  $\mathbf{C}^3$ , the space of three complex variables.

**Note:** In his later publications, Boris Delaunay used the Russian version of his surname, Delone.

### 1. Introduction

The scalars used by Delaunay (1932) in his formulation of Selling reduction (Selling, 1874) are (in the conventional order)  $b \cdot c$ ,  $a \cdot c$ ,  $a \cdot b$ ,  $a \cdot d$ ,  $b \cdot d$ ,  $c \cdot d$ , where  $d = -a - b - c$ . (As a mnemonic device, observe that the first three terms use  $\alpha$ ,  $\beta$ , and  $\gamma$ , in that order, and the following terms use  $a$ ,  $b$ ,  $c$ , in that order.)

Andrews *et al.* (2019a) chose to represent the scalars in the space  $\mathbf{S}^6$ ,  $\{s_1, s_2, s_3, s_4, s_5, s_6\}$  (defined in the order above), as a way to create a metric space for the measurement of the distance between lattices. They also consider the representation of this space as the space of three complex dimensions,  $\mathbf{C}^3$  or  $\{c_1, c_2, c_3\}$ .

In  $\mathbf{C}^3$ , in terms of the Delone scalars, a vector is  $\{(s_1, s_4), (s_2, s_5), (s_3, s_6)\}$ , where the real and imaginary parts of each are the “opposite” scalars according to the definition of Delaunay (1932) (see Andrews *et al.* (2019b)). As a mnemonic device, note that the complex components involve  $(\alpha, a)$ ,  $(\beta, b)$ , and  $(\gamma, c)$ .

Andrews *et al.* (2019a) considered the matrix representations of the reflections in  $\mathbf{C}^3$  (and in  $\mathbf{S}^6$ ). This paper will describe the boundary transformations at the edges of the fundamental unit of  $\mathbf{C}^3$ . In  $\mathbf{S}^6$ , the fundamental unit is the all negative orthant, which contains only and all of the reduced cells. In  $\mathbf{S}^6$  and  $\mathbf{C}^3$ , the boundaries are found where any scalar (or correspondingly the real or imaginary part) is equal to zero. The rationale for this work was that it might lend insights into the topology of the space of lattices.

## 2. Notation

Complex numbers will be represented in Cartesian format  $(x, y)$ , where  $x$  is the real part and  $y$  is the imaginary part.

We will represent a vector in  $\mathbf{C}^3$  by  $\{(x_1, y_1), (x_2, y_2), (x_3, y_3)\}$  as an alternative to  $\{(s_1, s_4), (s_2, s_5), (s_3, s_6)\}$ .

Next we define the operators in  $\mathbf{C}^3$  that will be used in the matrix descriptions of the transformations at the boundaries of the fundamental unit.

See Table 1.

Operator	Usage	Result	Name
$\mathfrak{M}_r$	$\mathfrak{M}_r(c_j)$	$(-x_j, -x_j + y_j)$	Minus real
$\mathfrak{M}_i$	$\mathfrak{M}_i(c_j)$	$(x_j - y_j, -y_j)$	Minus imag
$\mathfrak{P}_r$	$\mathfrak{P}_r(c_j)$	$(x_j, x_j)$	Plus real
$\mathfrak{P}_i$	$\mathfrak{P}_i(c_j)$	$(y_j, y_j)$	Plus imag
$\Re$	$\Re(c_j)$	$x_j$	Real
$\Im$	$\Im(c_j)$	$y_j$	Imaginary

### 3. Matrices of boundary transformations

For the boundary at  $s_1$ : (the real component of  $c_1$ ).

$$\begin{bmatrix} \mathfrak{M}_r & 0 & 0 \\ \mathfrak{P}_r & i\Re & \Re \\ \mathfrak{P}_r & i\Im & \Im \end{bmatrix} \begin{bmatrix} \mathfrak{M}_r & 0 & 0 \\ \mathfrak{P}_r & i\Im & \Im \\ \mathfrak{P}_r & i\Re & \Re \end{bmatrix} \begin{bmatrix} \mathfrak{M}_r & 0 & 0 \\ \mathfrak{P}_r & \Re & i\Re \\ \mathfrak{P}_r & \Im & i\Im \end{bmatrix} \begin{bmatrix} \mathfrak{M}_r & 0 & 0 \\ \mathfrak{P}_r & \Im & i\Im \\ \mathfrak{P}_r & \Re & i\Re \end{bmatrix}$$

For the boundary at  $s_4$ : (the imaginary component of  $c_1$ ).

$$\begin{bmatrix} \mathfrak{M}_i & 0 & 0 \\ \mathfrak{P}_i & i\Re & \Re \\ \mathfrak{P}_i & i\Im & \Im \end{bmatrix} \begin{bmatrix} \mathfrak{M}_i & 0 & 0 \\ \mathfrak{P}_i & i\Im & \Im \\ \mathfrak{P}_i & i\Re & \Re \end{bmatrix} \begin{bmatrix} \mathfrak{M}_i & 0 & 0 \\ \mathfrak{P}_i & \Re & i\Re \\ \mathfrak{P}_i & \Im & i\Im \end{bmatrix} \begin{bmatrix} \mathfrak{M}_i & 0 & 0 \\ \mathfrak{P}_i & \Im & i\Im \\ \mathfrak{P}_i & \Re & i\Re \end{bmatrix}$$

For the boundary at  $s_2$  (the real component of  $c_2$ ):

$$\begin{bmatrix} i\Re & \mathfrak{P}_r & \Re \\ 0 & \mathfrak{M}_r & 0 \\ i\Im & \mathfrak{P}_r & \Im \end{bmatrix} \begin{bmatrix} i\Im & \mathfrak{P}_r & \Im \\ 0 & \mathfrak{M}_r & 0 \\ i\Re & \mathfrak{P}_r & \Re \end{bmatrix} \begin{bmatrix} \Re & \mathfrak{P}_r & i\Re \\ 0 & \mathfrak{M}_r & 0 \\ \Im & \mathfrak{P}_r & i\Im \end{bmatrix} \begin{bmatrix} \Im & \mathfrak{P}_r & i\Im \\ 0 & \mathfrak{M}_r & 0 \\ \Re & \mathfrak{P}_r & i\Re \end{bmatrix}$$

For the boundary at  $s_5$ : (the imaginary component of  $c_2$ ).

$$\begin{bmatrix} i\Re & \mathfrak{P}_i & \Re \\ 0 & \mathfrak{M}_i & 0 \\ i\Im & \mathfrak{P}_i & \Im \end{bmatrix} \begin{bmatrix} i\Im & \mathfrak{P}_i & \Im \\ 0 & \mathfrak{M}_i & 0 \\ i\Re & \mathfrak{P}_i & \Re \end{bmatrix} \begin{bmatrix} \Re & \mathfrak{P}_i & i\Re \\ 0 & \mathfrak{M}_i & 0 \\ \Im & \mathfrak{P}_i & i\Im \end{bmatrix} \begin{bmatrix} \Im & \mathfrak{P}_i & i\Im \\ 0 & \mathfrak{M}_i & 0 \\ \Re & \mathfrak{P}_i & i\Re \end{bmatrix}$$

For the boundary at  $s_3$  (the real component of  $c_3$ ):

$$\begin{bmatrix} i\Re & \Re & \mathfrak{P}_r \\ i\Im & \Im & \mathfrak{P}_r \\ 0 & 0 & \mathfrak{M}_r \end{bmatrix} \begin{bmatrix} i\Im & \Im & \mathfrak{P}_r \\ i\Re & \Re & \mathfrak{P}_r \\ 0 & 0 & \mathfrak{M}_r \end{bmatrix} \begin{bmatrix} \Re & i\Re & \mathfrak{P}_r \\ \Im & i\Im & \mathfrak{P}_r \\ 0 & 0 & \mathfrak{M}_r \end{bmatrix} \begin{bmatrix} \Im & i\Im & \mathfrak{P}_r \\ \Re & i\Re & \mathfrak{P}_r \\ 0 & 0 & \mathfrak{M}_r \end{bmatrix}$$

For the boundary at  $s_6$  (the imaginary component of  $c_3$ ):

$$\begin{bmatrix} i\Re & \Re & \mathfrak{P}_i \\ i\Im & \Im & \mathfrak{P}_i \\ 0 & 0 & \mathfrak{M}_i \end{bmatrix} \begin{bmatrix} i\Im & \Im & \mathfrak{P}_i \\ i\Re & \Re & \mathfrak{P}_i \\ 0 & 0 & \mathfrak{M}_i \end{bmatrix} \begin{bmatrix} \Re & i\Re & \mathfrak{P}_i \\ \Im & i\Im & \mathfrak{P}_i \\ 0 & 0 & \mathfrak{M}_i \end{bmatrix} \begin{bmatrix} \Im & i\Im & \mathfrak{P}_i \\ \Re & i\Re & \mathfrak{P}_i \\ 0 & 0 & \mathfrak{M}_i \end{bmatrix}$$

#### 4. Basics

The standard representation of the identity operation is

$$\mathbf{c}' = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \mathbf{c}$$

The identity in  $\mathbf{C}^3$  can also be written:

$$\mathbf{c}' = \begin{bmatrix} 1 & 0 & 0 \\ 0 & \Re + i\Im & 0 \\ 0 & 0 & \Re + i\Im \end{bmatrix} \mathbf{c}$$

Delaunay (1932) does not consider the boundary transformations in detail. However, he uses them to define the process of Selling reduction. For example in  $\mathbf{S}^6$ , he lists the following as one of the possible results for a transformation on  $s_1$ .  $\{-s_1, -s_1+s_2, s_1+s_3, s_1+s_5, s_1+s_4, s_1+s_6\}$  The third matrix above (the third boundary transform for  $s_1$ ) implements this operation and interchanges the real part of  $c_3$  and the imaginary part of  $c_2$ .

$$\begin{bmatrix} \Re_r & 0 & 0 \\ \Im_r & \Re & i\Re \\ \Im_r & \Im & i\Im \end{bmatrix}$$

In terms of  $\mathbf{C}^3$ , Delone's alternate transformation for the  $s_1$  boundary would exchange the real of  $c_2$  with the imaginary part of  $c_3$ . That is the fourth matrix in the list for  $s_1$ . The other two transformations for  $s_1$  can be generated from the two we have just displayed by the "exchange operation" (Andrews *et al.*, 2019a) applied to the second and third  $\mathbf{C}^3$  elements. Delone did not describe the latter two transformation, perhaps because even a single transformation was adequate to implement reduction. He already listed two.

#### 5. Summary

The considerable regularity of the transformation matrices (above) highlights one of the important aspects of Selling reduction as used by Delaunay (1932) when compared

to Niggli reduction (Niggli, 1928). Where all the boundaries of the fundamental orthant in  $\mathbf{C}^3$  and  $\mathbf{S}^6$  are basically the same (and indeed are related by reflections), the boundaries formed in Niggli reduction are of multiple types, and the fundamental unit of the space representing Niggli reduced cells ( $\mathbf{G}^6$ , see Andrews & Bernstein (2014)) is non-convex.

The matrices display one of the properties of  $\mathbf{C}^3$  that make it a useful representation. For the scalar being transformed, it and its opposite scalar create a unique row in each matrix. Those rows contain only the minus operator and zeros, highlighting the unique relationship of that pair of scalars.

Several aspects of  $\mathbf{C}^3$  are obvious from inspecting the matrices above. First, we see that each boundary has four possible transformations that can be applied. Because each of the transformations at boundaries are self-inverse, they are the same transformations that would be used in the process of cell (lattice) reduction. Delaunay (1932) and Delone *et al.* (1975) give only two choices. Presumably for simplicity they did not list the transformations that used the "exchange" operator (see (Andrews *et al.*, 2019a)).

## 6. Availability of code

The  $C^{++}$  code for  $\mathbf{C}^3$  is available in github.com, in <https://github.com/duck10/LatticeRepLib.git>.

## Acknowledgements

Careful copy-editing and corrections by Frances C. Bernstein are gratefully acknowledged. Our thanks to Jean Jakoncic and Alexei Soares for helpful conversations and access to data and facilities at Brookhaven National Laboratory.

## Funding information

Funding for this research was provided in part by: US Department of Energy Offices

of Biological and Environmental Research and of Basic Energy Sciences (grant No. DE-AC02-98CH10886; grant No. E-SC0012704); U.S. National Institutes of Health (grant No. P41RR012408; grant No. P41GM103473; grant No. P41GM111244; grant No. R01GM117126, grant No. 1R21GM129570); Dectris, Ltd.

### References

- Andrews, L. C. & Bernstein, H. J. (2014). J. Appl. Cryst. **47**(1), 346 – 359.
- Andrews, L. C., Bernstein, H. J. & Sauter, N. K. (2019a). Acta Cryst. **A75**.  
<https://doi.org/10.1107/S2053273318015413>.
- Andrews, L. C., Bernstein, H. J. & Sauter, N. K. (2019b). Acta Cryst. **A75**(3), 593–599.
- Delaunay, B. N. (1932). Z. Krist. **84**, 109 – 149.
- Delone, B. N., Galiulin, R. V. & Shtogrin, M. I. (1975). J. Sov. Math. **4**(1), 79 – 156.
- Niggli, P., (1928). *Krystallographische und Strukturtheoretische Grundbegriffe*, Handbuch der Experimentalphysik, Vol. 7, part 1. Akademische Verlagsgesellschaft, Leipzig.
- Selling, E. (1874). Journal für die reine und angewandte Mathematik (Crelle's Journal), **1874**(77), 143 – 229.

---

### Synopsis

Selling reduction and Delone reduction are considered in a space of complex variables.

---