

Math 504 Homework: Advanced Linear Algebra

Tai Duc Nguyen

December 6, 2021 (Fall 2021)

1. Scoring Table

Problem # in notes	Problem # in this doc	Problem points	Problem grade
1.1.1	1	3	
1.1.2	2	3	
1.2.1	3	2	
1.5.1	4	1	
1.5.2	5	3	
1.5.3	6	2	
1.5.4	7	3	
1.5.5	8	5	
1.5.6	9	3	
1.5.7	10	5	
1.5.8	11	2	
1.8.1	12	4	
1.8.2	13	5	
2.1.1	14	2	
2.1.2	15	3	
2.1.3	16	2	

Problem # in notes	Problem # in this doc	Problem points	Problem grade
2.2.1	17	1	
2.3.2	18	2	
2.3.3	19	3	
2.5.1	20	2	
2.5.2	21	3	
2.5.3	22	2	
2.5.4	23	4	
2.5.5	24	4	
2.5.6	25	5	
2.5.7	26	4	
2.5.8	27	3	
2.6.1	28	2	
2.6.2	29	5	
2.6.3	30	2	
2.6.4	31	3	
2.6.5	32	3	
2.7.1	33	3	
2.7.2	34	3	
2.7.3	35	5	
2.8.1	36	2	
2.8.2	37	5	
3.4.1	38	3	
3.4.2	39	4	
3.5.1	40	6	
3.5.2	41	4	
Total points		131	

Exercises

Exercise 1. [3] (1.1.1) Let $x \in \mathbb{C}^n$ and $y \in \mathbb{C}^n$. Prove that

$$||x + y||^2 - ||x||^2 - ||y||^2 = \langle x, y \rangle + \langle y, x \rangle = 2 \cdot \operatorname{Re} \langle x, y \rangle.$$

Here, $\operatorname{Re} z$ denotes the real part of any complex number z .

Solution 1.

$$\begin{aligned} ||x + y||^2 &= \sqrt{\langle x + y, x + y \rangle}^2 - \sqrt{\langle x, x \rangle}^2 - \sqrt{\langle y, y \rangle}^2 \\ &= \langle x + y, x + y \rangle - \langle x, x \rangle - \langle y, y \rangle \end{aligned}$$

Due to the fact that dot product has distributive property:

$$\begin{aligned} &= \langle x, x \rangle + \langle x, y \rangle + \langle y, y \rangle + \langle y, x \rangle - \langle x, x \rangle - \langle y, y \rangle \\ &= \langle x, y \rangle + \langle y, x \rangle \end{aligned}$$

Since dot product has commutative property:

$$= 2 \cdot \langle x, y \rangle$$

And because the entire left hand side is all real (sum of squared terms)

$$= 2 \cdot \operatorname{Re} \langle x, y \rangle$$

QED.

Exercise 2. [3] (1.1.2) Prove triangular inequality theorem: Let $x \in \mathbb{C}^n$ and $y \in \mathbb{C}^n$. Then:

(a) The inequality $||x|| + ||y|| \geq ||x + y||$ holds.

(b) This inequality becomes an equality if and only if we have $y = 0$ or $x = \lambda y$ for some nonnegative real λ .

Solution 2. (a) We can prove the inequality: $||x|| + ||y|| \geq ||x + y||$ by squaring both sides.

Left side:

$$(||x|| + ||y||)^2 = ||x||^2 + ||y||^2 + 2 \cdot ||x|| \cdot ||y||$$

Right side:

$$\begin{aligned} ||x + y||^2 &= \langle x + y, x + y \rangle = \langle x, x \rangle + \langle y, y \rangle + \langle x, y \rangle + \langle y, x \rangle \\ &= ||x||^2 + ||y||^2 + \langle x, y \rangle + \langle y, x \rangle \end{aligned}$$

Using the Cauchy-Schwarz inequality: $\|x\| \cdot \|y\| \geq |\langle x, y \rangle|$, we have:

$$2 \cdot \|x\| \cdot \|y\| \geq 2 \cdot |\langle x, y \rangle|$$

Since $|\bar{z}| = |z|$, $z \in \mathbb{C}^n$,

$$2 \cdot \|x\| \cdot \|y\| \geq |\langle x, y \rangle| + |\langle y, x \rangle|$$

Hence, the left hand side is \geq right hand side.

(b)

The case for $y = 0$ is clear because $\|x\| + \|0\| = \|x + 0\| = \|x\|$

The other case if $x = \lambda y$, for some non-negative real λ , we can substitute in the last equation above

The left hand side:

$$\|x\| \cdot \|y\| = \|\lambda y\| \cdot \|y\| = |\lambda| \|y\| \cdot \|y\| = \lambda \|y\|^2$$

The right hand side:

$$|\langle x, y \rangle| = |\langle \lambda y, y \rangle| = \lambda |\langle y, y \rangle| = \lambda \|y\|^2$$

Hence, the left hand side = right hand side.

Exercise 3. [2] (1.2.1) Let (u_1, u_2, \dots, u_k) be an orthogonal k -tuple of nonzero vectors in \mathbb{C}^n . Then, we have $k \leq n$, and prove that we can find $n - k$ further nonzero vectors $u_{k+1}, u_{k+2}, \dots, u_n$ such that (u_1, u_2, \dots, u_n) is an orthogonal basis of \mathbb{C}^n .

Solution 3. Let $U_k = (u_1, u_2, \dots, u_k)$ be an orthogonal k -tuple of nonzero vectors in \mathbb{C}^n and $k < n - 1$.

From this Lemma,

Lemma 1.0.1. Let $k < n$. Let a_1, a_2, \dots, a_k be k vectors in \mathbb{C}^n . Then, there exists a nonzero vector $b \in \mathbb{C}^n$ that is orthogonal to each of a_1, a_2, \dots, a_k .

there exist a non-zero vector n_1 that is orthogonal to each vector u_i in U_k . Hence, $(u_1, u_2, \dots, u_k, n_1)$ forms a new tuple U_{k+1} of $k + 1$ orthogonal vectors in \mathbb{C}^n . Iteratively, we can find $n_1, n_2, n_3, \dots, n_l$ vectors, with $l = n - k$, that together with u_1, u_2, \dots, u_k forms a tuple of n orthogonal vectors $(u_1, u_2, \dots, u_k, n_1, n_2, \dots, n_l)$ in \mathbb{C}^n , which is an orthogonal basis of \mathbb{C}^n . QED.

Exercise 4. [1] (1.5.1) Let $A \in \mathbb{C}^{n \times k}$ be an isometry. Show that $n \geq k$.

Solution 4. Because A is an isometry, columns of A forms a tuple of k -orthogonal vectors in \mathbb{C}^n . If $n < k$, then there would be $l = k - n$ more equations than unknowns, which means that there are l vectors that are not orthogonal with each of the other vectors in the tuple. This contradicts, hence, QED.

Exercise 5. [3] (1.5.2) (a) Prove that the product AB of two isometries $A \in \mathbb{C}^{n \times m}$ and $B \in \mathbb{C}^{m \times k}$ is always an isometry.

(b) Prove that the product AB of two unitary matrices $A \in \mathbb{C}^{n \times n}$ and $B \in \mathbb{C}^{n \times n}$ is always unitary.

(c) Prove that the inverse of a unitary matrix $A \in \mathbb{C}^{n \times n}$ is always unitary.

Solution 5. (a) Let b_1, b_2, \dots, b_k be the columns in matrix B . Then, the product AB can be written as:

$$AB = C = \begin{pmatrix} | & & | \\ Ab_1 & \cdots & Ab_k \\ | & & | \end{pmatrix}$$

For any 2 columns c_i, c_j ($i \neq j$) in C , the dot product between them is:

$$\langle c_i, c_j \rangle = \langle Ab_i, Ab_j \rangle = \langle A, A \rangle \cdot \langle b_i, b_j \rangle = \|A\| \cdot \langle b_i, b_j \rangle$$

But because b_i is orthogonal to b_j , $\langle b_i, b_j \rangle = 0$. Hence,

$$\langle c_i, c_j \rangle = 0$$

Which means C is an isometry. QED.

(b)

$$\begin{aligned} (AB)(AB)^* &= (AB)(B^*A^*) \\ &= A(BB^*)A^* \\ &= AI_nA^* = AA^* = I_n \end{aligned}$$

Hence, the product AB is also unitary

(c) If A^{-1} is unitary, then we must prove:

$$A^{-1}(A^{-1})^* = I$$

From the fact that A is unitary, we have

$$AA^* = I_n$$

Invert both sides:

$$A^{-1}(A^*)^{-1} = I$$

Multiply both sides by A and A^{-1} :

$$A(A^{-1}(A^*)^{-1})A^{-1} = AIA^{-1}$$

Simplifying:

$$(AA^{-1})((A^*)^{-1}A^{-1}) = I$$

$$(A^*)^{-1}A^{-1} = I$$

Take the conjugate transpose of both sides:

$$((A^*)^{-1}A^{-1})^* = A^{-1}(A^{-1})^* = I$$

QED.

Exercise 6. [2] (1.5.3) Let $U \in \mathbb{C}^{n \times n}$ be a unitary matrix.

(a) Prove that $|\det U| = 1$.

(b) Prove that any eigenvalue λ of U satisfies $|\lambda| = 1$.

Solution 6. (a) Since U is unitary:

$$\det(U^*U) = \det(I) = 1$$

$$\det(U^*)\det(U) = 1$$

$$\det(U)^*\det(U) = 1$$

$$|\det(U)|^2 = 1$$

$$|\det(U)| = 1$$

(b) Let λ be an eigen value of U and x be the eigen vector associated with λ :

$$Ux = \lambda x$$

$$(Ux)^*Ux = (\lambda x)^*\lambda x$$

$$x^*U^*Ux = x^*\lambda^*\lambda x$$

$$x^*x = |\lambda|^2 x^*x$$

$$|x|^2 = |\lambda||x|^2$$

$$1 = |\lambda|$$

Exercise 7. [3] (1.5.4) Let $w \in \mathbb{C}^n$ be a nonzero vector. Then, $w^*w = \langle w, w \rangle > 0$. Thus, we can define an $n \times n$ -matrix

$$U_w := I_n - 2(w^*w)^{-1}ww^* \in \mathbb{C}^{n \times n}.$$

This is called a *Householder matrix*.

Show that this matrix U_w is unitary and satisfies $U_w^* = U_w$.

Solution 7. Symmetry:

$$\begin{aligned} U_w^* &= (I_n - 2 \frac{ww^*}{w^*w})^* \\ &= I_n - 2 \frac{(ww^*)^*}{(w^*w)^*} \\ &= I_n - 2 \frac{ww^*}{w^*w} = U_w \end{aligned}$$

Unitary:

$$\begin{aligned} U_w U_w^* &= (I_n - 2 \frac{ww^*}{w^*w})(I_n - 2 \frac{ww^*}{w^*w})^* \\ &= (I_n - 2 \frac{ww^*}{w^*w})(I_n - 2 \frac{ww^*}{w^*w}) \\ &= I_n - 4 \frac{ww^*}{w^*w} + 4 \frac{ww^*ww^*}{w^*ww^*w} \\ &= I_n - 4 \frac{ww^*}{w^*w} + 4 \frac{ww^*}{w^*w} \\ &= I_n \end{aligned}$$

Exercise 8. [5] (1.5.5) Let $S \in \mathbb{C}^{n \times n}$ be a skew-Hermitian matrix.

(a) Prove that the matrix $I_n - S$ is invertible.

[Hint: Show first that the matrix $I_n + S^*S$ is invertible, since each nonzero vector $v \in \mathbb{C}^n$ satisfies $v^* (I_n + S^*S) v = \underbrace{\langle v, v \rangle}_{>0} + \underbrace{\langle Sv, Sv \rangle}_{\geq 0} > 0$. Then, expand the

product $(I_n - S^*)(I_n - S)$.]

(b) Prove that the matrices $I_n + S$ and $(I_n - S)^{-1}$ commute (i.e., satisfy $(I_n + S) \cdot (I_n - S)^{-1} = (I_n - S)^{-1} \cdot (I_n + S)$).

(c) Prove that the matrix $U := (I_n - S)^{-1} \cdot (I_n + S)$ is unitary.

(d) Prove that the matrix $U + I_n$ is invertible.

(e) Prove that $S = (U - I_n) \cdot (U + I_n)^{-1}$.

Solution 8. (a) Let λ be an eigen value of S and v be its corresponding eigen vector. Then:

$$(I - S)v = Iv - Sv = Iv - \lambda v = (1 - \lambda)v$$

Since S is a skew-Hermitian matrix, λ must be purely imaginary, and $1 - \lambda$ cannot be 0. Hence, all the eigen values of $I - S$ cannot be 0, which means $I - S$ is invertible.

(b) We have:

$$(I + S) \cdot (I - S)^{-1} = I - S^{-1} + S + S \cdot (-S)^{-1}$$

and

$$(I - S)^{-1} \cdot (I + S) = I + S - S^{-1} + (-S)^{-1} \cdot S = I - S^{-1} + S + (-S)^{-1} \cdot S$$

From these 2 equations, we have to prove that: $S \cdot (-S)^{-1} = (-S)^{-1} \cdot S$. But this is obvious because:

$$S \cdot (-S)^{-1} = \frac{S}{-S} = -\frac{S}{S} = \frac{-S}{S} = (-S)^{-1} \cdot S$$

(c)

Let $W = I - S$, then, $(I - S)^{-1} = W^{-1}$ and $I + S = W^*$. Hence, $U = W^{-1}W^*$. We are going to prove $UU^* = I$.

$$UU^* = W^{-1}W^*(W^{-1}W^*)^* = W^{-1}W^*W(W^{-1})^*$$

Since W and W^* commutes (part b),

$$UU^* = (W^{-1}W) \cdot (W^*(W^{-1})^*) = I \cdot I = I$$

(d) Following the proof in part a, we can see that the matrix $I + S$ is also invertible because all of its eigen values cannot be 0 since $1 + \lambda$ (where λ is purely imaginary). Hence, the matrix U from part c above also have all of its eigenvalues be non-zero since its determinant is a product of two non-zero determinant: $\det((I - S)^{-1})$ and $\det(I + S)$. Therefore, again, let μ be an eigen value of U and v be its corresponding eigen vector. Then:

$$(U + I)v = Uv + Iv = \mu v + Iv = (\mu + 1)v$$

Since μ is non-zero, $\mu + 1$ is non-zero and $U + I$ is invertible.

(e) From the previous proofs, we can construct the following equation:

$$U(I - S) = (I - S)^{-1}(I + S)(I - S) = I \cdot (I + S) = I + S$$

Hence,

$$U - US = I + S$$

$$U - I = S + US$$

$$U - I = S(I + U)$$

Multiply both sides by $(I + U)^{-1}$:

$$(U - I)(I + U)^{-1} = S(I + U)(I + U)^{-1} = S$$

Exercise 9. [3] (1.5.6) Let $A \in \mathbb{C}^{n \times n}$ be a matrix. Prove the following:

(a) If A is unitary, then the matrix λA is unitary for each $\lambda \in \mathbb{C}$ satisfying

$$|\lambda| = 1.$$

(b) The matrix $\lambda A + I_n$ is invertible for all but finitely many $\lambda \in \mathbb{C}$.

[Hint: The determinant $\det(\lambda A + I_n)$ is a polynomial function in λ .]

(c) The set $\{U \in U_n(\mathbb{C}) \mid U + I_n \text{ is invertible}\}$ is dense in $U_n(\mathbb{C})$. (That is, each unitary matrix in $U_n(\mathbb{C})$ can be written as a limit $\lim_{k \rightarrow \infty} U_k$ of a sequence of unitary matrices U_k such that $U_k + I_n$ is invertible for each k .)

Solution 9. (a) We have:

$$\begin{aligned}\lambda A(\lambda A)^* &= \lambda A A^* \bar{\lambda} \\ &= \lambda I \bar{\lambda} = |\lambda|^2 I\end{aligned}$$

Hence, $|\lambda|^2 I = I$ only if $|\lambda|^2 = 1$, which means $|\lambda| = 1$

(b) Let μ be the eigen value of the matrix A and v be its corresponding eigen vector:

$$(\lambda A + I)v = \lambda Av + Iv = \lambda \mu v + Iv = (\lambda \mu + 1)v$$

In order for $\lambda A + I$ to be invertible, $(\lambda \mu + 1)$ has to be non zero. Hence, $\lambda \neq -1 - \mu$

(c) In order to show that the set $D = \{U \in U_n(\mathbb{C}) \mid U + I_n \text{ is invertible}\}$ is dense in $U_n(\mathbb{C})$, we will show that the complement of such set is nowhere dense in $U_n(\mathbb{C})$.

The complement of such set is that:

$$\bar{D} = \{U \in U_n(\mathbb{C}) \mid U + I_n \text{ is NOT invertible}\}$$

Again, let λ be an eigen value of U and v its corresponding eigen vector. Then:

$$(U + I)v = Uv + Iv = \lambda v + Iv = (\lambda + 1)v$$

Since λ is an eigen value of U , $|\lambda| = 1$. Also, because the matrix $U + I$ is not invertible, $\lambda + 1 = 0$. Hence, λ has to be -1 . And due to the fact that λ spans all of \mathbb{C} , the complementary set \bar{D} is not dense at all. Therefore, the set D is dense in $U_n(\mathbb{C})$.

Exercise 10. [5] (1.5.7) A *Pythagorean triple* is a triple (p, q, r) of positive integers satisfying $p^2 + q^2 = r^2$. (In other words, it is a triple of positive integers that are the sides of a right-angled triangle.) Two famous Pythagorean triples are $(3, 4, 5)$ and $(5, 12, 13)$.

(a) Prove that a triple (p, q, r) of positive integers is Pythagorean if and only if the matrix $\begin{pmatrix} p/r & -q/r \\ q/r & p/r \end{pmatrix}$ is unitary.

(b) Let $\begin{pmatrix} a & b \\ c & d \end{pmatrix}$ be any unitary matrix with rational entries. Assume that a and c are positive, and write a and c as p/r and q/r for some positive integers p, q, r . Show that (p, q, r) is a Pythagorean triple.

(c) Find infinitely many Pythagorean triples that are pairwise non-proportional (i.e., no two of them are obtained from one another just by multiplying all three entries by the same number).

[Hint: Use the $S \mapsto U$ construction from Exercise 8.]

Solution 10. (a) Let U be the matrix $\begin{pmatrix} p/r & -q/r \\ q/r & p/r \end{pmatrix}$. The product of $UU^* = UU^{-1}$ is:

$$\begin{pmatrix} p/r & -q/r \\ q/r & p/r \end{pmatrix} \times \begin{pmatrix} p/r & q/r \\ -q/r & p/r \end{pmatrix} = \begin{pmatrix} p^2/r^2 + q^2/r^2 & 0 \\ 0 & q^2/r^2 + p^2/r^2 \end{pmatrix}$$

But since $p^2 + q^2 = r^2$, which means (divide both sides by r^2) $q^2/r^2 + p^2/r^2 = 1$,

$$UU^* = UU^{-1} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} = I$$

This is the same for U^*U . Hence, $U^*U = UU^* = I$ and U is unitary.

The reverse direction is similar. If $UU^* = I$, then $q^2/r^2 + p^2/r^2$ has to be 1, so $p^2 + q^2 = r^2$.

(b)

Exercise 11. [2] (1.5.8) Let $A, B \in \mathbb{C}^{n \times n}$ be two skew-Hermitian matrices. Show that $AB - BA$ is again skew-Hermitian.

Solution 11. We have: $A^* = -A$ and $B^* = -B$. Then:

$$(AB - BA)^* = B^*A^* - A^*B^* = (-B)(-A) - (-A)(-B) = BA - AB$$

Exercise 12. [4] (1.8.1) Let $A \in \mathbb{C}^{n \times m}$ satisfy $n \geq m$ and $\text{rank } A = m$. Prove that there exists exactly one QR factorization (Q, R) of A such that the diagonal entries of R are positive reals.

Solution 12. Let $A \in \mathbb{C}^{n \times m}$, $n \geq m$ and $\text{rank } A = m$ (full rank). Suppose it has at least these 2 decompositions:

$$A = Q_1 R_1 = Q_2 R_2$$

Here, Q_1, Q_2 are isometries of size $n \times m$ and R_1, R_2 are invertible, upper-triangular of size $n \times n$. Shifting things around in the equation above, we get:

$$\begin{cases} Q_2^T Q_1 = R_2 R_1^T (*) \\ Q_1^T Q_2 = R_1 R_2^T (**) \end{cases}$$

In (*), since both R_2 and R_1 are upper-triangular, their product is upper-triangular. Hence $Q_2^T Q_1$ is upper-triangular. However, in (**), we can transpose both sides and get:

$$(Q_1^T Q_2)^T = (R_1 R_2^T)^T$$

$$Q_1 Q_2^T = Q_2^T Q_1 = R_1^T R_2$$

The product $R_1^T R_2$ results in a lower-triangular matrix. Hence, $Q_2^T Q_1 = R_2 R_1^T$ is both lower and upper-triangular, which means it has to be a diagonal matrix D . In addition, the equation $Q_1 R_1 = Q_2 R_2$ can also be written as:

$$Q_1 (R_1 R_2^T) = Q_2 R_2 R_2^T = Q_2$$

$$Q_1 D = Q_2$$

But, since:

$$I = (Q_1 D)^T (Q_1 D) = D^T Q_1^T Q_1 D$$

Because Q_1 is an isometry, $Q_1^T Q_1 = I$

$$I = D^T D = D^2$$

Therefore, D has to be a diagonal matrix with entries ± 1 . However, if we constrain that all the diagonal entries of R_1 and R_2 to be positive, the same must be true for those of D . Hence, $D = I$, which means that

$$Q_1 D = Q_1 I = Q_1 = Q_2$$

and,

$$R_1 = R_2$$

This means that the pair Q, R is unique.

Exercise 13. [5] (1.8.2) Prove Theorem 1.0.2.

[Hint: Reduce both cases $n > m$ and $n < m$ to the case $n = m$.]

Theorem 1.0.2 (QR factorization, unitary version). Let $A \in \mathbb{C}^{n \times m}$. Then, there exist a unitary matrix $Q \in \mathbb{C}^{n \times n}$ and an upper-triangular matrix $R \in \mathbb{C}^{n \times m}$ such that $A = QR$. Here, a rectangular matrix $R \in \mathbb{C}^{n \times m}$ is said to be *upper-triangular* if and only if it satisfies

$$R_{i,j} = 0 \quad \text{for all } i > j.$$

Solution 13.

Theorem 1.0.3 (QR factorization, isometry version). Let $A \in \mathbb{C}^{n \times m}$ satisfy $n \geq m$. Then, there exist an isometry $Q \in \mathbb{C}^{n \times m}$ and an upper-triangular matrix $R \in \mathbb{C}^{m \times m}$ such that $A = QR$.

We must consider 3 cases: $n < m$, $n > m$, and $n = m$.

Case 1 ($n = m$):

Using theorem 1.0.3, We can see that $A = QR$, in which Q is an isometry of size $n \times n$ (in other words, an unitary matrix) and R is an upper-triangular matrix also of size $n \times n$.

Hence, let Q_1, R_1 and Q_2, R_2 are 2 of decomposition of A . Then: $A = Q_1 R_1 = Q_2 R_2$.

Using the proof from the previous exercise 12, we know that:

$$Q_2 = Q_1 D$$

and,

$$R_2 = D R_1$$

in which D is a diagonal matrix with entries ± 1 . If we constrain the diagonal entries of R to be positive then the decomposition is unique.

Case 2 ($n < m$):

We have: $A = QR$, in which, Q is $n \times n$ and R is $n \times m$. We can write R as

$$A = QR = Q \begin{bmatrix} | & | \\ R_1 & N_1 \\ | & | \end{bmatrix} = [R_1 N_1] \text{ (in short)}$$

, where: R_1 is a square $n \times n$ matrix and N_1 is a rectangular $n \times m - n$ matrix. Assume that the matrix A has another QR decomposition, resulting in:

$$A = Q_1 [R_1 N_1] = Q_2 [R_2 N_2]$$

$$[(Q_1 R_1) (Q_1 N_1)] = [(Q_2 R_2) (Q_2 N_2)]$$

We can see that:

$$\begin{cases} Q_1 R_1 = Q_2 R_2 \\ Q_1 N_1 = Q_2 N_2 \end{cases}$$

From the previous case, we know that $Q_2 = Q_1 S$ and $R_2 = D R_1$. Hence, we can substitute in:

$$Q_1 N_1 = Q_1 D N_2$$

$$N_1 = D N_2$$

$$D^T N_1 = D N_1 = N_2$$

Hence, we have proved the theorem for $n < m$

Case 3 ($n > m$):

Similar to case 2, suppose the matrix A has decomposition $A = QR$, in which Q is an isometry of size $n \times n$ and R is an upper-triangular matrix of size $n \times m$. Since $n > m$, we can write Q as:

$$Q = [Q'M']$$

, where Q' is $n \times m$ and M' is $n \times n - m$. And we can write R as:

$$R = \begin{bmatrix} R' \\ 0 \end{bmatrix}$$

, with R' is $m \times m$.

Hence, we write the decomposition as:

$$A = [Q'M'] \begin{bmatrix} R' \\ 0 \end{bmatrix} = Q'R'$$

Similar to the proof in the previous exercise 12, assume A has 2 decomposition Q_1, R_1 and Q_2, R_2 , then:

$$A = Q_1R_1 = Q'_1R'_1 = Q'_2R'_2 = Q_2R_2$$

, and there exist a diagonal matrix D in which its entries are ± 1 so that:

$$\begin{cases} Q'_2 = Q'_1D \\ R'_2 = DR'_1 \end{cases}$$

Hence, we have proved the theorem for $n > m$

Exercise 14. 2 (2.1.1) Prove that the two matrices $\begin{pmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 \end{pmatrix}$ and

$\begin{pmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}$ are not similar.

Solution 14. Let $A = \begin{pmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 \end{pmatrix}$ and $B = \begin{pmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}$

Breaking both matrices into their Jordan blocks:

$$A = \left(\begin{array}{cc|cc} 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ \hline 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 \end{array} \right)$$

$$B = \left(\begin{array}{ccc|c} 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 \\ \hline 0 & 0 & 0 & 0 \end{array} \right)$$

Since the Jordan blocks of A have different sizes compared to those of B , A and B cannot be similar.

Exercise 15. [3] (2.1.2) Let $A \in \mathbb{C}^{n \times n}$ be a matrix that is similar to some unitary matrix. Prove that A^{-1} is similar to A^* .

Solution 15. Since A is similar to some unitary matrix U , we can write A as:

$$A = PUP^{-1}$$

Also, we can write U as:

$$U = P^{-1}AP$$

Now, take the conjugate transpose of both sides in the first equation:

$$A^* = (PUP^{-1})^* = (P^{-1})^*U^*P^*$$

And take the inverse of the second equation:

$$U^{-1} = (P^{-1}AP)^{-1} = P^{-1}A^{-1}P$$

Since U is unitary, we can substitute U^{-1} into U^* to obtain:

$$A^* = (P^{-1})^*(P^{-1}A^{-1}P)P^*$$

$$A^* = (P^{-1})^*P^{-1}A^{-1}PP^* = (PP^*)^{-1}A^{-1}PP^* = Q^{-1}A^{-1}Q$$

Hence, A^* is similar to A^{-1}

Exercise 16. [2] (2.1.3) Prove Proposition 1.0.4.

Proposition 1.0.4. Let \mathbb{F} be a field. Let $n \in \mathbb{N}$. For each $i \in [n]$, let A_i and B_i be two $n_i \times n_i$ -matrices (for some $n_i \in \mathbb{N}$) satisfying $A_i \sim B_i$. Then,

$$\begin{pmatrix} A_1 & 0 & \cdots & 0 \\ 0 & A_2 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & A_n \end{pmatrix} \sim \begin{pmatrix} B_1 & 0 & \cdots & 0 \\ 0 & B_2 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & B_n \end{pmatrix}.$$

Solution 16. Since $A_i \sim B_i$, $A_i = P_i B_i P_i^{-1}$. We can rewrite the block diagonal matrix A as:

$$A = \begin{pmatrix} P_1 B_1 P_1^{-1} & 0 & \cdots & 0 \\ 0 & P_2 B_2 P_2^{-1} & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & P_n B_n P_n^{-1} \end{pmatrix}$$

which is the same as:

$$\begin{aligned} A &= \begin{pmatrix} P_1 & 0 & \cdots & 0 \\ 0 & P_2 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & P_n \end{pmatrix} \begin{pmatrix} B_1 P_1^{-1} & 0 & \cdots & 0 \\ 0 & B_2 P_2^{-1} & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & B_n P_n^{-1} \end{pmatrix} \\ &= \begin{pmatrix} P_1 & 0 & \cdots & 0 \\ 0 & P_2 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & P_n \end{pmatrix} \begin{pmatrix} B_1 & 0 & \cdots & 0 \\ 0 & B_2 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & B_n \end{pmatrix} \begin{pmatrix} P_1^{-1} & 0 & \cdots & 0 \\ 0 & P_2^{-1} & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & P_n^{-1} \end{pmatrix} \end{aligned}$$

Hence,

$$\begin{pmatrix} A_1 & 0 & \cdots & 0 \\ 0 & A_2 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & A_n \end{pmatrix} \sim \begin{pmatrix} B_1 & 0 & \cdots & 0 \\ 0 & B_2 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & B_n \end{pmatrix}.$$

Exercise 17. [1] Prove Proposition 1.0.5.

Proposition 1.0.5. Let $n \in \mathbb{N}$. For each $i \in [n]$, let $A_i \in \mathbb{C}^{n_i \times n_i}$ and $B_i \in \mathbb{C}^{n_i \times n_i}$ be two $n_i \times n_i$ -matrices (for some $n_i \in \mathbb{N}$) satisfying $A_i \stackrel{\text{us}}{\sim} B_i$. Then,

$$\begin{pmatrix} A_1 & 0 & \cdots & 0 \\ 0 & A_2 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & A_n \end{pmatrix} \stackrel{\text{us}}{\sim} \begin{pmatrix} B_1 & 0 & \cdots & 0 \\ 0 & B_2 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & B_n \end{pmatrix}.$$

Solution 17. Similar to the prove for the exercise 16 above, the final equation becomes:

$$A = \begin{pmatrix} P_1 & 0 & \cdots & 0 \\ 0 & P_2 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & P_n \end{pmatrix} \begin{pmatrix} B_1 & 0 & \cdots & 0 \\ 0 & B_2 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & B_n \end{pmatrix} \begin{pmatrix} P_1^* & 0 & \cdots & 0 \\ 0 & P_2^* & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & P_n^* \end{pmatrix}$$

We can see that the left matrix and the right matrix are composed of diagonal blocks in which all of them are unitary. Hence, both the left and the right matrix are unitary. QED.

Exercise 18. [2] (2.3.2) Find a Schur triangularization of the matrix $\begin{pmatrix} 1 & 0 \\ i & 1 \end{pmatrix}$.

Solution 18. Let $A = \begin{pmatrix} 1 & 0 \\ i & 1 \end{pmatrix}$. First, we will construct the unitary matrix of the decomposition. The first column of such matrix can be found by finding the eigen vector of some eigen value of A. Since both diagonal entries of A is 1, A only has 1 eigen value = 1. Therefore:

$$\begin{pmatrix} 1-1 & 0 \\ i & 1-1 \end{pmatrix} \cdot u_1 = \begin{pmatrix} 0 & 0 \\ i & 0 \end{pmatrix} \cdot u_1 = 0$$

Trivially, we can see that u_1 is the column vector $[0 \ 1]^T$. Since A is a 2x2, we need to choose another vector that's orthorgonal to u_1 . Let this vector be $u_2 = [1 \ 0]^T$. Hence, we have:

$$A = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \cdot T \cdot \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}^* = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \cdot T \cdot \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$$

Solving for T, we obtain:

$$T = \begin{pmatrix} 1 & i \\ 0 & 1 \end{pmatrix}$$

Exercise 19. [3] (2.3.3) Find a Schur triangularization of the matrix $\begin{pmatrix} 1 & 1 & 1 \\ 1 & 1 & 1 \\ 1 & 1 & 1 \end{pmatrix}$.

Solution 19. Let $A = \begin{pmatrix} 1 & 1 & 1 \\ 1 & 1 & 1 \\ 1 & 1 & 1 \end{pmatrix}$. Similar to the previous exercise, we first find an eigen value, eigen vector pair of A to construct our unitary matrix U . From A , we see that the 3 eigen values are $\lambda_1 = 3, \lambda_2 = 0, \lambda_3 = 0$, and the 3 eigen vectors are: $v_1 = [1, 1, 1]^T, v_2 = [-1, 0, 1]^T, v_3 = [-1, 1, 0]^T$.

Since the 3 eigen vectors form a basis in \mathbb{R}^3 , we can use all 3 of them in our unitary matrix.

Take $u_1 = \frac{1}{\sqrt{3}}v_1$ as our first column of U and $u_2 = [0, -\frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2}}]^T$ as the second column. Then, we can construct our third column u_3 such that it is orthogonormal to both u_1, u_2 . Let $u_3 = [\sqrt{\frac{2}{3}}, \frac{1}{\sqrt{6}}, \frac{1}{\sqrt{6}}]^T$

Hence,

$$U = \begin{pmatrix} \frac{1}{\sqrt{3}} & 0 & \sqrt{\frac{2}{3}} \\ \frac{1}{\sqrt{3}} & -\frac{1}{\sqrt{2}} & \frac{1}{\sqrt{6}} \\ \frac{1}{\sqrt{3}} & \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{6}} \end{pmatrix}$$

Then we just solve for T with $T = U^*AU$:

$$T = \begin{pmatrix} 3 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}$$

Exercise 20. [2] (2.5.1) Find two normal matrices $A, B \in \mathbb{C}^{2 \times 2}$ such that neither $A + B$ nor AB is normal.

Solution 20. Let $A = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$ and $B = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$.

$$A + B = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} + \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} = \begin{pmatrix} 0 & 2 \\ 0 & 0 \end{pmatrix}$$

Since:

$$\begin{pmatrix} 0 & 2 \\ 0 & 0 \end{pmatrix} \begin{pmatrix} 0 & 0 \\ 2 & 0 \end{pmatrix} = \begin{pmatrix} 4 & 0 \\ 0 & 0 \end{pmatrix}$$

but

$$\begin{pmatrix} 0 & 0 \\ 2 & 0 \end{pmatrix} \begin{pmatrix} 0 & 2 \\ 0 & 0 \end{pmatrix} = \begin{pmatrix} 0 & 0 \\ 0 & 4 \end{pmatrix}$$

So $A + B$ is not normal. Similarly,

$$AB = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$$

We have:

$$(AB)(AB)^* = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix} = \begin{pmatrix} -1 & 0 \\ 0 & 1 \end{pmatrix}$$

and

$$(AB)^*(AB) = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$$

Because $(AB)(AB)^* \neq (AB)^*(AB)$, AB is not normal.

Exercise 21. [3] (2.5.2) Prove Proposition 1.0.6.

Proposition 1.0.6. Let $A \in \mathbb{C}^{n \times n}$ be a normal matrix. Let $p(x)$ be a polynomial in a single indeterminate x with coefficients in \mathbb{C} . Then, the matrix $p(A)$ is normal.

Solution 21. Let $p(A) = a_0A^0 + a_1A^1 + \cdots + a_dA^d \in \mathbb{F}^{n \times n}$

Then:

$$\begin{aligned} p(A)^* &= (a_0A^0 + a_1A^1 + \cdots + a_dA^d)^* \\ &= \bar{a}_0(A^0)^* + \bar{a}_1(A^1)^* + \cdots + \bar{a}_d(A^d)^* \end{aligned}$$

Since $(Z^n)^* = (Z^*)^n$:

$$\begin{aligned} p(A)^* &= \bar{a}_0(A^*)^0 + \bar{a}_1(A^*)^1 + \cdots + \bar{a}_d(A^*)^d \\ p(A)^* &= p(A^*) \end{aligned}$$

Since A is normal, A and A^* commutes. Hence, $p(A)$ and $p(A^*)$ commutes, which means that $p(A)$ is normal.

Exercise 22. [2] (2.5.3) Generalizing Proposition ?? (b), we might claim the following:

Let $A \in \mathbb{C}^{k \times k}$ be a normal matrix. Let $U \in \mathbb{C}^{n \times k}$ be an isometry. Then, the matrix UAU^* is normal.

Is this generalization correct?

Solution 22. From the proposition, since U^*U no longer reduce to I_k , we now have:

$$(UAU^*)(UAU^*)^* = UAU^*UA^*U^*$$

and,

$$(UAU^*)^*(UAU^*) = UA^*U^*UAU^*$$

Therefore, $(UAU^*)(UAU^*)^*$ only equal to $(UAU^*)^*(UAU^*)$ when we can commute U^*U with either A or A^*

Hence, this generalization is not correct.

Exercise 23. [4] (2.5.4) Let $A \in \mathbb{C}^{n \times n}$ be a normal matrix. Prove the following:

(a) We have $\|Ax\| = \|A^*x\|$ for each $x \in \mathbb{C}^n$.

(b) We have $\text{Ker } A = \text{Ker } (A^*)$.

(c) Let $\lambda \in \mathbb{C}$. Then, the λ -eigenvectors of A are the $\bar{\lambda}$ -eigenvectors of A^* .

Solution 23. (a) From the definition of vector norm: $\|z\|^2 = \langle z, z \rangle$, we have:

$$\|Ax\|^2 = \langle Ax, Ax \rangle = (Ax)^* Ax = \bar{x} A^* Ax$$

Since A is normal, A^* is normal and $A^*A = AA^*$

$$\|Ax\|^2 = \bar{x} AA^* x = (A^*x)^* A^* x = \langle A^*x \rangle$$

Hence,

$$\|Ax\|^2 = \|A^*x\|^2$$

Or,

$$\|Ax\| = \|A^*x\|$$

(b) Let x be an $n \times 1$ column vector in $\text{Ker } A$. Then we know that:

$$Ax = 0$$

Therefore,

$$\langle Ax, Ax \rangle = 0$$

Using the adjoint linear transformation:

$$\langle A^*Ax, x \rangle = 0$$

Since A is normal, $A^*A = AA^*$:

$$\langle AA^*x, x \rangle = \langle x, AA^*x \rangle = 0$$

$$\langle A^*x, A^*x \rangle = 0$$

Hence,

$$A^*x = 0$$

This means that x is also in the Kernel space of A^*

(c) Similar to the proof in the previous part (b), let λ be an eigen value of A and x be is corresponding eigen vector:

$$Ax = \lambda x$$

$$\begin{aligned}
xA^*Ax &= xA^*\lambda x \\
(xA)(A^*x) &= \lambda xA^*x \\
(A^*x^*)^*(\bar{\lambda}x) &= \lambda xA^*x
\end{aligned}$$

Conjugate transpose both sides:

$$\lambda x^*(A^*x^*) = \bar{\lambda}(x^*A)x^*$$

Shuffle around:

$$\begin{aligned}
A^*x^* &= \frac{\bar{\lambda}}{\lambda}(x^*A) \\
A^*x^* &= \frac{\bar{\lambda}}{\lambda}(A^*x)^* \\
A^*x^* &= \frac{\bar{\lambda}}{\lambda}(\bar{\lambda}x)^* \\
A^*x^* &= \frac{\bar{\lambda}}{\lambda}\lambda x^* \\
A^*x^* &= \bar{\lambda}x^*
\end{aligned}$$

Therefore, if A is normal and x is A 's eigen vector, then x^* is A^* 's eigen vector.

Exercise 24. [4] (2.5.5) (a) Let $A \in \mathbb{C}^{n \times n}$ and $B \in \mathbb{C}^{m \times m}$ be two normal matrices, and $X \in \mathbb{C}^{n \times m}$. Prove that $AX = XB$ if and only if $A^*X = XB^*$. (This is known as the (finite) *Fuglede–Putnam theorem*.)

(b) Let $A \in \mathbb{C}^{n \times n}$ and $X \in \mathbb{C}^{n \times n}$ be two matrices such that A is normal. Prove that X commutes with A if and only if X commutes with A^* .

[Hint: For part (a), set $C := AX - XB$ and $D := A^*X - XB^*$. Use Exercise ?? to show that $\text{Tr}(C^*C) = \text{Tr}(D^*D)$. Conclude using Exercise ?? (b). Finally, observe that part (b) is a particular case of part (a).]

Solution 24. (a) Let $C = AX - XB$ and $D = A^*X - XB^*$. We have:

$$\begin{aligned}
C^*C &= (AX - XB)^*(AX - XB) = (X^*A^* - B^*X^*)(AX - XB) \\
&= X^*A^*AX - X^*A^*XB - B^*X^*AX + B^*X^*XB
\end{aligned}$$

and,

$$\begin{aligned}
D^*D &= (A^*X - XB^*)^*(A^*X - XB^*) = (X^*A - BX^*)(A^*X - XB^*) \\
&= X^*AA^*X - X^*AXB^* - BX^*A^*X + BX^*XB^*
\end{aligned}$$

Since $\text{Tr}(AB) = \text{Tr}(BA)$ for any two matrix A and B in field F ($A \in F^{n \times m}$ and $B \in F^{m \times n}$), we can see that:

$$\begin{cases} \text{Tr}(X^* A^* A X) = \text{Tr}(X^* A A^* X) \\ \text{Tr}(X^* A^* X B) = \text{Tr}(X^* A X B^*) \\ \text{Tr}(B^* X^* A X) = \text{Tr}(B X^* A^* X) \\ \text{Tr}(B^* X^* X B) = \text{Tr}(B X^* X B^*) \end{cases}$$

Hence,

$$\begin{aligned} \text{Tr}(C^* C) &= \text{Tr}(X^* A^* A X) + \text{Tr}(X^* A^* X B) + \text{Tr}(B^* X^* A X) + \text{Tr}(B^* X^* X B) \\ &= \text{Tr}(X^* A A^* X) + \text{Tr}(X^* A X B^*) + \text{Tr}(B X^* A^* X) + \text{Tr}(B X^* X B^*) \\ &= \text{Tr}(D^* D) \end{aligned}$$

From this fact, we can clearly see that:

1. (The forward direction) If $AX = XB$, then $AX - XB = C = 0$. This means $\text{Tr}(C^* C) = 0$. Hence, $\text{Tr}(D^* D) = 0$ and $D = 0$, or $A^* X - X B^* = 0$, or $A^* X = X B^*$.
2. (The backward direction) Similary, If $A^* X = X B^*$ then $\text{Tr}(D^* D) = \text{Tr}(C^* C) = 0$. This means $C = AX - XB = 0$, or $AX = XB$.

(b) Using the result from part (a), we can see that part (b) is a special case of part (a), in which $B = A$. We also know that A^* is normal because A is normal. Hence, replacing B with A , we get:

$AX = XA$ if and only if $A^* X = X A^* = XA$. Therefore, X commutes with both A and A^* .

Exercise 25. [5] Let $A \in \mathbb{C}^{n \times n}$ and $B \in \mathbb{C}^{n \times n}$ be two normal matrices such that $AB = BA$. Prove that the matrices $A + B$ and AB are normal.

[Hint: Use Exercise 24 (b).]

Solution 25. Using the results from the previous exercise 24, we get these equalities:

$$\begin{cases} AB = BA & (1) \\ AB^* = B^* A & (A = A, X = B^*, B = A) & (2) \\ A^* B^* = B^* A^* & (A = A, X = B^*, B = A) & (3) \\ A^* B = B A^* & (A = A^*, X = B, B = A^*) & (4) \end{cases}$$

From these equalities, we have:

$$(AB)(AB)^* = ABB^* A^* \xrightarrow{3} ABA^* B^* \xrightarrow{1} BAA^* B^* = BA^* AB^* \xrightarrow{2} BA^* B^* A \xrightarrow{4}$$

$$A^*BB^*A = A^*B^*BA = (AB)^*(AB)$$

Hence, AB is normal.

Now, we need to prove that $A + B$ is also normal. We have:

$$(A + B)(A + B)^* = (A + B)(A^* + B^*) = AA^* + AB^* + BA^* + BB^*$$

and,

$$(A + B)^*(A + B) = (A^* + B^*)(A + B) = A^*A + A^*B + B^*A + B^*B$$

Since:

$$\begin{cases} AA^* = A^*A \\ AB^* = B^*A \quad (2) \\ BA^* = A^*B \quad (4) \\ BB^* = B^*B \end{cases}$$

,

$$(A + B)(A + B)^* = (A + B)^*(A + B)$$

and $A + B$ is normal.

Exercise 26. 4 (2.5.7) Let $A \in \mathbb{C}^{n \times n}$.

(a) Show that there is a **unique** pair (R, C) of Hermitian matrices R and C such that $A = R + iC$.

(b) Consider this pair (R, C) . Show that A is normal if and only if R and C commute (that is, $RC = CR$).

[Hint: For part (a), apply the “conjugate transpose” operation to $A = R + iC$ to obtain $A^* = R - iC$.]

Solution 26. (a)

Since $A = R + iC$, we have $A^* = R - iC$. Hence, let $B = \frac{A+A^*}{2}$ and $C = \frac{A-A^*}{2i}$. Then:

$$B^* = \left(\frac{A+A^*}{2}\right)^* = \frac{A^*+A}{2} = B$$

Hence B is Hermitian. Also,

$$C^* = \left(\frac{A-A^*}{2i}\right)^* = \frac{A^*-A}{-2i} = \frac{A-A^*}{2i} = C$$

Hence, C is Hermitian.

However,

$$B + iC = \frac{A+A^*}{2} + i\frac{A-A^*}{2i} = \frac{2A}{2} = A$$

Since both B and C are constructed from A , hence, they are an unique pair of hermitian matrices. QED.

(b)

$$A^*A = (R - iC)(R + iC) = R^2 + C^2 + iRC - iCR$$

and,

$$AA^* = (R + iC)(R - iC) = R^2 + C^2 - iRC + iCR$$

Hence, A^*A can only be equal to AA^* if and only if $iRC = iCR$, or $RC = CR$.

Exercise 27. [3] (2.5.8) (a) Let $T \in \mathbb{C}^{n \times n}$ be an upper-triangular matrix. Prove that

$$\sum_{i=1}^m (TT^* - T^*T)_{i,i} = \sum_{i=1}^m \sum_{j=m+1}^n |T_{i,j}|^2$$

for each $m \in \{0, 1, \dots, n\}$.

(b) Use this to give a direct proof (i.e., not a proof by contradiction) of Lemma 1.0.7.

Lemma 1.0.7. Let $T \in \mathbb{C}^{n \times n}$ be a triangular matrix. Then, T is normal if and only if T is diagonal.

Solution 27. (a) Let examine the entries of T (empty entries are 0):

$$T = \begin{pmatrix} a_{1,1} & a_{1,2} & \cdots & a_{1,n} \\ & a_{2,2} & \cdots & a_{2,n} \\ & & \ddots & \vdots \\ & & & a_{n,n} \end{pmatrix}$$

Hence, the entries of T^* is:

$$T^* = \begin{pmatrix} \bar{a}_{1,1} & & & \\ \bar{a}_{1,2} & \bar{a}_{2,2} & & \\ \vdots & \vdots & \ddots & \\ \bar{a}_{1,n} & \bar{a}_{2,n} & \cdots & \bar{a}_{n,n} \end{pmatrix}$$

We can see that the diagonal entries of the product of TT^* are simply:

$$(TT^*)_{i,i} = \sum_{k=i}^n a_{i,k} \bar{a}_{i,k} = \sum_{k=i}^n |a_{i,k}|^2$$

and those of the product of T^*T are:

$$(T^*T)_{i,i} = \bar{a}_{i,i} a_{i,i} = |a_{i,i}|^2$$

Therefore,

$$(TT^* - T^*T)_{i,i} = \sum_{k=i}^n |a_{i,k}|^2 - |a_{i,i}|^2 = \sum_{k=i+1}^n |a_{i,k}|^2$$

and finally,

$$\sum_{i=1}^m (TT^* - T^*T)_{i,i} = \sum_{i=1}^m \sum_{k=i+1}^n |a_{i,k}|^2$$

(b)

Direction 1: If T is normal $\rightarrow T$ is diagonal.

Since T is normal, T^* is also normal and $TT^* - T^*T = 0$. Hence,

$$\sum_{i=1}^m (TT^* - T^*T)_{i,i} = 0$$

$$\sum_{i=1}^m \sum_{k=i+1}^n |a_{i,k}|^2 = 0$$

This means that all elements $a_{i,k}$ where $k > i$ are 0. Also, because T is upper-triangular, $a_{i,k}$ where $k < i$ are also 0. Hence, T is diagonal.

Direction 2: If T is diagonal $\rightarrow T$ is normal.

This is obvious since T^* does not change the order of the diagonal elements.

QED.

Exercise 28. [2] (2.5.9) Let $A \in \mathbb{C}^{n \times n}$ be a normal matrix that is nilpotent. Prove that $A = 0$.

Solution 28. Let k be a positive integer such that $A^k = 0$. Let λ be an eigenvalue of A and x its corresponding eigenvector. Then we have:

$$A^k x = \lambda^k x$$

But since $A^k = 0$,

$$0 = \lambda^k x$$

and x is a non-zero vector, $\lambda = 0$.

Let consider the fact that A is also a normal matrix. So A is diagonalizable. Write A as: $A = UDU^*$ where U is unitary and D is a diagonal matrix with entries are the eigenvalues of A (spectral theorem for normal matrices). However, since $\lambda = 0$, $D = 0$.

$$A = UDU^* = U0U^* = 0$$

Exercise 29. [2] (2.6.1) Prove Proposition 1.0.8.

Proposition 1.0.8. Let $A \in \mathbb{C}^{n \times n}$. Let $U \in U_n(\mathbb{C})$ be a unitary matrix and $D \in \mathbb{C}^{n \times n}$ a diagonal matrix. Assume that for each $i \in [n]$, we have $AU_{\bullet,i} = D_{i,i}U_{\bullet,i}$ (that is, the i -th column of U is an eigenvector of A for the eigenvalue $D_{i,i}$). Then, $A = UDU^*$, so that (U, D) is a spectral decomposition of A .

Solution 29. Since the columns of U are the eigenvectors of A . Denote u_i to be the i -th eigenvector of A . Then $U = [u_1, u_2, \dots, u_n]$. Let $\lambda_1, \lambda_2, \dots, \lambda_n$ be A 's respective eigenvalues. We have:

$$AU = A[u_1, u_2, \dots, u_n] = [Au_1, Au_2, \dots, Au_n] = [\lambda_1 u_1, \lambda_2 u_2, \dots, \lambda_n u_n]$$

Construct a diagonal matrix D with entries $\lambda_1, \lambda_2, \dots, \lambda_n$, then:

$$AU = UD$$

$$AUU^* = UDU^*$$

Since U is unitary,

$$A = UDU^*$$

Exercise 30. 5 (2.6.2) (a) Find a spectral decomposition of the normal matrix $\begin{pmatrix} 1 & 1+i \\ 1+i & 1 \end{pmatrix}$.

(b) Find a spectral decomposition of the Hermitian matrix $\begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}$.

(c) Find a spectral decomposition of the skew-Hermitian matrix $\begin{pmatrix} 0 & i \\ i & 0 \end{pmatrix}$.

(d) Find a spectral decomposition of the unitary matrix $\frac{1}{\sqrt{2}} \begin{pmatrix} 1 & 1 \\ 1 & -1 \end{pmatrix}$.

Solution 30. (a) Let $A = \begin{pmatrix} 1 & 1+i \\ 1+i & 1 \end{pmatrix}$. Then, A 's eigenvalues are $\lambda_1 = 2 + i, \lambda_2 = -i$ and A 's eigenvectors are $x_1 = [1, 1]^T, x_2 = [-1, 1]^T$

Therefore, A 's spectral decomposition is:

$$A = \begin{pmatrix} 1/\sqrt{2} & -1/\sqrt{2} \\ 1/\sqrt{2} & 1/\sqrt{2} \end{pmatrix} \begin{pmatrix} 2+i & 0 \\ 0 & -i \end{pmatrix} \begin{pmatrix} 1/\sqrt{2} & 1/\sqrt{2} \\ -1/\sqrt{2} & 1/\sqrt{2} \end{pmatrix}$$

(b) Let $A = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}$. Then, A 's eigenvalues are $\lambda_1 = -1, \lambda_2 = 1$ and A 's eigenvectors are $x_1 = [i, 1]^T, x_2 = [-i, 1]^T$

Therefore, A 's spectral decomposition is:

$$A = \begin{pmatrix} i/\sqrt{2} & -i/\sqrt{2} \\ 1/\sqrt{2} & 1/\sqrt{2} \end{pmatrix} \begin{pmatrix} -1 & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} -i/\sqrt{2} & 1/\sqrt{2} \\ i/\sqrt{2} & 1/\sqrt{2} \end{pmatrix}$$

(c) Let $A = \begin{pmatrix} 0 & i \\ i & 0 \end{pmatrix}$. Then, A 's eigenvalues are $\lambda_1 = i, \lambda_2 = -i$ and A 's eigenvectors are $x_1 = [1, 1]^T, x_2 = [-1, 1]^T$

Therefore, A 's spectral decomposition is:

$$A = \begin{pmatrix} 1/\sqrt{2} & -1/\sqrt{2} \\ 1/\sqrt{2} & 1/\sqrt{2} \end{pmatrix} \begin{pmatrix} i & 0 \\ 0 & -i \end{pmatrix} \begin{pmatrix} 1/\sqrt{2} & 1/\sqrt{2} \\ -1/\sqrt{2} & 1/\sqrt{2} \end{pmatrix}$$

(d) Let $A = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & 1 \\ 1 & -1 \end{pmatrix}$. Then, A 's eigenvalues are $\lambda_1 = -1, \lambda_2 = 1$ and A 's eigenvectors are $x_1 = [1 - \sqrt{2}, 1]^T, x_2 = [1 + \sqrt{2}, 1]^T$

Therefore, A 's spectral decomposition is:

$$A = \begin{pmatrix} \frac{1-\sqrt{2}}{\sqrt{1+(\sqrt{2}-1)^2}} & \frac{1+\sqrt{2}}{\sqrt{1+(\sqrt{2}+1)^2}} \\ \frac{1}{\sqrt{1+(\sqrt{2}-1)^2}} & \frac{1}{\sqrt{1+(\sqrt{2}+1)^2}} \end{pmatrix} \begin{pmatrix} -1 & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} \frac{1-\sqrt{2}}{\sqrt{1+(\sqrt{2}-1)^2}} & \frac{1}{\sqrt{1+(\sqrt{2}-1)^2}} \\ \frac{1+\sqrt{2}}{\sqrt{1+(\sqrt{2}+1)^2}} & \frac{1}{\sqrt{1+(\sqrt{2}+1)^2}} \end{pmatrix}$$

Exercise 31. [2] (2.6.3) Describe all spectral decompositions of the $n \times n$ identity matrix I_n .

Solution 31. Let $I = UDU^*$. Since I is the identity matrix, all of its eigenvalues are 1s, so D is a diagonal matrix with its entries = 1. In other words, $D = I$. So,

$$I = UIU^*$$

Hence, we can see that UIU^* has to be equal to I . But this is true because U is unitary. Therefore the spectral decomposition of I is $I_n = UI_nU^*$ with U being any unitary matrix of size $n \times n$

Exercise 32. [3] Let $A \in \mathbb{C}^{n \times n}$ and $B \in \mathbb{C}^{n \times n}$ be two normal matrices such that $A \sim B$. Prove that $A \overset{us}{\sim} B$.

Solution 32. Let (Q, D) be the spectral decomposition of A and (P, F) be the spectral decomposition of B :

$$\begin{cases} A = QDQ^* & (1) \\ B = PFP^* & (2) \end{cases}$$

Also, since A is similar to B , we can write A as: $A = MBM^{-1}$ where M is an invertible matrix. However, since both A and B are square matrices, M must be square. This plus the fact that the columns of M are linearly independent, M is unitary. Therefore, $M^{-1} = M^*$ and $A = MBM^{-*}$. Using equation (2), we have:

$$MBM^* = MPFP^*M^*$$

$$MBM^* = (MP)F(P^*M^*) = (MP)F(MP)^*$$

In other words,

$$A = (MP)F(MP)^*$$

Because both M and P are unitary, (MP) is unitary. Therefore, A is also unitarily similar to F . Since $B \stackrel{\text{us}}{\sim} F$, $A \stackrel{\text{us}}{\sim} B$

Exercise 33. [3] (2.6.5) Prove Proposition 1.0.9 and Corollary 1.0.10.

Proposition 1.0.9. Let $A \in \mathbb{C}^{n \times n}$ be a skew-Hermitian matrix, and let (U, D) be a spectral decomposition of A . Then, the diagonal entries of D are purely imaginary.

Corollary 1.0.10. An $n \times n$ -matrix $A \in \mathbb{C}^{n \times n}$ is skew-Hermitian if and only if it is unitarily similar to a diagonal matrix with purely imaginary entries.

Solution 33. (a) Proof of proposition:

Let $A = UDU^*$, U is unitary, D is diagonal. We have:

$$A^* = UD^*U^*$$

But since A is skew-Hermitian, $A^* = -A$. Therefore,

$$UD^*U^* = -UDU^*$$

In other words,

$$D^* = -D$$

Since the entries of D^* are $\bar{\lambda}$, the only way for $\bar{\lambda} = -\lambda$ is for λ to be pure imaginary. QED.

(b) Proof of corollary:

\implies : Assume that A is skew-Hermitian. Then, A is normal. Hence, using the spectral theorem for normal matrices, $A = UDU^*$ for some unitary matrix $U \in U_n(\mathbb{C})$ and some diagonal matrix $D \in \mathbb{C}^{n \times n}$. Hence, using the proposition above, the entries of D must be pure imaginary. This proves the " \implies " direction of Corollary 1.0.10.

\impliedby : Assume that A is unitarily similar to a diagonal matrix with imaginary entries. In other words, $A = UDU^*$ for some unitary matrix $U \in U_n(\mathbb{C})$ and some diagonal matrix $D \in \mathbb{C}^{n \times n}$ that has imaginary entries. Therefore, it's easy to see that $D^* = -D$

Now, from $A = UDU^*$, we obtain

$$A^* = UD^*U^* = -UDU^* = -A.$$

In other words, the matrix A is skew-Hermitian. This proves the " \impliedby " direction of Corollary 1.0.10.

Exercise 34. [3] (2.7.1) Let \mathbb{F} be a field. Let n be a positive integer. Let $A \in \mathbb{F}^{n \times n}$ be an invertible matrix with entries in \mathbb{F} . Prove that there exists a polynomial f of degree $n - 1$ in the single indeterminate t over \mathbb{F} such that $A^{-1} = f(A)$.

Solution 34. Let A has the characteristic polynomial p_A : $p_A(t) = a_0t^0 + a_1t^1 + a_2t^2 + \cdots + a_{n-1}t^{n-1} + t^n$. Since A is invertible, $\det(A) \neq 0$. Hence, we have:

$$p_A(A) = a_0I + a_1A^1 + a_2A^2 + \cdots + a_{n-1}A^{n-1} + A^n = 0$$

$$a_0I + A(a_1 + a_2A^1 + \cdots + a_{n-1}A^{n-2} + A^{n-1}) = 0$$

$$A(a_1 + a_2A^1 + \cdots + a_{n-1}A^{n-2} + A^{n-1}) = -a_0I$$

$$-\frac{A}{a_0}(a_1 + a_2A^1 + \cdots + a_{n-1}A^{n-2} + A^{n-1}) = I$$

$$-\frac{A}{a_0A}(a_1 + a_2A^1 + \cdots + a_{n-1}A^{n-2} + A^{n-1}) = IA^{-1}$$

$$-a_0^{-1}(a_1 + a_2A^1 + \cdots + a_{n-1}A^{n-2} + A^{n-1}) = A^{-1}$$

Exercise 35. [3] (2.7.2) Let \mathbb{F} be a field. Let $A \in \mathbb{F}^{n \times n}$ be a square matrix with entries in \mathbb{F} . Prove that for any nonnegative integer k , the power A^k can be written as an \mathbb{F} -linear combination of the first n powers A^0, A^1, \dots, A^{n-1} .

Solution 35. Using the characteristic polynomial of $A \in \mathbb{F}^{n \times n}$ and Cayley-Hamilton theorem, we clearly see that if we want to find the value of the matrix A^k where $0 \leq k \leq n$, then:

$$a_0I + a_1A^1 + a_2A^2 + \cdots + a_kA^k + \cdots + a_{n-1}A^{n-1} + A^n = 0$$

$$a_0I + a_1A^1 + a_2A^2 + \cdots + a_{k-1}A^{k-1} + a_{k+1}A^{k+1} + \cdots + a_{n-1}A^{n-1} + A^n = -a_kA^k$$

$$A^k = -a_k^{-1}(a_0I + a_1A^1 + a_2A^2 + \cdots + a_{k-1}A^{k-1} + a_{k+1}A^{k+1} + \cdots + a_{n-1}A^{n-1})$$

Hence, A^k is a linear combination of the first n powers A^0, A^1, \dots, A^{n-1}

If $n < k$, then we have:

$$a_0I + a_1A^1 + a_2A^2 + \cdots + a_{n-1}A^{n-1} + A^n = 0$$

$$A^{k-n}(a_0I + a_1A^1 + a_2A^2 + \cdots + a_{n-1}A^{n-1} + A^n) = A^{k-n} \cdot 0 = 0$$

$$a_0A^{k-n} + a_1A^{k-n+1} + a_2A^{k-n+2} + \cdots + a_{n-1}A^{k-1} + A^k = 0$$

$$A^k = -A^{k-n}(a_0I + a_1A^1 + a_2A^2 + \cdots + a_{n-1}A^{n-1})$$

Then, A^{k-n} can be further broken down until $k - n < n$ using the same process. Hence, QED.

Exercise 36. [5] (2.7.3) Let a_1, a_2, \dots, a_k be k numbers. Let (x_0, x_1, x_2, \dots) be any (a_1, a_2, \dots, a_k) -recurrent sequence of numbers. Let d be a positive integer. Show that there exist k integers b_1, b_2, \dots, b_k such that each $i \geq kd$ satisfies

$$x_i = b_1 x_{i-d} + b_2 x_{i-2d} + \dots + b_k x_{i-kd}.$$

(This means that the sequences $(x_{0+u}, x_{d+u}, x_{2d+u}, x_{3d+u}, \dots)$ are (b_1, b_2, \dots, b_k) -recurrent for all $u \geq 0$.)

[Hint: For each $j \geq 0$, define the column vector v_j by $v_j = \begin{pmatrix} x_j \\ x_{j+1} \\ \vdots \\ x_{j+k-1} \end{pmatrix} \in \mathbb{R}^k$.

Let A be the $k \times k$ -matrix $\begin{pmatrix} 0 & 1 & 0 & \dots & 0 \\ 0 & 0 & 1 & \dots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \dots & 1 \\ a_k & a_{k-1} & a_{k-2} & \dots & a_1 \end{pmatrix} \in \mathbb{R}^{k \times k}$. Start by showing that $Av_j = v_{j+1}$ for each $j \geq 0$.]

Definition 1.0.11. Let a_1, a_2, \dots, a_k be k numbers. A sequence (x_0, x_1, x_2, \dots) of numbers is said to be (a_1, a_2, \dots, a_k) -recurrent if each integer $i \geq k$ satisfies

$$x_i = a_1 x_{i-1} + a_2 x_{i-2} + \dots + a_k x_{i-k}.$$

Solution 36. Let a column vector $v_j \in \mathbb{R}^k$ contains k entries of x from x_j to x_{j+k-1} .

For example, $v_0 = \begin{pmatrix} x_0 \\ x_1 \\ \vdots \\ x_{k-1} \end{pmatrix}$. Hence, v_{j+1} is $\begin{pmatrix} x_{j+1} \\ x_{j+2} \\ \vdots \\ x_{j+k} \end{pmatrix}$. But since we know from

the definition of x that: $x_{j+k} = a_1 x_{j+k-1} + a_2 x_{j+k-2} + \dots + a_k x_j$, we can construct a matrix A , such that the product $Av_j = v_{j+1}$. It's easy to see that:

$$A = \begin{pmatrix} 0 & 1 & 0 & \dots & 0 \\ 0 & 0 & 1 & \dots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \dots & 1 \\ a_k & a_{k-1} & a_{k-2} & \dots & a_1 \end{pmatrix}$$

. It's easy to see that $A^{-1} = A$ and since the entries a_1, a_2, \dots, a_k are real, $A^* = A$.

Therefore, A is Hermitian and normal. Also, it is important to note that:

$$v_1 = Av_0$$

$$v_2 = Av_1 = A(Av_0) = A^2v_0$$

$$v_3 = Av_2 = A(A^2v_0) = A^3v_0$$

Hence,

$$v_m = A^m v_0 \text{ for some } m \geq 0$$

In other words, to get v_m , we need to calculate A^m . For that, the characteristic polynomial of A is:

$$p_A(t) = \det(tI - A) = t^k + a_{k-1}t^{k-1} + a_{k-2}t^{k-2} + \cdots + a_1t + a_0t^0$$

Using the Cayley-Hamilton theorem and proof from the previous exercise 35, we know that A^m can always be evaluated as a linear combination of the first n powers A^0, A^1, \dots, A^{n-1} . Hence, we can always calculate v_m to get x_{m+1} .

Now, however, we must also prove that, for a positive integer d , if the sequence (x_0, x_1, x_2, \dots) satisfy:

$$x_i = a_1x_{i-1} + a_2x_{i-2} + \cdots + a_kx_{i-k} \text{ for all } i \geq k$$

then it must also satisfy,

$$x_i = b_1x_{i-d} + b_2x_{i-2d} + \cdots + b_kx_{i-kd} \text{ for all } i \geq kd$$

Again, let's consider A 's characteristic polynomial:

$$p_A(t) = t^k + a_{k-1}t^{k-1} + a_{k-2}t^{k-2} + \cdots + a_1t + a_0t^0$$

Let $b_i = -a_{k-i}$, we have:

$$p_A(t) = t^k - b_1t^{k-1} - b_2t^{k-2} - \cdots - b_{k-1}t - b_k$$

$$p_A(t) = t^k - \sum_{j=1}^k b_j t^{k-j}$$

Using the Cayley-Hamilton theorem, substitute t with A^d :

$$p_A(A^d) = (A^d)^k - \sum_{j=1}^k b_j (A^d)^{k-j} = A^{kd} - \sum_{j=1}^k b_j A^{(k-j)d} = 0$$

$$A^{kd} = \sum_{j=1}^k b_j A^{(k-j)d}$$

Let i be an integer such that $i \geq kd$. Multiplying both side of the equation above with $A^{i-kd}v_0$:

$$A^{kd}A^{i-kd}v_0 = \left(\sum_{j=1}^k b_j A^{(m-j)d}\right)A^{i-kd}v_0$$

$$A^i v_0 = \sum_{j=1}^k b_j A^{i-jd} v_0$$

But since we know that $A^{i-jd}v_0 = v_{i-jd} = \begin{pmatrix} x_{i-jd} \\ x_{i-jd+1} \\ \vdots \\ x_{i-jd+k-1} \end{pmatrix}$. Hence,

$$A^i v_0 = \sum_{j=1}^k b_j \begin{pmatrix} x_{i-jd} \\ x_{i-jd+1} \\ \vdots \\ x_{i-jd+k-1} \end{pmatrix} = \begin{pmatrix} \sum_{j=1}^k b_j x_{i-jd} \\ \sum_{j=1}^k b_j x_{i-jd+1} \\ \vdots \\ \sum_{j=1}^k b_j x_{i-jd+k-1} \end{pmatrix}$$

Now, the left hand side $A^i v_0 = v_i = \begin{pmatrix} x_i \\ x_{i+1} \\ \vdots \\ x_{i+k-1} \end{pmatrix}$. Therefore:

$$\begin{pmatrix} x_i \\ x_{i+1} \\ \vdots \\ x_{i+k-1} \end{pmatrix} = \begin{pmatrix} \sum_{j=1}^k b_j x_{i-jd} \\ \sum_{j=1}^k b_j x_{i-jd+1} \\ \vdots \\ \sum_{j=1}^k b_j x_{i-jd+k-1} \end{pmatrix}$$

We can see that from the first element of both sides' vectors:

$$x_i = \sum_{j=1}^k b_j x_{i-jd}$$

In other words,

$$x_i = b_1 x_{i-d} + b_2 x_{i-2d} + \cdots + b_k x_{i-kd}$$

Hence, QED.

Exercise 37. [2] (2.8.1) Let $A \in \mathbb{C}^{n \times m}$, $B \in \mathbb{C}^{n \times m}$ and $C \in \mathbb{C}^{n \times p}$ be three complex matrices. Prove that there exists a matrix $X \in \mathbb{C}^{m \times p}$ such that $AX - BX = C$ if and only if each column of C belongs to the image (= column space) of $A - B$.

Solution 37. Let $\text{Col}(M)$ denotes the column space of matrix M .

Since X is the left side of both A and B , we can write:

$$AX - BX = (A - B)X = C$$

WLOG, let v be a column vector in $\text{Col}((A - B)X)$. Then by definition, there's some vector x for which $v = ((A - B)X)x$. Set $y = Xx$, then $v = (A - B)y$. This means v is also a vector in $\text{Col}(A - B)$.

On the other hand, let v be a column vector in $\text{Col}(A - B)$. By definition, there's some vector y for which $v = (A - B)y$. Let x be some vector x in which $Xx = y$. Then we can write v as $v = (A - B)Xx$, which is equivalent to saying v is $\in \text{Col}((A - B)X)$.

From the above two proofs, $\text{Col}(A - B) = \text{Col}((A - B)X) = \text{Col}(C)$. Therefore, $A - B$ and C contain the same vectors \rightarrow columns of $C \in \text{Col}(A - B)$.

Exercise 38. [5] (2.8.2) Let A , B and C be as in Theorem 1.0.12.

(a) Let the linear map L be as in the above proof of the $\mathcal{V} \implies \mathcal{U}$ part of Theorem 1.0.12. Prove that if $\lambda \in \sigma(A)$ and $\mu \in \sigma(B)$, then $\lambda - \mu$ is an eigenvalue of L (that is, there exists a nonzero matrix $X \in \mathbb{C}^{n \times m}$ satisfying $L(X) = (\lambda - \mu)X$).

(b) Prove the implication $\mathcal{U} \implies \mathcal{V}$ in Theorem 1.0.12 (thus completing the proof of the theorem).

Theorem 1.0.12. Let $A \in \mathbb{C}^{n \times n}$ be an $n \times n$ -matrix, and let $B \in \mathbb{C}^{m \times m}$ be an $m \times m$ -matrix (both with complex entries). Let $C \in \mathbb{C}^{n \times m}$ be an $n \times m$ -matrix. Then, the following two statements are equivalent:

- \mathcal{U} : There is a **unique** matrix $X \in \mathbb{C}^{n \times m}$ such that $AX - XB = C$.
- \mathcal{V} : We have $\sigma(A) \cap \sigma(B) = \emptyset$.

Solution 38. (a) Consider the linear map

$$\begin{aligned} L : \mathbb{C}^{n \times m} &\rightarrow \mathbb{C}^{n \times m}, \\ X &\mapsto AX - XB. \end{aligned}$$

We can write L as a function of X such that: $L(X) = AX - XB$. Now, let λ be an eigenvalue of A and u be its corresponding eigenvector. Also, let μ be an eigenvalue of B and v be its corresponding eigenvector. Then, $\lambda \in \sigma(A)$ and $\mu \in \sigma(B)$. We have:

$$\begin{cases} Au = \lambda u \\ Bv = \mu v \end{cases}$$

Now, we denote a matrix $Z = uv^*$. Hence,

$$\begin{aligned} L(Z) &= AZ - ZB = Auv^* - uv^*B \\ &= \lambda uv^* - u(B^*v)^* \\ &= \lambda uv^* - u(\mu^*v)^* = \lambda uv^* - \mu uv^* = (\lambda - \mu)uv^* \\ &= (\lambda - \mu)Z \end{aligned}$$

Since $Z = uv^*$, Z must be nonzero. Hence if we let $X = Z$, then we have found a non-zero matrix X in which $L(X) = (\lambda - \mu)X$. Therefore, $\lambda - \mu$ must be an eigenvalue of L .

(b) Now, we are going to assume the contradiction (*): **assume both A and B share an eigenvalue δ** and we are going to show that if there exist a non-zero matrix $X \in \mathbb{C}^{n \times m}$ such that $AX - XB = C$, then we contradict our assumption.

Let $X = uv^*$ where:

$$\begin{cases} Au = \delta u \\ Bv = \delta v \end{cases}$$

We have:

$$L(X) = AX - XB = Auv^* - uv^*B = \delta uv^* - u(\delta^*v)^* = \delta uv^* - \delta uv^* = 0$$

Since $AX - XB = 0$, X must be 0. However, we defined X as $X = uv^*$, which is non-zero. Hence, X cannot exist (contradict our initial assumption(*)). Therefore, A and B cannot share any eigenvalue. QED.

Exercise 39. [3] (3.4.1) Compute the Jordan canonical form of the matrix

$$\begin{pmatrix} 1 & 0 & 2 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{pmatrix}.$$

Solution 39. Let $A = \begin{pmatrix} 1 & 0 & 2 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{pmatrix}$. Since A is already upper-triangular, A 's eigenvalues can be found on its diagonal. We can see that A only has 1 eigenvalue $\lambda = 1$. Because of that, A 's characteristic polynomial is $p_A(\lambda) = (1 - \lambda)^3$. Hence, the algebraic multiplicity of $\lambda = 1$ is 3. Therefore, the JCF of $A = J$ has

only 01 Jordan block of size 3.

Next, we determine the $\dim(\text{Ker}(A - \lambda I))$ to get the geometric multiplicity of λ , denoted as $\gamma(\lambda)$.

$$A - \lambda I = A - 1I = \begin{pmatrix} 0 & 0 & 2 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{pmatrix}$$

$$\text{Ker}(A - 1I) = \text{Ker} \begin{pmatrix} 0 & 0 & 2 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{pmatrix} = \text{Ker} \begin{pmatrix} 0 & 0 & 2 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}$$

Since $\begin{pmatrix} 0 & 0 & 2 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}$ has 1 pivot and 2 free variables, $\dim(\text{Ker}(A - 1I)) = 2 < 3 = \alpha(\lambda = 1)$.

Hence, we need to calculate $\text{Ker}((A - 1I)^2)$:

$$\text{Ker}((A - 1I)^2) = \text{Ker} \begin{pmatrix} 0 & 0 & 2 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{pmatrix}^2 = 0$$

Because $\text{Ker}((A - 1I)^2) = 0$, $\dim(\text{Ker}((A - 1I)^2)) = 3 = \alpha(\lambda = 1)$, then our Jordan block contains 2 smaller boxes of size 1×1 and size 2×2 :

$$J = \begin{pmatrix} J_1(1) & \\ & J_2(1) \end{pmatrix} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{pmatrix}$$

Now, we find the matrix S such that $AS = SJ$. We can write $S = [s_1, s_2, s_3]$ where s_1, s_2, s_3 are its columns. we have:

$$AS = [As_1, As_2, As_3] = SJ = [Se_1, Se_2, S(e_2 + e_3)] = [s_1, s_2, s_2 + s_3]$$

Therefore, we have:

$$\begin{cases} As_1 = s_1 \\ As_2 = s_2 \\ As_3 = s_2 + s_3 \end{cases}$$

Hence, s_1 and s_2 is in $\text{span}(\text{Ker}(A - 1I)) = \text{span} \left(\begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix}, \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix} \right)$

Let $s_1 = \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix}$. Let $s_2 = \alpha_1 \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix} + \alpha_2 \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}$ Hence,

$$As_2 = A \left(\alpha_1 \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix} + \alpha_2 \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix} \right) = \alpha_1 \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix} + \alpha_2 \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix} = \begin{pmatrix} \alpha_2 \\ \alpha_1 \\ 0 \end{pmatrix}$$

Since s_3 has to be orthogonormal to both s_1 and s_2 , $s_3 = \begin{pmatrix} 0 \\ 0 \\ \alpha_3 \end{pmatrix}$. Assume $\alpha_3 = 1$, then $s_3 = \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix}$. Hence, we have:

$$As_3 = A \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix} = \begin{pmatrix} 2 \\ 1 \\ 1 \end{pmatrix}$$

Then, s_2 is:

$$s_2 = As_3 - s_3 = \begin{pmatrix} 2 \\ 1 \\ 1 \end{pmatrix} - \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix} = \begin{pmatrix} 2 \\ 1 \\ 0 \end{pmatrix}$$

Therefore, S is:

$$S = \begin{pmatrix} 0 & 2 & 0 \\ 1 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

Hence,

$$A = \begin{pmatrix} 0 & 2 & 0 \\ 1 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} -1/2 & 1 & 0 \\ 1/2 & 0 & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

Exercise 40. [4] (3.4.2) Let $A \in \mathbb{C}^{n \times n}$ be a matrix. Prove that the following three statements are equivalent:

- \mathcal{A} : The matrix A is nilpotent.
- \mathcal{B} : We have $A^n = 0$.
- \mathcal{C} : The only eigenvalue of A is 0 (that is, we have $\sigma(A) = \{0\}$).

Solution 40. $\mathcal{A} \implies \mathcal{C}$:

Proof:

Let A be a nilpotent-matrix. Let λ be a non-zero eigenvalue of A and x its corresponding non-zero eigenvector. Then:

$$Ax = \lambda x$$

Now, let k be a positive integer in which $A^k = 0$. Then:

$$A^k x = \lambda^k x = 0$$

However, since x is non-zero, $\lambda^k = 0$, in other words, $\lambda = 0$. However, this contradicts our assumption. Hence, the only eigenvalue of A is 0.

$\mathcal{C} \implies \mathcal{B}$:

Because the only eigenvalue of A is 0, $p_A(t) = 0$. In addition, using the Cayley-Hamilton theorem, $p_A(A)$ is also 0. However, any polynomial p of degree n with 0 as the only root is of the type $p(x) = cx^n$. Hence, $A^n = 0$.

$\mathcal{C} \Leftarrow \mathcal{B}$:

Proof:

$\mathcal{A} \implies \mathcal{C}$ implies $\mathcal{B} \Leftarrow \mathcal{C}$ just by replacing k with n .

$\mathcal{A} \Leftarrow \mathcal{C}$:

Proof:

$\mathcal{C} \implies \mathcal{B}$ implies $\mathcal{A} \Leftarrow \mathcal{C}$ because if $A^n = 0$ then A is nilpotent.

$\mathcal{A} \iff \mathcal{B}$: This is obvious from the definition of nilpotent matrices.

Exercise 41. [6] (3.5.1) Let $\lambda \in \mathbb{C}$. Let n and k be two positive integers. Prove the following:

(a) If a $k \times k$ -matrix C has eigenvalue λ with algebraic multiplicity k and geometric multiplicity 1, then $C \sim J_k(\lambda)$.

(b) We have $(J_k(\lambda))^n \sim J_k(\lambda^n)$ if $\lambda \neq 0$.

(c) If $A \in \mathbb{C}^{k \times k}$ is an invertible matrix such that A^n is diagonalizable, then A is diagonalizable.

Solution 41. (a) Let A be a matrix $\in \mathbb{C}^{k \times k}$, which has eigenvalue λ with algebraic multiplicity k and geometric multiplicity 1. Therefore, A has a single eigenvalue and a single eigenvector. Hence, the JCF of A is exactly a $k \times k$ Jordan block of λ : $J_k(\lambda)$:

$$J_k(\lambda) = \begin{pmatrix} \lambda & * & \cdots & 0 & 0 \\ & \lambda & * & \cdots & 0 \\ & & \ddots & \ddots & \vdots \\ & & & \lambda & * \\ & & & & \lambda \end{pmatrix}$$

No matter how the structure of the sub-blocks inside $J_k(\lambda)$, it is by definition that A is similar to its JCF, $J_k(\lambda)$.

(b) Let $\lambda \neq 0$. Let a Jordan block be $J_k(\lambda) = \begin{pmatrix} \lambda & 1 & \cdots & 0 & 0 \\ & \lambda & 1 & \cdots & 0 \\ & & \ddots & \ddots & \vdots \\ & & & \lambda & 1 \\ & & & & \lambda \end{pmatrix}$. Hence,

$$J_k(\lambda^n) = \begin{pmatrix} \lambda^n & 1 & \cdots & 0 & 0 \\ & \lambda^n & 1 & \cdots & 0 \\ & & \ddots & \ddots & \vdots \\ & & & \lambda^n & 1 \\ & & & & \lambda^n \end{pmatrix}$$

Theorem 1.0.13. Let \mathbb{F} be a field. Let $k > 0$ and $\lambda \in \mathbb{F}$. Let $C = J_k(\lambda)$. Let $m \in \mathbb{N}$. Then, C^m is the upper-triangular $k \times k$ -matrix whose (i, j) -th entry is $\binom{m}{j-i} \lambda^{m-j+i}$ for all $i, j \in [k]$. (Here, we follow the convention that $\binom{m}{\ell} \lambda^{m-\ell} := 0$ when $\ell \notin \mathbb{N}$. Also, recall that $\binom{n}{\ell} = 0$ when $n \in \mathbb{N}$ and $\ell > n$.)

Using theorem 1.0.13, $J_k(\lambda)^n$ is:

$$J_k(\lambda)^n = \begin{pmatrix} \lambda^n & \binom{n}{1} \lambda^{n-1} & \binom{n}{2} \lambda^{n-2} & \dots & \dots & \dots & \dots \\ & \lambda^n & \binom{n}{1} \lambda^{n-1} & \binom{n}{2} \lambda^{n-2} & \dots & \dots & \dots \\ & & \lambda^n & \binom{n}{1} \lambda^{n-1} & \binom{n}{2} \lambda^{n-2} & \dots & \dots \\ & & & \lambda^n & \binom{n}{1} \lambda^{n-1} & \dots & \dots \\ & & & & \lambda^n & \dots & \vdots \\ & & & & & \ddots & \vdots \\ & & & & & & \lambda^n \end{pmatrix}$$

Now, since λ is the eigenvalue of A , then λ^n is the eigen value of A^n . Hence A^n is similar to $J_k(\lambda^n)$.

Moreover,

$$A = S J_k(\lambda) S^{-1}$$

$$A^n = S J_k(\lambda)^n S^{-1}$$

So A^n is also similar to its JCF - $J_k(\lambda)^n$. Therefore, $J_k(\lambda^n)$ is similar to $J_k(\lambda)^n$.

(c) From the previous part (b),

$$A^n = S J_k(\lambda)^n S^{-1}$$

Therefore, if A^n is diagonalizable, then $J_k(\lambda)^n$ is a diagonal matrix. Since $J_k(\lambda)^n$ has the form:

$$\begin{pmatrix} \lambda^n & \binom{n}{1} \lambda^{n-1} & \binom{n}{2} \lambda^{n-2} & \dots & \dots & \dots & \dots \\ & \lambda^n & \binom{n}{1} \lambda^{n-1} & \binom{n}{2} \lambda^{n-2} & \dots & \dots & \dots \\ & & \lambda^n & \binom{n}{1} \lambda^{n-1} & \binom{n}{2} \lambda^{n-2} & \dots & \dots \\ & & & \lambda^n & \binom{n}{1} \lambda^{n-1} & \dots & \dots \\ & & & & \lambda^n & \dots & \vdots \\ & & & & & \ddots & \vdots \\ & & & & & & \lambda^n \end{pmatrix}$$

But since $J_k(\lambda)^n$ is a diagonal matrix,

$$J_k(\lambda)^n = \begin{pmatrix} \lambda^n & & & & \\ & \lambda^n & & & \\ & & \lambda^n & & \\ & & & \lambda^n & \\ & & & & \lambda^n \\ & & & & & \ddots \\ & & & & & & \lambda^n \end{pmatrix}$$

Hence it's easy to see that:

$$J_k(\lambda) = \begin{pmatrix} \lambda & & & & \\ & \lambda & & & \\ & & \lambda & & \\ & & & \lambda & \\ & & & & \lambda \\ & & & & & \ddots \\ & & & & & & \lambda \end{pmatrix}$$

And because $A^n = SJ_k(\lambda)^nS^{-1}$, $A = SJ_k(\lambda)S^{-1}$, or:

$$A = S \begin{pmatrix} \lambda & & & & \\ & \lambda & & & \\ & & \lambda & & \\ & & & \lambda & \\ & & & & \lambda \\ & & & & & \ddots \\ & & & & & & \lambda \end{pmatrix} S^{-1}$$

Therefore, A is diagonalizable.

Exercise 42. 4 (3.5.2) Let \mathbb{F} be a field. Compute $(J_k(\lambda))^{-1}$ for any nonzero $\lambda \in \mathbb{F}$ and any $k > 0$.

Solution 42. Let $J_k(\lambda) = \begin{pmatrix} \lambda & 1 & \cdots & 0 & 0 \\ & \lambda & 1 & \cdots & 0 \\ & & \ddots & \ddots & \vdots \\ & & & \lambda & 1 \\ & & & & \lambda \end{pmatrix}$. We can see that $J_k(\lambda) =$

$\lambda I + B$, where B is:

$$\begin{pmatrix} 0 & 1 & \cdots & 0 & 0 \\ & 0 & 1 & \cdots & 0 \\ & & \ddots & \ddots & \vdots \\ & & & 0 & 1 \\ & & & & 0 \end{pmatrix}$$

Let $C = I - tB$. Hence, according to the Cayley-Hamilton theorem, $p_C(C)$ is:

$$p_C(C) = c_0I + c_1C^1 + c_2C^2 + \cdots + c_{k-1}C^{k-1} + C^k = 0$$

$$-c_0I = C(c_1 + c_2C^1 + \cdots + c_{k-1}C^{k-2} + C^{k-1})$$

$$C^{-1} = -\frac{1}{c_0}(c_1 + c_2C^1 + \cdots + c_{k-1}C^{k-2} + C^{k-1})$$

Hence,

$$(I - tB)^{-1} = I - tB + t^2B^2 - \cdots + (-t)^{k-1}B^{k-1}$$

Therefore, replacing t with λ we have:

$$J_k(\lambda)^{-1} = \frac{1}{\lambda}(I - \frac{1}{\lambda}B + \frac{1}{\lambda^2}B^2 - \cdots + \frac{1}{(-\lambda)^{n-1}}B^{k-1})$$