

Plant Protection Product Evaluation Report

Issuing authority: Fictional Crop Science Authority (FCSA) · Report ID: FCSA-PPR-2026-0042 · Issued on: 2026-04-18 (v1.0)

Overall summary. This report covers 2 products; 1 of them under conditional approval. Average efficacy across all applications: 86.3%.

AgriShield 250 EC

HIGH RISK NOTICE
This product carries a composite risk score of 78 out of 100 and requires elevated regulatory scrutiny. Consult the Restrictions section before authorising any application.

Active substance: **Fictoxazole** (250 g/L) · Formulation: Emulsifiable concentrate (EC) · Manufacturer: Demo CropChem GmbH

Germany: Zugelassen gem. PflSchG — Status: conditional. Valid from 2025-01-01 to 2027-12-31.

This product is registered for 3 crops with an average efficacy of **83.7%**.

Applications

Crop	Target pest	Stage (BBCH)	Dose	Max/season	Efficacy	PHI (days)
Winter wheat	Septoria tritici	31-39	1 L/ha	2	88%	35
Winter barley	Ramularia collo-cygni	37-49	1.25 L/ha	2	84%	35
Winter oilseed rape	Phoma lingam	14-20	0.8 L/ha	1	79%	56


Toxicity Indicators

Ld50 Oral Rat Mg Per Kg	1450
Ld50 Dermal Rat Mg Per Kg	2000
Fish Lc50 96H Mg Per L	0.89
Bee Ld50 Oral Microg Per Bee	12.4
Earthworm Lc50 14D Mg Per Kg	145

Risk Components

Human Health	45
Ecotoxicology	82
Environmental Fate	78
Residue In Food	30

Efficacy by crop



Restrictions

- Do not apply within 10 m of surface water bodies.
- Use closed-cab application equipment.
- Maximum two applications per growing season.
- Field re-entry only after spray deposit has dried.

Environmental notes

- Persistent in soil under cool, dry conditions; field DT50 = 92 days.
- Low risk of leaching to groundwater under typical European conditions.
- Moderate risk to non-target arthropods at edge-of-field.

Verdura 75 WG

Active substance: **Lumicarb-methyl** (750 g/kg) · Formulation: Water-dispersible granules (WG) · Manufacturer: Demo CropChem GmbH

Country: FR · Approval status: approved. Valid from 2024-06-01 to 2029-05-31.

This product is registered for 2 crops with an average efficacy of **89.0%**.

Applications

Crop	Target pest	Stage (BBCH)	Dose	Max/season	Efficacy	PHI (days)
Tomato (greenhouse)	<i>Botrytis cinerea</i>	61-89	0.6 kg/ha	3	91%	7
Grapevine	<i>Erysiphe necator</i> (powdery mildew)	53-81	0.4 kg/ha	4	87%	21


Toxicity Indicators

Ld50 Oral Rat Mg Per Kg	3200
Ld50 Dermal Rat Mg Per Kg	5000
Fish Lc50 96H Mg Per L	4.6
Bee Ld50 Oral Microg Per Bee	89
Earthworm Lc50 14D Mg Per Kg	480

Risk Components

Human Health	38
Ecotoxicology	51
Environmental Fate	36
Residue In Food	44

Efficacy by crop



Environmental notes

- Rapid degradation in soil; field DT50 = 8 days.
- Low bioaccumulation potential.
- Acceptable risk to bees under good agricultural practice.