

Quantum LTC With Positive Rate

David Ponnarovsky

August 11, 2022

preamble. preamble.

Claim for any $[[n, k, d]]$ CSS code property 1 holds

. **Proof.** let $y \in \{0, 1\}^n$ be a vector such $y \in G_z^\delta$, let assume that $|y|_{c^{\perp}} \leq C_2 d$ then for any $c \in C_x^\perp$:

$$\delta r_z \geq |H_z y| = |H_z (y + c)|$$

Robusstness Let $\omega \leq \Delta^2$. Let C_A and C_B be codes of length Δ with minimum distance d_A and d_B . We shall say that the dual tensor code $C = C_A \otimes \mathbb{F}_2^B + \mathbb{F}_2^A \otimes C_B$ is ω -robust, if for any codeword $c \in C$ of Hamming weight $|c| \leq \omega$, there exist $A' \subset A, B' \subset B, |A'| \leq |c|/d_B, |B'| \leq |c|/d_A$, such that $c_{ab} = 0$ whenever $a \notin A', b \notin B'$.

Definition. Sub-Tensor Pair We will say that C'_A, C'_B are sub-tensor pair of C_A, C_B if each of the code is subspace of C_A, C_B respectively and in addition one of the minimal codeword in C_A is also contained in C'_A (and similar to C'_B).

Note that the distance of each subcode is equal to the one from which it is derived. And also such code can be generated efficiently by choosing Δ non trivial coordinates of one of the minimal codewords and setting check nodes over them. (Assuming that Δ is even and that there is at least one different codeword in the code which has an overlap with that minimal codeword).

Claim. Subcode Robusstness. Consider the sub-tensor pair $C'_A \subset C_A, C'_B \subset C_B$, such that the dual tensor of C_A, C_B is ω -robust then the dual tensor of C'_A, C'_B is also ω -robust.

Proof. Let c be a codeword in the dual tensor of C'_A, C'_B then it's clear that c is also in the dual tensor of C_A, C_B and therefore there exists V, U subsets of A, B respectively such that c is supported only on them, and their size is less than $|c|/d_B, |c|/d_A$. As the length of the space of each of the subcodes is identical to their container, and by the fact that the distance of each of the subcodes is equal to the one which contains it, it follows that (1) $U \subset A' = A$ and (2) $|c|/d_A = |c|/d_{A'}$.

Existence Of Sub-Tensor Pair [COMMENT]

Try to prove existence by the probabilistic method.