

Hardness of Computing Fault Tolerance.

David Ponnarovsky

June 22, 2025


Introduction

- ▶ Brief overview of the topic
- ▶ Importance and relevance
- ▶ Objectives of the presentation

Key Points

- ▶ Main point 1
- ▶ Main point 2
- ▶ Main point 3

Nosiy Circuit.



Threshold Theorem.

Pippenger's Construction.

Encode each bit with the repetition code $0 \mapsto 0^m, 1 \mapsto 1^m$. Now observe that any logical operation, without decoding, can be made in $O(1)$ depth.

For example, $\text{OR}(\bar{x}, \bar{y})$ can be computed by applying in parallel $\text{OR}(x_i, y_i)$ for each i .

The 'Decoding' trick.

Instead of completely decoding, we would apply only a single step of partial decoding. We assume that in each code block the bits are partitioned into random disjoint triples, and we will apply a local correction to each of the triples by majority.

Claim

There are constants $\alpha, \eta \in (0, 1)$ such that for any bit string x at a distance $\leq \alpha n$ from the code (Repetition Code), one cycle of local correction on x yields x' such that:

$$d(x', C) \leq d(x, C)$$

The 'Decoding' trick.

Suppose that a bit absorb a bit flip with probability p . So in expectation we expect that entire block at length n will absorb pn flips.

$$\eta(\beta + p)n \leq \beta n$$
$$\beta \geq \frac{p}{1 - \eta}$$

From now on, we will assume that the graphs are bipartite and we will denote the right and the left vertices by V^- and V^+ . Notice that such expanders near Ramanujan exist, see for example [?]. The partition into two subsets enable us to come with a simple efficient decoder.

Expanders code are known for having good decoders, beneath, in ?? , we introduce a procedure to reduce an error. In overall, we alternately let to the right and then the left vertices to correct their own local view. In Theorem 1 we prove that when the applied error has size at most βn , for some constant β then the error's weight reduced by $\frac{1}{2}$. Repeating over the procedure $\Theta(\log(n))$ times completely correct the error.

We will call to the first stage, when only the right vertices suggest correction the right round, and to the second stage a left round.
For the whole procedure, we will call a single correction round.

Data: $x \in \mathbb{F}_2^n$

Result:

$$\arg \min \{y \in C : |y + x|\}$$

if $d(y, C) <$

1 **for** $v \in V^+$ **do**

2 $x'_v \leftarrow$

$$\arg \min \{y \in C_0 : |y + x|_v|\}$$

3 **end**

4 **for** $v \in V^-$ **do**


5 $x'_v \leftarrow$

$$\arg \min \{y \in C_0 : |y + x|_v|\}$$

6 **end**

7 **return** x

: Single decoding round



(b) location.

Lemma

If the error is at wight less than βn then a single round of the

Denote by $S^{(0)} \subset V^+$ and $T^{(0)} \subset V^-$ the subsets of left and right vertices adjacent to the error. And denote by $T^{(1)} \subset T^{(0)}$ the right vertices such any of them is connect by at least $\frac{1}{2}\delta_0\Delta$ edges to vertices at $S^{(0)}$. Note that that any vertex in $V^-/T^{(1)}$ has on his local view less than $\frac{1}{2}\delta_0\Delta$ faulty bits, So it corrects into his right local view in the first right correction round. Therefore after the right correction round the error is set only on $T^{(1)}$'s neighbourhood, namely at size at most $\Delta|T^{(1)}|$. We will show that this amount is strictly lower by a constant factor than $|e|$.

First, let's use the expansion property (??) for getting an upper bound on $T^{(1)}$ size:

$$\begin{aligned}\frac{1}{2}\delta_0\Delta|T^{(1)}| &\leq \Delta\frac{|T^{(1)}||S^{(0)}|}{n} + \lambda\sqrt{|T^{(1)}||S^{(0)}|} \\ \left(\frac{1}{2}\delta_0\Delta - \frac{|S^{(0)}|}{n}\Delta\right)|T^{(1)}| &\leq \lambda\sqrt{|T^{(1)}||S^{(0)}|} \\ |T^{(1)}| &\leq \left(\frac{1}{2}\delta_0\Delta - \frac{|S^{(0)}|}{n}\Delta\right)^{-2} \lambda^2|S^{(0)}|\end{aligned}$$

Since any left vertex adjoins to at most Δ faulty bits we have that $\Delta|S^{(0)}| \leq |e|$. Combing with the inequality above we get:

$$\Delta|T^{(1)}| \leq \left(\frac{1}{2}\delta_0\Delta - \frac{|e|}{n}\right)^{-2} \lambda^2|e|$$

Hence for $|e|/n \leq \beta = \frac{1}{2}\delta_0\Delta - \sqrt{2\lambda}$ it holds that $\Delta|T^{(1)}| \leq \frac{1}{2}|e|$.
Namely the error is reduced by half.

The Franch's Construction.




Figure: Caption for the image




Figure: Caption for the image




Figure: Caption for the image

Disjointness.