

Topological Memory.

David Ponarovsky

November 19, 2025

Introduction

Today:

- ▶ Proof.
 1. Walkthrough the math
 2. A
 3. B
- ▶ New experimental results.
- ▶ Answering the code teleportation.

Our Code

Code

In this session, we consider LDPC codes, where each bit adjoins to Δ_1 checks, and each check adjoins to Δ_2 bits. We denote by $\Delta = \Delta_1 \Delta_2$.

Local Decoder

definition

We say that D is a local decoder if when applied against $2p(1-p)$ -depolarized channel:

- ▶ There exists function $M : (0, 1) \rightarrow (0, 1)$ such that the probability of each bit being flipped is $M(2p(1-p)) < p^{\frac{100}{99}}$.
- ▶ There is a constant l such that for any bit, there are only $(\Delta)^l$ bits on which the event of it being flipped depends.

For example, if we take k large enough and consider the (n, k) -Toric, then both the swift rule-and the flip upon majority are local decoders.

Storing Logical State

Logical State

For a random variable X we denote by:

1. $\mathbf{E}_{\sim p}[X]$ - its expectation when measured after a single application of the p -depolarizing channel.
2. $\mathbf{E}_{\sim q^{(t)}}[X]$ - its expectation when measured after t rounds, each initiated by a p -depolarized channel followed by an application of local decoder D .

The Theorem

Theorem

There is $p_{\text{th}} \in (0, 1)$ such that for any $p < p_{\text{th}}$ and any bounded function f with a supremum $|f|$ it follows:

$$\mathbf{E}_{\sim q(t)}[f] \leq \mathbf{E}_{\sim p}[f] + t|f|e^{-\Theta(\sqrt{n})}$$

Lemma

There exists a constant $c > 0$, such:

$$\mathbf{E}_{\sim 2p(1-p)}[f \circ D] \leq \mathbf{E}_{\sim p}[f \circ D] + |f|e^{-c\sqrt{n}}$$


Theorem Proof

By induction on t .

$$\begin{aligned}\mathbf{E}_{\sim q^{(t)}}[f] &= \mathbf{E}_{\sim q^{(t-1)}}[\mathbf{E}_{\sim p}[f \circ D]] \\ &\leq \mathbf{E}_{\sim p}[\mathbf{E}_{\sim p}[f \circ D]] + (t-1)|\mathbf{E}_p[f \circ D]|e^{-c\sqrt{n}} \\ &\leq \mathbf{E}_{\sim 2p(1-p)}[f \circ D] + (t-1)|f|e^{-c\sqrt{n}} \\ &\leq \mathbf{E}_{\sim p}[f] + t|f|e^{-c\sqrt{n}}\end{aligned}$$

When in the last transition we used the lemma.

Dependence



$$\begin{aligned} & \Pr[(x, y) \in e | z \text{ is not flipped}] \\ &= \Pr[x \in e | z \text{ is not flipped}] \Pr[y \in e | z \text{ is not flipped}] \\ &\leq \Pr[x \in e] \Pr[y \in e] \end{aligned}$$

Dependence

Claim

Consider the application of a $2p(1-p)$ -depolarizing channel, followed by decoding D . For any subset of qubits S , the probability that there are more than $\frac{1}{100}$ dependent qubits in S is less than $e^{-c\sqrt{n}}$.

Proof

First, for an arbitrary S , denote by J the qubits that depend on qubits in S . Then, the expected number of dependent qubits in S is:

$$\begin{aligned} & \mathbf{E}_{\sim 2p(1-p)} [\text{dependent qubits in } S] \\ & \leq \mathbf{E}_{\sim 2p(1-p)} [|J|] \\ & = 2p(1-p)\Delta' |S| \rightarrow \frac{1}{100} |S| \end{aligned}$$

Dependence

We are going to use the following inequality:

Theorem 1 (McDiarmid's inequality). *Let $X_1, \dots, X_n$ be independent random variables, where X_i has range $\mathcal{X}_i$. Let $f: \mathcal{X}_1 \times \dots \times \mathcal{X}_n \rightarrow \mathbb{R}$ be any function with the $(c_1, \dots, c_n)$ -bounded differences property: for every $i = 1, \dots, n$ and every $(x_1, \dots, x_n), (x'_1, \dots, x'_n) \in \mathcal{X}_1 \times \dots \times \mathcal{X}_n$ that differ only in the i -th coordinate ($x_j = x'_j$ for all $j \neq i$), we have*

$$|f(x_1, \dots, x_n) - f(x'_1, \dots, x'_n)| \leq c_i.$$

For any $t > 0$,

$$\Pr(f(X_1, \dots, X_n) - \mathbb{E}[f(X_1, \dots, X_n)] \geq t) \leq \exp\left(-\frac{2t^2}{\sum_{i=1}^n c_i^2}\right).$$

Dependence

Thus, the probability that the number of dependent qubits in S is $|S|^{\frac{3}{4}}$ far from the expectation:

$$\Pr_{\sim 2p(1-p)} [\# - \mathbf{E} [\#]] \leq e^{-2 \frac{|S|^{\frac{2}{3}}}{n\Delta^I}}$$

So, for any $|S| > \alpha n$ we have:

$$\Pr_{\sim 2p(1-p)} [\# - \mathbf{E} [\#]] \leq e^{-\frac{2\alpha^{\frac{2}{3}}}{\Delta^I} \sqrt{n}}$$

Dependence

So, for any $|S| > \alpha n$, with probability greater than $1 - e^{-\frac{2\alpha^{\frac{2}{3}}}{\Delta'}\sqrt{n}}$, the number of dependent bits is less than:

$$2p(1-p)\Delta'|S| + |S|^{\frac{3}{4}} \rightarrow \frac{|S|}{100}$$

Lemma Proof


Denote by S the subset of the flipped qubits, and denote by $de(S)$ the number of dependent qubits in S .

$$\begin{aligned}\mathbf{E}_{\sim 2p(1-p)}[f \circ D] &= \sum f \circ D(S) \cdot \mathbf{Pr}[S] \\ &\leq \sum_{|S| \geq \alpha n} f \circ D(S) \cdot \mathbf{Pr}[S] + |f|e^{-c\sqrt{n}} \\ &\leq \sum_{|S| \geq \alpha n, de(S) < \frac{1}{100}|S|} f \circ D(S) \cdot \mathbf{Pr}[S] + |f|e^{-c\sqrt{n}} + |f|e^{-c\sqrt{n}} \\ &\leq \sum_{|S| \geq \alpha n, de(S) < \frac{1}{100}|S|} f(S)M(2p(1-p))^{\frac{99}{100}|S|} + 2|f|e^{-c\sqrt{n}}\end{aligned}$$


Lemma Proof

$$\begin{aligned} &\leq \sum_{|S| \geq \alpha n, de(S) < \frac{1}{100}|S|} f(S)p^{|S|} + 2|f|e^{-c\sqrt{n}} \\ &\leq \sum_S f(S)p^{|S|} + 2|f|e^{-c\sqrt{n}} \\ &\leq \mathbf{E}_{\sim p}[f(S)] + 2|f|e^{-c\sqrt{n}} \end{aligned}$$

v_2 is flipped.



v_2 is not faulty.



If the majority of bits in the local view of v_2 are flipped, then after decoding, v_2 will be also (remain) flipped.