

# SWEEP rule on the KD-Toric code

February 22, 2026

## 1 SWEEP Rule Decoder on the Toric

The decoder works as follows: each vertex  $v$  looks for a suggestion  $\hat{v}$  on  $\uparrow(v) \cap lk_2$  that matches  $\sigma_v^+$ . If there are more than one, then pick the one which has the minimal weight.




Figure 1: SWEEP rule

## 2 Span Expansion

Define the potentials functions  $L_i : \mathbb{R}^d \rightarrow \mathbb{R}$  as follow<sup>1</sup>:

$$L_i(\mathbf{x}) = \begin{cases} \mathbf{x}_i & i < d + 1 \\ -\langle \mathbf{1}, \mathbf{x} \rangle & i = d + 1 \end{cases}$$

The **Size** of a subset  $S$  is defined as:

$$\mathbf{Size}(S) = \sum_i \max_{\mathbf{x} \in S} L_i(\mathbf{x})$$

Numerically, we show that the span property holds for the checks. Namely, let  $e$  be an unsatisfied check at time  $t$ , and let  $H_e$  be the unsatisfied checks at time  $t - 1$ . Then there is a set  $H'_e \subset H_e$  such that:

---

<sup>1</sup>Notice that we use opposite signs relative to Berman and Peter Gacs, yet we agree with Perskil.

1. For any  $i < d + 1$  there exists  $e' \in H'_e$  such  $L_i(e') \geq L_i(e)$ .
2. There exists  $e' \in H'_e$  such that  $L_{d+1}(e') \leq L_{d+1}(e) + 1$ .

*Proof.* Consider a vertex  $v$  which at time  $t$  adjoins to an unsatisfied edge  $e$ , let  $u, w$  be vertices which have a positive faces that adjoin to  $e$ .

1. If  $\hat{u}|_e \neq \emptyset$ , then the parallel edge to  $e$  going out from  $u$  belongs to  $H_e$ . Denote it by  $e_u$ , taking  $e_u$  satisfies at least two (three in 4D) of the first  $d$  inequalities and the last  $d + 1$ th inequality. Denote by  $x \in [d]$  the index of the inequality which isn't satisfied by  $e_u$ . So it remains to prove the existence of  $e' \in H_e$  for which  $L_x(e') \geq L_x(e)$ .

Consider the assignment of bits at time  $t - 1$  over the complex, and consider the graph  $G = (\mathcal{F}, E)$  induced by associating vertices with any face that is set to 1 by the assignment. Two vertices are connected if their origin faces share an edge. Let  $K \subset \mathcal{F}$  be a connected component of faces that contains the faces supported on  $u$ .

We say that  $K$  is  $d$  far from  $u$  if:

$$\max_{f \in K} |f_x - u_x| \leq d$$

Now, we will prove by induction on  $d$  that there is a face  $f \in K$  with  $f_x \geq v_x$  such one of its edge is not satisfied.

- (a) **Base.** The edge from  $v$  which point at the  $\hat{z}$  direction must not be satisfied.




Figure 2: The induction base: In red, we have the faces on the support of  $u$ . The blue edge has the same  $x$ -coordinate as  $e$ . Any attempt to zero out the syndrome on the edge results in adding a face that is at least 2-far from  $u$ . Any configuration which is at most 1-far from  $u$  has to have that local view, yet might be connected to more faces which don't adjoin the blue edge.

- (b) **Assumption.** Assume that for any  $K$  which is at most  $d - 1$  far from  $u$  there is a face, with adjoins edge that is both has a  $x$ -coordinate that is greater than  $u_x$  and is not satisfied.

(c) **Induction Step.** Consider a connected component  $K$  that is  $d$ -far from  $u$ . Notice that if by substitute a co-boundary (stabilizer) from  $K$  we get  $K'$  that is at most  $d'$  far from  $u$  for some  $d' < d$  then we done. That because:  $\partial_2 K' = \partial_2(K + z) = \partial_2 K$  for some  $z \in \ker \partial_2$ .

On the other hand, [\[COMMENT\]](#) Finish that.

2. Else, if for any  $u \in \downarrow(u) \cap lk_1$  it holds that  $\hat{u}|_e = \emptyset$ , then it must be that  $e \in H_e$  because  $\hat{v}$  can only decrease the syndrome on  $v$ . Moreover, it follows that  $\hat{v} = \emptyset$  (otherwise, the suggestion would zero out the syndrome on  $\uparrow(v) \cap lk_2$ ). Thus,  $\sigma_v \not\subset \uparrow(v) \cap lk_2 \Rightarrow$  there is also an edge  $e' \in H_e$  where  $e' \in \downarrow(v)$ . In that case, take  $e \in H_e$  as the edge satisfies the first  $d$  equalities and  $e'$  for satisfying the  $d + 1$ th inequality.

□

### 3 Investigation

[COMMENT] To do: understands how the investigation should look like. Plot an example to such tree. The model goes as follow, at each even iteration we toss a noise, then in each odd iteration we perform a single SWEEP rule iteration. That allows to track over the history of the errors.

We say that a check  $e \in \text{Error}$ , if there is a face  $f$  that adjoin to  $e$  and belongs to the Error.

**Definition 3.1** (Investigation). For any unsatisfied check  $e$  at time  $t$ , we define  $V$  (the set of checks which participated in 'misleading' of  $e$ ), and two sets of edges on  $V$ , Arrows and Forks via the following algorithm:

```

1 let  $V = \{e\}$ , Arrows =  $\emptyset$ , Forks =  $\emptyset$ 
2 let  $Q \leftarrow \text{queue } \{e\}$ 
3 while  $Q$  is not empty do
4    $e' \leftarrow \text{dequeue } Q$ 
5   if  $e' \notin \text{Error}$  then
6      $e_1, \dots, e_l \leftarrow \text{Excuse}(e')$ 
7     Insert  $\{e_i, e'\}$  to Arrows
8     Insert  $\{e_i, e_j\}$  to Forks
9   end
10 end

```




Figure 3: blabla

## 4 Space-Time Graph

**Definition 4.1.** Consider a graph  $G = (V, A, F)$ , where  $V$  is a set of vertices, and  $A, F$  are sets of undirected edges. For  $v \in V$ , denote by  $\text{pre}(v) = \{u \in V : \{u, v\} \in A\}$ , we extend the notion for subsets of vertices  $X \subset V$  by  $\text{pre}(X) = \bigcup_{v \in X} \text{pre}(v)$ . Let  $T : V \rightarrow \mathbb{N}$ , We say that  $(G, T)$  is a time graph iff:

1.  $\{x, y\} \in A \Rightarrow T(y) = T(x) \pm 1$
2. For  $x, y \in V$   $\{x, y\} \in F$  iff there exists  $z \in V$  such both  $\{x, z\}, \{y, z\} \in A$ .

We will call to  $T$  time-function, to  $A$  arrows, and to  $F$  forks. Observe that from the second requirement it follows that forks can connect only vertices that have the same time value (set by  $T$ ).

**Definition 4.2** (History, slice.). For a time graph  $(G, T)$  and a vertex  $u \in V$ , let  $G_u$  be the subgraph of  $(V, A)$  induced by  $\{v \in V : T(v) \leq T(u)\}$ . We will denote by  $H_u$  the 'History of  $u$ ', which is the set of vertices that are connected to  $u$  in  $G_u$ . A slice is an equivalence class of the relation  $u \equiv v$  iff  $H_u = H_v$ . Notice that the time function has to be constant on a slice, to see this consider vertices  $x, y$  with  $T(x) > T(y)$ , thus  $x \notin G_y$  and therefore  $x \in H_x/H_y$ . So they are belong to different classes. Thus, we can extend the time function and the history to slices and write for a slice  $K$ :  $T(K)$  and  $H_K$ .

**Claim 4.1.** Consider slices  $K_1, K_2$  such that  $T(K_1) = T(K_2)$ , then either  $K_1 = K_2$  or  $H_{K_1}, H_{K_2}$  are disjoint.

*Proof.* Assume a non-empty intersection, so there exists  $z \in H_{K_1} \cap H_{K_2}$  such that for any  $x \in K_1$  and  $y \in K_2$ , there exist arrow-paths  $z \rightarrow x$  and  $z \rightarrow y$ . Thus, for any  $w \in H_{K_1}$ , one can concatenate a path  $w \rightarrow x \rightarrow z \rightarrow y$  and conclude that  $y \in H_{K_2}$ .  $\square$

**Definition 4.3** (The graph of slice). Let  $K$  be a slice. The graph of  $K$ , denoted by  $G_K = (V_K, E_K)$ , is the graph whose vertices are slices  $R \subset H_K$  with  $T(R) = T(K) - 1$ . Two vertices are connected by an edge if and only if there is a fork whose ends belong to the slices associated with them. Namely,  $S, R \in V_K$  are connected in  $G_K$  if and only if there is  $\{s, r\} = f \in F$  such that  $s \in S$  and  $r \in R$ .

**Claim 4.2.**  $G_K$  is connected.

*Proof.* Let  $R, S \in V_K$  and consider  $r, s \in V$  such that  $r \in R, s \in S$ . Recall that  $R, S \subset H_K$  and therefore  $r, s \in H_K$ . Thus, by definition of the 'History' there exists a path, passing through arrows only, from  $r$  to  $s$  in  $H_K$ . Consider such a path that is minimal in the number of points  $x$  belong to it, with  $T(x) = T(K)$ . By induction on the number  $k$  of such points we prove that  $R$  and  $S$  are connected in  $G_K$ .

1. Base.  $k = 0$ , In this case, the path from  $r$  to  $s$  lies within  $G_r (= G_s)$ . Hence  $H_r = H_s$ , So  $R = S$ .

2. Induction step. There exists  $y$  on the path from  $r$  to  $s$  such that  $T(y) = T(K)$ . Let  $x$  be the point which precedes  $y$  on the path, and  $z$  the point that follows  $y$ . Since the values of  $T$  at the two ends of an arrow differ by 1, and since  $y$  gets the maximal  $T$  value in  $H_K$  it follows that:

- (a)  $\{x, z\} \subset \text{pre}(y) \Rightarrow \{x, z\} \in F$ .
- (b) The slices of  $x$  and  $z$ , say  $K_x$  and  $K_z$  has time value  $T(K_x) = T(K_z) = T(K) - 1$ , Namely they are both members of  $V_K$ .

Since the path connecting  $r, x$  in  $H_K$  has  $k - 1$  vertices with  $T(\cdot) = T(K)$ , then by the induction hypothesis,  $R$  and  $K_x$  are connected in  $G_K$ , and by similar argument so are  $K_z$  and  $S$ . On the other hand,  $\{x, z\} \Rightarrow \{K_x, K_z\} \in E_K$ . Thus,  $R$  and  $S$  are connected in  $G_K$ .

□

**Definition 4.4.** Let  $f : G \hookrightarrow \mathbb{R}^d$  be an embedding that maps each element of  $V \cup A \cup F$  to a point in  $\mathbb{R}^d$ . We say that  $(G, \mathbb{R}^d)$  is a space-time graph iff:

- 1.
2. The point set by  $f$  for an edge is the mean of its endpoints' images; that is, for  $e = \{x, y\} \in A \cup F$ , we have  $f(e) = \frac{1}{2}(f(x) + f(y))$ .

**Definition 4.5.** Define  $\text{pre}_i(v) = \arg \max_{u \in \text{pre}(v)} L_i(f(u))$

**Claim 4.3.** Let  $\mathbf{S} = \{S_1, S_2, \dots, S_k\}$  where  $S_i \subset \mathbb{R}^d$ , introduce an edge between  $S_i$  and  $S_j$  iff  $S_i \cap S_j \neq \emptyset$ . Assume that  $\mathbf{S}$  is connected. Then  $\mathbf{Size}(\cup_{i=1}^k S_i) \leq \sum_{i=1}^k \mathbf{Size}(S_i)$ .

*Proof.* It suffices to prove that if  $R_1 \cap R_2 \neq \emptyset$ , then  $\mathbf{Size}(R_1 \cup R_2) \leq \mathbf{Size}(R_1) + \mathbf{Size}(R_2)$ . Let  $a_{i,j} = \max_{v \in R_j} L_i(v)$ . Then for any  $w \in R_1 \cap R_2$  we have:

$$\begin{aligned} \mathbf{Size}(R_1 \cap R_2) &= \sum_i^{d+1} \max a_{i,1}, a_{i,2} = \sum_i^{d+1} a_{i,1} + a_{i,2} - \min a_{i,1}, a_{i,2} \\ &\leq \sum_i^{d+1} a_{i,1} + a_{i,2} - L_i(w) = \mathbf{Size}(R_1) + \mathbf{Size}(R_2) - \mathbf{Size}(w) \\ &= \mathbf{Size}(R_1) + \mathbf{Size}(R_2) \end{aligned}$$

□

**Definition 4.6** (Spanned Set). Consider an embedding  $f : \star \hookrightarrow \mathbb{R}^d$ , a subset of  $f$ 's domain  $S \subset \star$ , and  $d + 1$  points  $x_1, \dots, x_{d+1}$ . We say that  $\langle S, x_1, \dots, x_{d+1} \rangle$  is a spanned set of  $S$  if for any  $i \in [d + 1]$  and  $s \in S$ , it holds that  $L(f(s)) \leq L_i(x_i)$  and in addition there are  $v_1, \dots, v_{d+1} \in V$ , not necessary different, that achieve the equalities  $L_i(x_i) = L_i(f(v_i))$ . We name  $x_1, \dots, x_{d+1}$  the poles of  $S$ . Furthermore, we extend the notation to a spanned slice when  $S$  is a slice of some time-graph. Notice that saying  $\langle S, x_1, \dots, x_{d+1} \rangle$  is a spanned set means that  $\mathbf{Size}(S) = \mathbf{Size}(\{x_1, \dots, x_{d+1}\})$ .

Observe that in the case where the subset  $S$  is a slice. Each of these poles can be associated with a vertex of  $G$  that gives the equality. If there are multiple vertices whose images by  $f$  give equality, associate the point with one of them arbitrarily. We will use the notion of applying functions originally defined over vertices to the poles, referring to the application over the vertices associated with them. For example  $\text{pre}(x_i) = \text{pre}(v)$  for the vertex  $v \in S$  which satisfies  $L_i(f(v)) = L_i(x_i)$  and therefore is associated with the pole  $x_i$ . Another example is referring to  $x_i$  as a vertex, so when saying 'For  $x_i \in V_k$ ' or 'For  $\{x_i, u\} \in F$ ', we refer to the associated vertex or the edge connecting it to  $u$ .

**Claim 4.4.** Consider a time graph  $(G, T)$ , an embedding  $f : G \rightarrow \mathbb{R}^d$ , such that.. , Let  $K$  be a slice in  $G$ , and let  $\hat{K} = \langle K, x_1, \dots, x_{d+1} \rangle$  be a spanned slice of  $K$  such that  $K \neq H_K$ . Then for some integer  $k$  there exist spanned slices  $\hat{K}_0 = \langle K_0, \cdot \rangle, \dots, \hat{K}_k = \langle K_k, \cdot \rangle$  and Forks  $F_1, \dots, F_k$  such that:

1. For  $i \in [d+1]$   $\text{pre}_i(x_i)$  belongs to the poles  $\hat{K}_0, \dots, \hat{K}_k$ .
2. Introduce an edge between  $\hat{K}_i$  and  $\hat{K}_j$  if and only if one of the Forks  $F_1, \dots, F_k$  consists of a pole of  $\hat{K}_i$  and a pole of  $\hat{K}_j$ . Then the graph formed from  $\hat{K}_0, \dots, \hat{K}_k$  is connected.
3.  $\sum_{i=0}^k \text{Size}(\hat{K}_i) + \sum_{i=1}^k \text{Size}(F_i) \geq \text{Size}(\hat{K}) + 1$

*Proof.* First Notice that  $K$  doesn't contain leaves in  $(V, A)$ , since a leaf is connected only to itself via arrows, and therefore its 'slice' equals to its 'History'. Thus  $\text{pre}_i(x_i)$  is not empty for  $i = 1, \dots, d+1$ . Next consider the graph  $G_K = (V_K, E_K)$  from definition 4.3. Since  $\{x_i, \text{pre}_i(x_i)\}$  is an arrow  $\text{pre}_i(x_i)$  belongs to a slice in  $V_K$ . for  $i = 1, \dots, d+1$  denote by  $R_i$  the slice in  $V_K$  that contains  $\text{pre}_i(x_i)$ .

By claim 4.2  $G_K$  is connected and therefore there exist a minimal collection  $\{K_0, \dots, K_k\}$  of slices from  $V_K$  which contains  $R_1, R_2, \dots, R_{d+1}$ , and which is connected in  $G_K$ . We pick a set of forks  $\mathbf{F} = \{F_1, \dots, F_k\}$  which realizes a spanning tree for this collection of slices. Later we will identify edges from this spanning tree with the forks from  $\mathbf{F}$ .

For  $i = 1, \dots, d+1$  we set  $\text{pre}_i(v_i)$  to be the  $i$ th pole of  $R_i$ , This assures (1). The set of remaining poles for  $\hat{K}_0, \dots, \hat{K}_k$  will be equal to  $\cup_{i=k}^k F_i$ . This will assure (ii), We will also select poles for  $F_1, \dots, F_k$  to form 'spanned forks'  $\hat{F}_1, \dots, \hat{F}_k$  in such way that:

$$\sum_{i=0}^k \text{Size}(\hat{K}_i) + \sum_{i=1}^k \text{Size}(\hat{F}_i) \geq \sum_{i=1}^{d+1} L_i(\text{pre}_i(v_i)) \geq \text{Size}(\hat{K}) + 1$$

This will assure (3).

Let us denote by  $J_0, \dots, J_{2k}$  the enumerated union  $K_0, \dots, K_k, F_1, \dots, F_k$ . Observes that for any choise of  $(d+1)(2k+1)$  elements  $z_0^1, z_0^2, z_0^3, \dots, z_0^{d+1}, \dots, z_{2k}^{d+1}$  in  $\mathbb{R}^d$ , such that  $z_j^i \in J_j$  it holds that:

$$\sum_{i=0}^k \text{Size}(\hat{K}_i) + \sum_{i=1}^k \text{Size}(\hat{F}_i) \geq \sum_{j,i} L_i(z_j^i)$$

We will choose the  $z_j^i$  by the following process. Pick a  $z_j^i$  that has not been set yet. If  $J_j = R_i$ , then set  $z_j^i \leftarrow \text{pre}_i(x_i)$ ; otherwise, consider the path from  $J_j$  to  $R_i$  (by connectivity it has to be such one, and by minimality there is exactly one). Let  $J_t$  be the successor on the path, then pick  $z_j^i \in J_j \cap J_t$  according to some deterministic rule dependent on  $J_j, J_t$ .

Observes that  $z_t^i$  will not be set to  $J_j \cap J_t$  since  $J_j$  can't be the successor of  $J_t$  on the path to  $R_i$ . Thus for a fixed  $i$  there is no more  $j'$  such  $z_{j'}^i = z_j^i$ .

Now consider a non trivial intersrion  $\bigcap_{j \in X} J_j$ , we claim that for any  $i$ , there is  $t \in X$  such that  $J_t$  is the succcessor of  $J_j$  on the path to  $R_i$  for any  $j \in X/\{t\}$ .

By induction, on the number of the poles:  $(d + 1)$ .

1. Base for  $d + 1 = 3$ , proven in the original work of Berman.
2. Induction Step. Consider  $d + 1$ , Consider the collection  $J'$  botained by erasion minimal number of spannsed sets (including spanned forks) such that the  $d + 1$  th pole is not connected any more. Form the intersection, we removed at most one.  $|X| = x$ ,

First, for any  $i$  there is  $t$ , and at least one  $j$ , such that  $J_t$  is the succcessor of  $J_j$ . For see that just consider a path from the  $i$ th pole that pass trough the junction  $\bigcap_{j \in X} J_j$ , since the graph is connected and covers all the poles, then there is must to be path from  $i$ th to any of the intrsected spanned-sets, denote them as  $P_1$  and  $P_2$ . In addition, since the graph, is minimal. if there is sach a path, then the two spanned-sets consist the

If not, namely there are  $j_1, j_2$  such that  $J_t$  is the succcessor to  $J_{j_1}$  but not the succcessor of  $J_{j_2}$  then one can close a cycle by concatinating  $J_t \rightarrow R_i \rightarrow J_{j_2} \rightarrow J_t$  which is a contradiction to the minimality. Thus,  $z_X$  will appers  $|X| - 1$  in  $\{z_j^i : j \in X\}$  for any  $i$ .

consider the following routine, we start in the  $i$ th pole, and then start to scan the graph in BFS manner. Then we associate a coloeer either 0 or 1 accoridng the following rule, until uncovering one onf the nodes form  $X$ , we assign 0, after the uncovering of one the vertices in the junction, we first colorize the first uncovered nodes by 0 and then we coloraize any of the other nodes do be meet with 1.

Now, since, the graph, is tree, the spanning tree reuturned by the BFS precedure is alawas the same. Thus, no matther the what is the local edges orination, the  $t \in X$  for which  $J_t$  is colored by 0 is always the same. Pick it to be  $J_t$ .

claim, suppose that  $\{u, v\} \in E$  and  $d(u, s) = d(v, s)$  then there is a cycle in the graph.

Consider the graph  $G' = (V', E')$ , such that  $V' = \{J_j\}_j \cup v$ . where to any  $x \in X$  we have  $\{J_x, v\} \in E'$  and for any  $x, y$  such  $y \notin X$ , and  $J_x \cap J_y \neq \emptyset$ , then  $\{J_x, J_y\} \in E'$ .  
[COMMENT] Should split any junction.

Since, the graph is tree,

□

## 5 Explantation of a Spanned Slice

[COMMENT] Here we follow the original proof, but change the triminalogy,  $m$  will be the number of leaves in the explanation, then in the end we conclude that the number of faults is at least  $m/\Delta$ . So  $|R| \leq f(\frac{m}{\Delta})$ .

**Definition 5.1.** Given sets  $W$  of points,  $R$  of arrows and  $F$  of forks, we define  $U$  as the set of vertices of the graph induced by the edges of  $R \cup F$ . The sets  $W, R$  and  $F$  form an explanation of a spanned slice  $K = \langle K, v_1, \dots, v_{d+1} \rangle$  iff:

1.  $W \cup \{v_1, \dots, v_{d+1}\} \subset U \subset H_K$  and the graph  $\langle U, R \cup F \rangle$  is connected.
2. All elements of  $W$  are errors.
3. For  $m = |W|$  and  $s = \mathbf{Span}(K)$  we have  $|R| \leq \alpha(m-1) - \beta s$  and  $|F| < m-1$ .

**Claim 5.1.** Every spanned slice  $K = \langle K, v_1, \dots, v_{d+1} \rangle$  has an explanation.

*Proof.* Let  $h = T(K) - \min_{u \in H_K} T(u)$ . The proof is by induction on  $h$ .

1. Base.  $h = 0$ . Then no arrows can connect points of  $H_K$ , Thus  $H_K$ , as well as  $\mathbf{K}$  consists of a single point  $u$  such that  $u \in \text{Error}$  [COMMENT] Leaves. In that case  $U = W = \{u\}$ ,  $m = 1, s = 0$  and  $|R|, |F| = 0$  satisfies the required.
2. Induction Step. Now, we know that  $K \neq H_K$ . Thus, by ??, we know that for some  $k$  there exist spanned slices  $K_0, \dots, K_k$  and forks  $F_1, \dots, F_k$  that satisfy ??.

Let  $s_i = \mathbf{Span}(K_i)$ . By the inductive hypothesis, we know that  $K_i$  has an explanation  $\mathbf{W}_i, \mathbf{R}_i, \mathbf{F}_i$ . For  $m_i = |\mathbf{W}_i|$ , the sets  $R_i$  and  $F_i$  have at most  $\alpha m_i - \beta s_i$  and  $m_i - 1$  leaves. We define:

- (a)  $\mathbf{W} = \cup_{i=0}^k \mathbf{W}_i$ , with  $m = \sum_{i=0}^k m_i$  leaves. Notice that  $\mathbf{W}_i \subset K_i$  and that  $T(K_i) = T(K_j)$  for all  $i, j$ , thus by ??  $\mathbf{W}$  is a union of disjoint set.
- (b)  $\mathbf{F} = \{F_1, \dots, F_k\} \cup_{i=0}^k \mathbf{F}_i$  with at most  $k + \sum_{i=0}^k m_i - 1 = m - 1$  elements.
- (c)  $\mathbf{R} = \{\{v_i, \mathbf{Excuse}_i(v_i)\}\} \cup_{i=0}^k \mathbf{R}_i$  with at most

$$d+1 + \sum_{i=0}^k \alpha(m_i - 1) - \beta s_i \leq d+1 + \alpha m - \alpha(k+1) - \beta(s - 1 + k \cdot \mathbf{Span}(\text{fork}))$$

$$= \alpha m - \beta s + (d+1 - (k+1)\alpha + \beta - \beta k \cdot \mathbf{Span}(\text{fork}))$$

So its enough to pick  $\alpha, \beta$  such that for any  $k \geq 1$ :

$$d+1 - (k+1)\alpha + \beta - \beta k f \leq 0$$

$$\Rightarrow \beta = 1, \alpha = (d+1)^3(1+f)$$

It is easy to see that  $\mathbf{W}, \mathbf{R}, \mathbf{F}$  form an explanation of  $K$ .

□

## 6 Counting Subtrees.

**Claim 6.1.** Let  $G = (V, E)$  be a  $\Delta + 1$ -regular graph, and let  $k$  be an integer less than  $|E|$ . Fix a vertex  $r \in V$ . The number of subtrees rooted at  $r$  is at most  $\alpha^k$ .

*Proof.* Denote by  $T_\Delta$  and  $T_2$  the infinity  $\Delta$  and binary tree. There is an injection between the subtrees of  $G$ , rooted at  $v$ , with at most  $k$  edges, to the subtrees of  $T_\Delta$  rooted at  $v$ , with at most  $k$  edges. Moreover, there is an injection from the subtrees of  $T_\Delta$  rooted at  $v$ , with at most  $k$  edges to the subtrees of  $T_2$ , rooted at  $v$ , with at most  $k \log \Delta$  edges.

**[COMMENT]** In the second map, we think on replacing each of the vertex with  $\Delta - 1$ -leaves binary tree, each leaf is associated with a outgoing edge. .

Thus, it's enough to bound the number of subtrees in  $T_2$  rooted at  $v$ , with at most  $k \log \Delta$  edges, that number is less than the number of binary tree with  $k \log \Delta + 1$  leaves. For exact leaves number we get the  $k \log \Delta$  Catalan number, which its limit is:

$$\sim \frac{4^{k \log \Delta}}{(k \log \Delta)^{\frac{3}{2}} \sqrt{\pi}} \leq \frac{\Delta^{2k}}{(k \log \Delta)^{\frac{3}{2}} \sqrt{\pi}}$$

Therefore we can bound by:

$$\sum_{j=1}^k \frac{\Delta^{2j}}{(j \log \Delta)^{\frac{3}{2}} \sqrt{\pi}} \leq \frac{\Delta^{2k}}{\sqrt{\pi} \log^{\frac{3}{2}} \Delta} \sum_{j=1}^k \frac{1}{j^{\frac{3}{2}}} \leq \zeta\left(\frac{3}{2}\right) \cdot \frac{\Delta^{2k}}{\sqrt{\pi} \log^{\frac{3}{2}} \Delta}$$

□

Now, from here we get an upper bound on the number of subgraphs that connected  $l$  vertices at the top level. We bound it from above by counting combination of rooted subtrees, such the vertices are their root and the overall edges number is summed up to  $k$ . Thus:

$$\leq \zeta\left(\frac{3}{2}\right) \binom{k-1}{l-1} \frac{\Delta^{2k}}{\sqrt{\pi} \log^{\frac{3}{2}} \Delta}$$

## 7 Tree Separator Lemma

## 8 Tanner - History Connected Equivalency

Imagine the following process, we apply  $t$  noisy iterations of the SWEEP rule, and then a single ideal (without noise) iteration. Denote by  $\xi$  a deviated configuration (can be either bits or checks), and by  $\xi'$  the result of applying an ideal SWEEP rule on  $\xi$ . Notice that if  $\xi$  is connected in the Tanner graph then  $\xi'$  is connected in the History graph [\[COMMENT\] Require proof, For the classical case enumerate all the options can be done by hand.](#) Thus:

$$\begin{aligned}
\Pr[\xi \text{ connected in Tanner}] &\leq \Pr[\xi' \text{ connected in Time} \mid \text{no noise at the } t+1 \text{ iteration}] \\
&\leq \sum_m^\infty |\{\text{explanation to } \xi' \text{ over } m \text{ leaves restricted to no noise at the final iteration}\}| p^{\frac{m}{\Delta}} \\
&\leq \sum_m^\infty |\{\text{explanation to } \xi' \text{ over } m \text{ leaves}\}| p^{\frac{m}{\Delta}} \\
&\leq \Pr[\xi' \text{ connected in Time} - \text{Upper Bound}] \leq q^{\text{Span}(\xi')} \leq q^{\text{Span}(\xi)-1}
\end{aligned}$$

## 9 Intermediate Step Lemma

[\[COMMENT\]](#) The idea of this step is also to bound the case that all the bits are flipped at once, So checks are satisfied (and the argue above is not applied directly).

Let  $\xi$  be a subset that connected component in the Tanner graph with a span  $> \frac{d}{4}$  at time  $t$ , and let  $\sigma_i$  the event that at time  $i$  a connected with span at least  $\sqrt{d}$  belongs to the error. Then:

$$\Pr[\xi] = \Pr[\xi \cap \cup_i^t \sigma_i] + \Pr[\xi \cap \cap_i^t \bar{\sigma}_i]$$

So denote by  $\tau$  the number of bits that require to be flipped in order to merge the flipped connected component to a one with a span greater than  $d/4$ , So  $\tau$  has to be at least:

$$\tau(\sqrt{d} + 1) \geq \frac{1}{4}d \Rightarrow \tau \geq \frac{1}{8}\sqrt{d}$$

Therefore:

$$\begin{aligned}
\Pr[\xi \cap \cap_i^t \bar{\sigma}_i] &\leq \sum_{\xi^{(t-1)}} \Pr[\xi | \xi^{(t-1)}] \Pr[\xi^{(t-1)} \cap \cap_i^t \bar{\sigma}_i] \\
&\leq \sum_{\xi^{(t-1)}} p^{\frac{1}{8}\sqrt{d}} \Pr[\xi^{(t-1)} \cap \cap_i^t \bar{\sigma}_i] \leq p^{\frac{1}{8}\sqrt{d}}
\end{aligned}$$

$$\begin{aligned}
\Pr[\xi] &= \Pr[\xi \cap \cup_i^t \sigma_i] + \Pr[\xi \cap \cap_i^t \bar{\sigma}_i] \\
&\leq t\Pr[\sigma_1] + p^{\frac{1}{8}\sqrt{d}}
\end{aligned}$$

## 10 Configuration up to $C_z^\perp$