

# Quantum Fault Tolerance Meets Toom's Contours

Michael Ben-Or   David Ponnarovsky

May 5, 2026

## 1 Notations.

We follow the standard notations commonly used in the literature. For an integer  $n$ , we denote by  $[n]$  the set of integers  $\{1, 2, \dots, n\}$ . To present a countable set, which might or might not be finite, where the number of elements does not matter, we use curly brackets with a subscript denoting the first index of the set. For example,  $\{u_i\}_{i=1}$  represents the set  $u_1, u_2, \dots$ ; For finite sets, we might add a superscript to present the size of the set. We use a colon inside curly brackets to specify a set of elements satisfying a condition, i.e., for sets  $A, B$ , the notation  $\{a : a \in A, a \in B\}$  equals  $A \cap B$ . Curly brackets without subscripts or conditions refer to singletons. For example, for a given integer  $i$ , the set  $\{u_i\}$  is the set that contains only the element  $u_i$ . For an integer  $i$ , we denote by  $1_i$  the vector consisting of one at the  $i$ th coordinate and zero elsewhere.

When not otherwise mentioned, we consider quantum states over qubits, each lying in a two-dimensional Hilbert space  $\mathcal{H}_2$ , and denote by  $\mathcal{H}_{2^n}$  the Hilbert space of an  $n$ -qubit system. Pure quantum states and their dual presentations are denoted by  $|\psi\rangle, \langle\psi|$ , and in the matrix representation, we use  $|\psi\rangle\langle\psi|$  to denote its density matrix. We will use  $\rho$  to denote a mixed state, whose density matrix is a trace one PSD matrix.

Below, a semi proof that in CSS code partial correction  $Z^e \rightarrow Z^{\hat{e}}$  doesn't add bit flip errors. It might adds a global phase. I should enter it in a section about CSS coding.

$$\begin{aligned} & X^f Z^e |w + C_z^\perp\rangle \\ &= X^f Z^e X^w |C_z^\perp\rangle \\ \text{apply } H^{\otimes n} &\mapsto Z^f X^e Z^w |C_z\rangle \\ &= (-1)^{e \cdot w} Z^f Z^w X^e |C_z\rangle \\ \text{partial correction } X^{e+\hat{e}} &\mapsto (-1)^{(e+\hat{e})(f+w)} (-1)^{e \cdot w} Z^f Z^w X^{\hat{e}} |C_z\rangle \\ \text{apply } H^{\otimes n} &\mapsto (-1)^{(e+\hat{e})(f+w)} (-1)^{e \cdot w} X^f X^w Z^{\hat{e}} |C_z^\perp\rangle \\ &= (-1)^{(e+\hat{e})(f+w)} (-1)^{e \cdot w} (-1)^{\hat{e} \cdot w} X^f Z^{\hat{e}} X^w |C_z^\perp\rangle \\ &= (-1)^{(e+\hat{e})f} X^f Z^{\hat{e}} |w + C_z^\perp\rangle \end{aligned}$$

## 2 Computation Model

Informally, a quantum circuit is a placement of a collection of quantum gates, taken from a finite set, in different space-time locations. Each space-time location, associated with a gate, encodes the time of the application and the qubits in the support of the gate. Formally

Separate between an element gate  $g \in \mathcal{G}$  and a gate, which also has a space-time location.

**Definition 2.1** (Quantum Circuit). A space-time location  $l$  is a tuple that encodes in its first coordinate time, and in its second coordinate a subset of integers referring to a support of (q)bits. For a collection of quantum gates  $\mathcal{G} = \{g: L(\mathcal{H}_2^{\otimes \Delta}) \rightarrow L(\mathcal{H}_2^{\otimes \Delta})\}$ , and integers  $w, d \in \mathbb{N}$  a quantum circuit  $C$ , with depth  $d$  and width  $w$ , is a collection of tuples  $\{u_i = (g_i, l_i)\}_{i=1}$ , where  $u_i$  are the gates of  $C$  and each  $u_i$  is composed of a gate  $g_i \in \mathcal{G}$  and a space-time location  $l_i \in \mathbb{Z}_d \times \mathbb{Z}_w^\Delta$ . The gates of  $C$  have to satisfy the constraint that for every two gates with the same time coordinate, there cannot be a non-empty intersection in their space coordinate, namely for  $u_i = (g_i, l_i), u_j = (g_j, l_j) \in C$ , the equality  $l_{i,1} = l_{j,1}$  implies  $l_{i,2} \cap l_{j,2} = \emptyset$ . We define the  $t$ -cut of  $C$  to be  $C|_t := \{u_i = (g_i, l_i): u_i \in C \text{ and } l_{i,1} = t\}$ .

To define the running of a quantum circuit  $C$  on a given (mixed) quantum state, we will need to have the notion of the directed acyclic graph associated with  $C$ .

**Definition 2.2** (Associated Computation DAG). Let  $C = \{u_i\}_{i=1}$  be a quantum circuit with depth  $d$  and width  $w$ . We define the associated computation DAG of  $C$ , denoted by  $\pi_C$ , to be the acyclic directed graph  $\pi_C = (\mathcal{U}, E)$  as follows:

1. The vertex set  $\mathcal{U}$  is the gate set of  $C$ , namely  $\mathcal{U} = \{u_i\}_{i=1}$ .
2. For  $u_i = (g_i, l_i), u_j = (g_j, l_j)$ , There is an edge  $\{u_i, u_j\} \in E$  if and only if:

$$j \in \arg \min_k \{l_{k,1} : (g_k, l_k) \in \mathcal{U}, l_{k,1} > l_{i,1} \text{ and } l_{k,2} \cap l_{i,2} \neq \emptyset\}$$

Note that there might be multiple indices that get the minimum, so the out degree might be greater than one.

In addition, suppose that  $J = \{j_1, j_2, \dots, j_k\} \subset [|C|]$  is an order of the indices of the gates in  $C$  according to the topological order induced by  $\pi_C$ . Then, we say that  $J$  is an order of indices that agrees with  $\pi_C$  and define the series of circuit  $C_J = \{C_{J,i}\}_{i=1}$  as follows:

1. The first element:  $C_{J,1} = \{u_{j_1}\}$
2. Then,  $C_{J,i+1} = C_{J,i} \cup \{u_{j_i}\}$ . We name  $C_{J,i}$  the partial computation circuit until step  $i$ .

**Definition 2.3.** Let  $\rho$  be a density matrix over  $w$  qubits, and let  $C = \{u_i\}_{i=1}$  be a quantum circuit with depth  $d$  and width  $w$ . For any density matrix  $\sigma$ , quantum gate

$g \in \mathcal{G}$ , and location  $l$ , the application of the gate-location tuple  $(g, l)$  on  $\sigma$ , denoted by  $(g, l) \circ \sigma$ , is the application of  $g$  on  $\sigma$  where wires are ordered such that the first qubit in the input to  $g$  is  $l_{2,1}$ , the second is  $l_{2,2}$ , and so on. The result of the application of  $C$  on  $\rho$ , denoted by  $C \circ \rho$ , is given by iterative applications of the gates  $\{u_i\}_{i=1}$  on  $\rho$ , according to a topological order induced by  $\pi_C$ .

We say that  $\rho_1, \rho_2, \dots, \rho_k$  is a computation prefix if  $\rho_1 = \rho$  and  $\rho_{i+1} = u'_i \circ \rho_i$ , where  $\{u'_i\}_{i=1}$  is a prefix of a topological ordering of  $\{u_i\}_{i=1}$ . If  $\{u'_i\}_{i=1}$  contains all the gates in  $\{u_i\}_{i=1}$ , then we call  $\rho_1, \rho_2, \dots, \rho_k$  a running.

**Definition 2.4** (*C Subjected to  $\mathcal{E}$  at frequency  $r$* ). Let  $C = \{u_i\}_{i=1}$  be a quantum circuit with depth  $d$  and width  $w$ . For a set of fault locations  $\mathcal{E} \subset [d] \times [w]$ , and a number  $r \in (0, 1)$  satisfies  $r^{-1} \in \mathbb{N}$ , we define the quantum circuit  $C$  *subjected to  $\mathcal{E}$  at frequency  $r$* , denoted by  $C[\mathcal{E}]_r = \{u'_i\}_{i=1}$ , to be the quantum circuit that is the result of the following map:

$$u_i = (g_i, (l_{i,1}, l_{i,2})) = \begin{cases} (g_i, (l_{i,1} + \lfloor r l_{i,1} \rfloor + 1, l_{i,2})) & u_i \in C \\ (g_i, (r^{-1} l_{i,1}, l_{i,2})) & u_i \in \mathcal{E} \end{cases}$$

Where  $r$  is not mentioned, we mean  $r = \frac{1}{2}$ , and the result of the subsection is the circuit whose gates on the odd time coordinates are the gates of  $C$ , and whose gates on the even time coordinates are the gates of  $\mathcal{E}$ . We will denote the associated computation DAG of  $C$  affected by  $\mathcal{E}$  as  $\pi[C, \mathcal{E}]$ .

**Definition 2.5** (*Faults Propagation to a Step  $\tau$* ). For a quantum circuit  $C = \{u_i\}_{i=1}$ , with depth  $d$ , an order  $J = \{j_1, j_2, \dots, j_k\} \subset [|C|]$  that agrees with  $\pi_C$ , and gate  $u_i$  in  $C$ , a *propagation of a gate  $u_i \in C$  to step  $\tau$*  is the quantum operator  $X: L(\mathcal{H}_2^w) \rightarrow L(\mathcal{H}_2^w)$  obtained by the follow process:

1. First, check if  $i$  precedes  $\tau$  in  $J$ . If not, then return  $X = I$ . Otherwise, denote  $j_l = i$ , and look for an operator  $X_1$  such that  $u_{j_l} u_{j_{l+1}} = u_{j_{l+1}} X_1$ .
2. Then, for  $k \leq \tau - l$ , look for an operator  $X_k: L(\mathcal{H}_2^w) \rightarrow L(\mathcal{H}_2^w)$ , such that  $X_{k-1} u_{j_{l+1}} = u_{j_{l+1}} X_k$ .
3. At the final stage of the process, namely where  $k = \tau - l$ , we define  $X := X_k$  to be the propagation of  $u_j$  to step  $\tau$ .

We define the *propagated computation graph*  $(\mathcal{U}', E')$ , resulted by the propagation of  $u_i$  to step  $\tau$ , as follow: Let  $u_i = u_{j_l}, \dots, u_{j_\tau}$ , be the gates at steps between  $i$  to  $\tau$ , and let  $X$  be the propagation of  $u_i$  to step  $\tau$ . Consider the tree rooted at  $u_i$  restricted to  $u_{j_l}, \dots, u_{j_\tau}$ , denote it by  $T$  and it leaves by  $L_T$ . Set  $\mathcal{U}' = \mathcal{U}/u_i \cup X$  and:

$$\begin{aligned} E' = & E \\ & / \{ \{u, u_i\} : \{u, u_i\} \in E \} \\ & / \{ \{u, v\} : u \in L_T, \{u, v\} \in E \} \\ & \cup \{ \{X, v\} : u \in L_T, \{u, v\} \in E \} \\ & \cup \{ \{u, X\} : u \in L_T \} \\ & \cup \{ \{u, v\} : \{u, u_i\}, \{u_i, v\} \in E \} \end{aligned}$$

For a quantum circuit  $C$ , a set of faults  $\mathcal{E} = \{e_i\}_{i=1}$ , an order  $J \subset [|C[\mathcal{E}]|]$  that agrees with  $\pi_{C[\mathcal{E}]}$ , and a step  $\tau \in J$ , denote by  $X_{e_i}^\tau$  the propagation of  $e_i \in \mathcal{E}$  in  $C[\mathcal{E}]$  to step  $\tau$ . Let  $j'_1, j'_2, \dots, j'_l$  be the indices of the faults in  $\mathcal{E}$  such that their corresponding positions in  $J$  precede  $\tau$ , ordered according to the reverse topological order induced by  $\pi_{\mathcal{E}}$ . The propagated error to step  $\tau$  is defined by

$$X^\tau := \prod_i X_{e_{j'_i}}^\tau$$

**Definition 2.6** (Layer Propagation.). Let  $C = \{u_i\}_{i=1}$  be a quantum circuit, and let  $U_1, U_2, \dots, U_d$  be a partition of the gates in  $C$  according to their time coordination. We name such a partition the *partition of  $C$  into layers*. The *forward layer propagation* of  $U_i$  is defined to be the action that results in the quantum circuit such that its partition into layers is  $U_1, \dots, U_{i-1}, U_{i+1}, V, U_{i+2}, \dots, U_d$ , where  $V$  is a collection of quantum gates, such that

$$\prod_{u \in U_i} u \prod_{u \in U_{i+1}} u = \prod_{u \in U_{i+1}} u \prod_{u \in V} u$$

For  $i' > i$ , the *forward layer propagation* of  $U_i$  to time  $i'$  is defined by propagating  $U_i$  forward for  $i' - i$  times. In the resulting circuits of the layer propagation actions, we update the time coordinates to match their application turn.

**Claim 2.7.** The result of forward layer propagation has a bounded fan-in.

[COMMENT] Prove it.

**Definition 2.8** ( $C \sim_P C'$ ). We define the equivalence relation over quantum circuits  $\sim_P$  to be satisfied if any of them can be obtained from the other by propagating layers. Namely, for quantum circuits  $C, C'$ , such that the partition of  $C$  into layers is  $U_1, \dots, U_d$ , we have  $C \sim_P C'$  if there is a subset of layers  $\{U_i\}_{i=1} \subset \{U_1, \dots, U_d\}$  and subsets of time points  $\{t_i\}_{i=1}$  such that  $C'$  is the result of propagating each of the  $U_i$  to time  $t_i$  in  $C$  according to the order indexed by  $i$ .

Note that, by definition, we find that equivalence between quantum circuits by  $\sim_P$  implies an equivalence of their action. More specifically, for a mixed state  $\rho$  and quantum circuits  $C$  and  $C'$  such that  $C \sim_P C'$ , we have:

$$C\rho C^\dagger = C'\rho C'^\dagger$$

**Definition 2.9** (Deviation). For a quantum circuit  $C$ , a set of faults  $\mathcal{E}$ , and step  $\tau$ , let  $X^\tau$  be the propagated error at step  $\tau$ . We define the deviation of  $C[\mathcal{E}]$  from  $C$  at step  $\tau$  by the set of indices on which  $X^\tau$  acts non-trivially.

**Definition 2.10.** Using the same notations as in Definition 2.9, for a set of unitary operators  $\mathcal{S} = \{s_i: \mathcal{H}_2^w \rightarrow \mathcal{H}_2^w\}_{i=1}$ , we define the propagated error up to stabilizers in  $\mathcal{S}$  as  $X^\tau[\mathcal{S}] := \min_{s \in \mathcal{S}} \{w(sX)\}$ . Then, we define the deviation of  $C[\mathcal{E}]$  from  $C$  at step  $\tau$  up to stabilizers in  $\mathcal{S}$  to be the subset of coordinates on which  $X^\tau[\mathcal{S}]$  acts non-trivially.

The following, extends the notion of Tanner graphs to the context of quantum circuits.

**Definition 2.11** (Tanner Graph of a Register). For a quantum circuit  $C$ , a register  $R$  is a subset of qubits in  $C$ . The *Tanner graph* at step  $\tau$  of register  $R$ , denoted by  $\mathcal{T}_{R,\tau}$ , is the graph defined as follows:

1. If there is a code  $\mathcal{C}$ , such that at turn  $\tau$ , the qubits in  $C$  are an encoding of  $\mathcal{C}$ , then  $\mathcal{T}_{R,\tau}$  is the Tanner graph of  $\mathcal{C}$ .
2. Otherwise,  $\mathcal{T}_{R,\tau}$  is the empty graph.

Where we consider a range of turns in which  $R$  is encoded in the same code, and the turn is clear from the context, we will be brief by omitting the turn subscript.

**Definition 2.12** (Generalized Tanner Graph of a Circuit). For a quantum circuit  $C$ , and registers  $a$  and  $b$  in  $C$ , the *generalized Tanner graph* of the register obtained by taking their union  $a \cup b$  is  $\mathcal{T}_{a \cup b} := \mathcal{T}_a \cup \mathcal{T}_b$ . For a partition of the qubits into registers  $\{R_i\}_{i=1}$ , the *generalized Tanner graph* of a quantum circuit  $C$ , at step  $\tau$ , according to the partition  $\{R_i\}_{i=1}$ , is the graph obtained by:

$$\mathcal{T}_{C,\{R_i\}_{i=1},\tau} := \bigcup_{i=1} \mathcal{T}_{R_i,\tau}$$

When the partition is clear, we will brief and use  $\mathcal{T}_C$  to define the generalized Tanner graph of  $C$ .

**Definition 2.13** (Clustered Deviation). For a quantum circuit  $C$ , a register  $R$  of  $C$ , and a set of faults  $\mathcal{E}$ , let  $\mathcal{T}_R$  be the Tanner graph of  $R$ , and let  $\mathcal{D}_\tau$  be the deviation of  $C[\mathcal{E}]$  from  $C$  on  $R$  at step  $\tau$ . Denote by  $\mathcal{T}_R|_{\mathcal{D}_\tau}$  the subgraph of  $\mathcal{T}_R$  induced by the restriction of the left vertices (bits) to the indices in  $\mathcal{D}_\tau$ . Let us abuse and extend the connection notion to Tanner graphs by defining that two left vertices are connected if and only if they share a right neighbor. We define the deviation clusters at step  $\tau$  to be the connected components of  $\mathcal{T}_R|_{\mathcal{D}_\tau}$ . The size of each deviation cluster is the number of right vertices it contains.

**Claim 2.14.** Let  $C$  be a quantum circuit composed of gates with bounded fan degree  $\Delta \in \mathbb{N}$ . In addition, let  $R$  be a quantum register encoded by a quantum error correction code  $\mathcal{C}$  with Tanner graph  $\mathcal{T}$  that has left and right degrees  $\Delta_1$  and  $\Delta_2$ . Fix a set of faults  $\mathcal{E}$ , and an order  $J$  that agrees with  $\pi_{C[\mathcal{E}]}$ . For any  $\tau$ , denote by  $Y_1^{(\tau)}, Y_2^{(\tau)}, \dots, Y_l^{(\tau)}$  the deviation clusters at step  $\tau \in J$ . Assume that at step  $\tau_1 \in J$  there is a deviation cluster  $Y_i^{(\tau_1)}$  of size  $|Y_i^{(\tau_1)}|$ . Then there is a deviation cluster at step  $\tau_1 - 1$ , let's denote it by  $Z$ , such  $Z \cap Y_i^{(\tau_1)} \neq \emptyset$  and  $Z$  is of size at least:

$$|Z| \geq \frac{1}{\Delta \Delta_1 \Delta_2} |Y_i^{(\tau_1)}|$$

Furthermore there is a deviation cluster at step  $\tau_1 - 1$ , let's denote it by  $Z'$ , such  $Z' \cap Y_i^{(\tau_1)} \neq \emptyset$ , and  $Z'$  is of size at most:

$$|Z'| \leq \Delta \Delta_1 \Delta_2 |Y_i^{(\tau_1)}|$$

Add: that when it isn't mentioned, we assume that the error at time  $t$  can be decomposed into a product  $\Rightarrow$  at each step, a fault is supported on at most one (q)bit.

*Proof.* Denote by  $u = (g, l)$  the gate at step  $\tau$ . If  $u$  acts trivially on the (q)bits in  $Y_i^{(\tau_1)}$  then they also belong to deviation on register  $R$  at step  $\tau_1 - 1$ , Since they are connected in  $\mathcal{T}$  they form a deviation cluster on register  $R$  at step  $\tau_1 - 1$  and we got the desired. Thus it is left to handle the case in which  $u$  acts non-trivially on (q)bits in  $Y_i^{(\tau_1)}$ . Since  $u$  acts on at most  $\Delta$  qubits, and each qubit has at most  $\Delta_1(\Delta_2 - 1)$  neighbors at  $\mathcal{T}$  it follows that  $u$  acts non trivially on at most

$$\Delta + \Delta \Delta_1 (\Delta_2 - 1) \leq \Delta \Delta_1 \Delta_2$$

deviation clusters at step  $\tau_1 - 1$ . Let  $l$  be integer no greater than  $\Delta \Delta_1 \Delta_2$  and let  $Y_1, Y_2, \dots, Y_l$  be the deviation cluster, on register  $R$ , at step  $\tau_1 - 1$ , on which  $u$  acts non trivially, since  $Y_i^{(\tau_1)} \subset \bigcup_j Y_j$ . We have that:

$$\begin{aligned} |Y_i^{(\tau_1)}| &\leq \sum_j |Y_j| \leq \Delta \Delta_1 \Delta_2 \max_j |Y_j| \\ \Rightarrow \max_j |Y_j| &\geq \frac{1}{\Delta \Delta_1 \Delta_2} |Y_i^{(\tau_1)}| \end{aligned} \tag{1}$$

Complete the second inequality. Should consider to erase what have already written about it. To get the second inequality, we use the fact that  $u$  is a unitary gate, in particular,  $u$  is reversible. Let  $C_{J, \tau_1} = u_1, \dots, u_{\tau_1}$  be the partial computation circuit until step  $\tau_1$ . Observe that  $C_{J, \tau_1 - 1}$  is obtained by stacking  $u_{\tau_1}^\dagger$  as a final stage to  $C_{J, \tau_1}$ , denoted by  $u^\dagger \circ C_{J, \tau_1}$ . Thus, the deviation from  $C_{J, \tau_1}$  at step  $\tau_1 - 1$  equals the deviation from  $u^\dagger \circ C_{J, \tau_1}$  at step  $\tau_1 + 1$ . By eq. (1),  $\square$

**Corollary 2.15** (Intermediate Value Theorem). Let  $\alpha = 1/(\Delta \Delta_1 \Delta_2)$ , and let  $x$  be an integer. Assume that at the initial step  $\tau_o$ , there is no deviation cluster of size more than  $\alpha x$ , and that at a step  $\tau_1 \in J$ , such that  $\tau_1 > \tau_o$ , there is a deviation cluster  $Y^{(\tau_1)}$  of size  $|Y^{(\tau_1)}| > x$ . Then there is a step  $\tau_\star \in J$  such that  $\tau_o < \tau_\star < \tau_1$  for which there is a deviation cluster  $Z$  at step  $\tau_\star$ , satisfying that the size of  $Z$  is in the range  $(\alpha x, x)$ .

*Proof.* Assume through contradiction that for any turn  $\tau \in (\tau_o, \tau_1)$ , all the deviation clusters are of size smaller than  $\alpha x$ . Particularly, at turn  $\tau_1 - 1$ , all the deviation clusters are of size smaller than  $\alpha x$ . Yet, by Claim 2.14, the existence of the deviation cluster  $Y^{(\tau_1)}$ , at size  $d$ , implies the existence of a deviation cluster at turn  $\tau_1 - 1$ , of size at least  $\alpha x$ , which leads to a contradiction.  $\square$

**Claim 2.16.** Let  $C$  be a quantum circuit with order  $J$  and depth  $d$ , and let  $\mathcal{E} \sim \mathcal{N}_{\mathcal{E}}$ , so the circuit  $C[\mathcal{E}]$  is a random variable, and it is fan-bounded by  $\Delta$ . For a time  $t \leq 2d$  and an integer  $x \in \mathbb{N}$ , let  $A$  be the event in which, until time  $t$ , there is a time  $t' \leq t$  such that there is a deviation cluster of size greater than  $x$ . For any step  $\tau \in J$ , let  $B_{\tau}$  be the event that step  $\tau$  is the first step to have a deviation cluster of size in the range  $(\alpha x, x)$ . Then:

$$\Pr[A] \leq 2wd \max_{\tau \in [2wd]} \Pr[B_{\tau}]$$

*Proof.* By Corollary 2.15, we have that the occurrence of a deviation cluster of size greater than  $x$  before time  $t$  implies that there is a step  $\tau \in J$ , such that  $\tau \leq 2wt$ , in which there is a deviation cluster of size greater than  $\alpha x$ . Thus,

$$A \subset \bigcup_{\tau \in [2wt]} B_{\tau} \subset \bigcup_{\tau \in [2wd]} B_{\tau}$$

So by using the union bound we get:

$$\begin{aligned} \Pr[A] &\leq \Pr\left[\bigcup_{\tau \in [2wd]} B_{\tau}\right] \leq \sum_{\tau \in [2wd]} \Pr[B_{\tau}] \\ &\leq 2wd \max_{\tau \in [2wd]} \Pr[B_{\tau}] \end{aligned}$$

□

**Definition 2.17** (Noise Channel and The Local Stochastic Noise Property). A *noise channel*  $\mathcal{N}_{\mathcal{E}}$  is a distribution over sets of faults, we will denote the outcome of  $\mathcal{N}_{\mathcal{E}}$  by  $\mathcal{E}$ . For  $p \in (0, 1)$  we say that a noise channel  $\mathcal{N}_{\mathcal{E}}$  is a *local stochastic*, with noise rate  $p$ , if the following holds: For every subset of locations  $S \subset \mathbb{N} \times \mathbb{N}$ , whose locations have the same time coordinate, the probability that  $S$  is covered by  $\mathcal{E}$  is less than:

$$\Pr_{\mathcal{E} \sim \mathcal{N}_{\mathcal{E}}}[S \subset \mathcal{E}] \leq p^{|S|}$$

Observes that a quantum circuit  $C$  and a noise channel  $\mathcal{N}_{\mathcal{E}}$  induce a subjection  $C[\mathcal{E}]$ , which is a random variable.

**Claim 2.18.** Let  $\mathcal{N}_{\mathcal{E}}$  be a local stochastic noise channel with noise rate  $p$ , let  $C$  be a quantum circuit with bounded fan-in  $\Delta$ , and let  $c$  be an integer. We define  $q$  to be the number that depends on  $p$ ,  $\Delta$ , and  $c$  as:  $q := (\Delta^c p)^{\frac{1}{\Delta^c}}$ . There is a local stochastic noise channel  $\mathcal{N}_{\mathcal{E}_q}$ , with noise rate  $q$  that induces a subjection  $C[\mathcal{E}_q]_{\frac{1}{c}}$  such that:

$$C[\mathcal{E}]_{\frac{1}{2}} \sim_P C[\mathcal{E}_q]_{\frac{1}{c}}$$

*Proof.* We need to show that any subjection outcome  $C[\mathcal{E}]$  can be transformed by propagation to a circuit  $C'$  that is also a result of subjecting  $C$  to a set of faults  $\mathcal{E}'$  at

frequency  $\frac{1}{c}$ , and that  $\mathcal{E}'$  satisfies the local noise property with a noise rate  $q$ . Assume that  $c$  is even, and consider the layer propagations  $U_i \rightarrow i + c$ .

$$\begin{aligned} U_0, U_1, U_2, \dots, U_{c-1}, U_c, U_{c+1}, \dots, U_c &\mapsto \\ U_0, U_1, U_3, U_5, \dots, U_{c-1}, V_2, V_4, \dots, V_{c-2}, U_c, U_{c+1}, U_{c+3}, \dots &\mapsto \\ V^{(0)}, U_1, U_3, U_5, \dots, U_{c-1}, V^{(1)}, U_{c+1}, U_{c+3}, \dots &\mapsto \end{aligned}$$

We grouped the errors together. □

For that we need the follow property, for every subset of qubits  $S$ , there is subset of qubits  $S'$  at size at least  $(1 - \gamma)|S|$  such that for  $S$  to be deviated, all the qubits in  $S'$  have to absorb fault. Now suppose that the gate is at depth at most  $\Delta$ , and the fanning is at most  $c_f$ , then we get:

$$\begin{aligned} \Pr[S \in \mathcal{D}] &\leq (c_f^\Delta p)^{|S'|} \leq (c_f^\Delta p)^{(1-\gamma)|S|} \\ \Rightarrow p &\rightarrow (c_f^\Delta p)^{1-\gamma} \end{aligned}$$

A trivial lower bound for  $1 - \gamma$  is .

Change letters: the  $d$  in the model section refers to a size that is comparable to the code distance where here  $d$  refers to the dimension of the Toric complex

### 3 The $(d, d')$ -Dimensional Toric Code and the SWEEP Rule

**Definition 3.1** ( $d$ -Dimensional Toric Complex). For integers  $d$  and  $n$ , denote the group resulting from the  $d$ -directed power of the cyclic group of order  $n$  by  $\mathcal{G}_n^d = \mathbb{Z}_n^{\times d}$ , and denote the standard generating set of  $\mathcal{G}_n^d$  by  $S = \{1_i : i \in [d]\}$ . We define the  $d$ -dimensional toric complex to be the hypergraph obtained from the Cayley graph  $\text{Cay}(\mathcal{G}_n^d, S)$  by taking its vertices as the 0-faces and for any  $i \in [d - 1]$  we define the  $i$ -faces as follow: First, for a vertex  $v \in \mathcal{G}_n^d$  and for a subset of generators  $S' \subset S$  of size  $|S'| = i$ , the  $i$ -face rooted at  $v$  and spanned by  $S'$ , denoted by  $\theta_{v, S'}$ , is:

$$\theta_{v, S'} := \bigcup_{I \subset S'} \left\{ \left( \prod_{g \in I} g \right) v \right\}$$

The  $i$ -faces of the complex are the collection of all the possible rooted spanned  $i$ -faces induced by  $\text{Cay}(\mathcal{G}_n^d)$ . We will denote them by  $\mathcal{F}_i$ .

Additionally, we name  $n$  and  $d$ , the *side length* and the *dimension* of the complex.

**Definition 3.2** (Positive Edges and Cubes). For  $u, v \in \mathcal{G}_n^d$ , and  $S'_1, S'_2 \subset [d]$ , with sizes  $|S'_1| = |S'_2| = i$ , we say that the  $i$ -faces  $\theta_{v, S'_1}$  and  $\theta_{u, S'_2}$  are adjacent if the intersection  $\theta_{v, S'_1} \cap \theta_{u, S'_2}$  is an  $i - 1$ -face. In addition, if  $\theta_1$  and  $\theta_2$  are  $i$  and  $i - 1$ -faces such that  $\theta_2 \subset \theta_1$ , then we say that  $\theta_2$  is a *positive edge relative to face*  $\theta_1$  if they are rooted at the same vertex. Furthermore, we say that  $\theta_1$  is a *positive cube* of  $\theta_2$ . When the  $i$ -face is clear from the context, we will abbreviate and say that  $\theta_2$  is *positive*.



**Example 3.3.** For example, let  $v, u \in \mathcal{G}_n^d$  and  $g_1 \neq g_2 \in S$ . Consider the 2-face  $\theta = \{v, g_1v, g_2g_1v, g_2v\}$ . Then, for  $x, y \in \mathcal{G}_n^d$ , we say that an edge (1-face)  $\{x, y\}$  is *positive relative to the face  $\theta$*  if  $x = v$  and  $y$  is either  $g_1v$  or  $g_2v$ .

We denote the complex by  $G = (V, E, F)$ , where  $V$ ,  $E$ , and  $F$  are the 0, 1, and 2 faces.

Add a paragraph that explains that we choose not to use a chain-complex since we earn nothing from it i.e don't care about topology properties. Yet mention that chain-complex formulation proved itself as an effective tool to compute the properties of the codes we use.

**Definition 3.4** (Assignment on The  $(d', d)$ -Toric Complex). Let  $\mathcal{T}$  be a  $d$ -dimensional complex. Each of its  $i$ -face sets naturally induces a vector space  $\mathbb{F}_2^{|\mathcal{F}_i|}$ . We call a binary vector  $z \in \mathbb{F}_2^{|\mathcal{F}_i|}$  an assignment on the  $i$ -faces of  $\mathcal{T}$ . For a vertex  $v$  and a subset of generators  $S' \subset S$  of size  $|S'| = i$ , we use the notation  $\theta_{\mathbf{v}, S'}$  to represent the unit vector in  $\mathbb{F}_2^{|\mathcal{F}_i|}$  that has 0 in every coordinate except the one associated with the  $i$ -face  $\theta_{v, S'}$ . Then

$$\left\{ \theta_{\mathbf{v}, S'} : v \in \mathcal{F}_0, S' \subset S \text{ s.t } |S'| = i \right\}$$

is a base to  $\mathbb{F}_2^{|\mathcal{F}_i|}$  **Not exactly base, they are linear dependent.** Let  $z$  be an assignment on the  $i$ -faces. Let  $v_1, \dots, v_k \in \mathcal{F}_0$  and  $S_1, \dots, S_k \subset S^{\times k}$  be the generator subsets such that  $\theta_{v_1, S'_1}, \theta_{v_2, S'_2}, \dots, \theta_{v_k, S'_k}$  are the collection of  $i$ -faces that support  $z$ , namely:

$$z = \sum_{i=1}^k \theta_{\mathbf{v}_i, S'_i}$$

We refer to  $\theta_{v_1, S'_1}, \theta_{v_2, S'_2}, \dots, \theta_{v_k, S'_k}$  as the collection of faces that form  $z$ , or briefly, the faces form  $z$ . For  $i > 0$ , we define the linear map  $\partial : \mathbb{F}_2^{|\mathcal{F}_i|} \rightarrow \mathbb{F}_2^{|\mathcal{F}_{i-1}|}$  such that

$$\partial \theta_{\mathbf{v}, S'} := \sum_{g \in S'} \theta_{\mathbf{v}, S'/g} + \sum_{g \in S'} \theta_{\mathbf{g}\mathbf{v}, S'/g}$$

For  $i < d$ , we define the linear map  $\partial^* : \mathbb{F}_2^{|\mathcal{F}_i|} \rightarrow \mathbb{F}_2^{|\mathcal{F}_{i+1}|}$  such that

$$\partial^* \theta_{\mathbf{v}, S'} := \sum_{g \in S/S'} \theta_{\mathbf{v}, S' \cup g} + \sum_{g \in S'} \theta_{\mathbf{g}^{-1}\mathbf{v}, S' \cup g}$$

We define the *restriction of  $z$  to a vertex  $v$*  to be the linear map  $|_v : \mathbb{F}_2^{|\mathcal{F}_i|} \rightarrow \mathbb{F}_2^{|\mathcal{F}_i|}$  such that:

$$\theta_{\mathbf{u}, S'}|_v := \begin{cases} \theta_{\mathbf{u}, S'} & v \in \theta_{\mathbf{u}, S'} \\ 0 & \text{else} \end{cases}$$

We define the *front of  $z$  to a vertex  $v$*  to be the linear map  $|_{v+} : \mathbb{F}_2^{|\mathcal{F}_i|} \rightarrow \mathbb{F}_2^{|\mathcal{F}_i|}$  such that:

$$\theta_{\mathbf{u}, S'}|_{v+} := \begin{cases} \theta_{\mathbf{u}, S'} & u = v \\ 0 & \text{else} \end{cases}$$

We say that an assignment  $z$  is *connected* if the graph that is induced by taking the faces that support  $z$  as vertices and defining the edges to be the adjacency relation between them is connected. We define the connected components of  $z$  to be its subsets such that each of them is a connected assignment. For a  $j$ -face  $f$ , we say that  $f \in z$  if  $z$  has a support on  $\theta_{\mathbf{v}, \mathbf{S}'}$  such that  $f \in \theta_{\mathbf{v}, \mathbf{S}'}$ . Consider an assignment  $z \in \mathbb{F}_2^{\mathcal{F}_i}$  and a vertex  $v \in \mathcal{G}_n^d$ . We say that  $z$  is a *rooted assignment* at vertex  $v$  if there are subsets  $S'_1, \dots, S'_k \subset S^{\times k}$  such that each of  $S'_j$  is at size  $i$  and  $z$  is decomposed as:

$$z = \sum_j^k \theta_{\mathbf{v}, \mathbf{S}'_j}$$

Put differently,  $z$  is the result of taking a collection of  $i$ -faces that are rooted at  $v$ .

Add a proof for  $\partial\theta_{\mathbf{v}, \mathbf{S}'} \cdot \partial^*\theta_{\mathbf{u}, \mathbf{S}''} = 0 \Rightarrow$  gives a CSS code.

Need to decide on what should I elaborate, for example should I provide a proof for the distance and the rate of the code? [Den+02]

**Definition 3.5** ( $(d, d')$  Toric Code). Let  $d, d'$  be integers such that  $1 < d' < d$ . For any integer  $n$ , we construct the  $n$ th instance of the quantum code family  $(d, d')$  *Toric Code* as follows: Let  $G = (\mathcal{F}_0, \dots, \mathcal{F}_d)$  be the  $d$ -dimensional toric complex obtained from  $\mathcal{G}_n^d$  as defined in Definition 3.1. We associate the (q)bits with the  $\mathcal{F}_{d'}$ , the  $X$ -checks with the  $\mathcal{F}_{d'-1}$ , and the  $Z$ -checks with the  $\mathcal{F}_{d'+1}$ . An  $X$ -check,  $c_x$ , is satisfied if the parity of (q)bits set on the  $d'$ -face containing  $c_x$  is even. Similarly, a  $Z$ -check,  $c_z$ , is satisfied if the parity, in the Hadamard basis, of the (q)bits set on the  $d'$ -faces contained in  $c_z$  is even.

**Definition 3.6** (Sweep Rule). Let  $z$  be an assignment over the  $d'$ -faces, The *sweep rule decoding procedure* at vertex  $v \in \mathcal{G}_n^d$ , defining as follow:

1. For the  $\mathcal{C}_X$  code. Look for a rooted assignment  $\tilde{z}$  at  $v$  such that  $(\partial\tilde{z})|_{v+} = \partial(z)|_{v+}$ . Namely, the  $X$ -checks rooted at  $v$  get the same syndromes for  $z$  and  $\tilde{z}$ . If there is one, then flip  $\tilde{z}$ .
2. Apply the Hadamard gate on any  $d'$  face containing  $v$ .
3. For the  $\mathcal{C}_Z$  code. Look for a rooted assignment  $\tilde{z}$  at  $v$  such that  $(\partial^*\tilde{z})|_{v+} = \partial^*(z)|_{v+}$ . Namely, the  $Z$ -checks rooted at  $v$  get the same syndromes for  $z$  and  $\tilde{z}$ . If there is one, then flip  $\tilde{z}$ .
4. Apply the Hadamard gate again on any  $d'$  face containing  $v$  to reverse the second stage.

Stages 1 and 3 involve measurement of stabilizers.

**Definition 3.7** (Decoding Gadget). Consider a quantum circuit  $C$ , for a quantum gate  $u$  in  $C$ , with fun-in  $\Delta$ , we say that  $u$  is a *decoding gadget* if it has the following form:

1. The wires of  $u$  wires can be decomposed into two subsets  $A$  and  $B$  such that the wires in  $A$  are reset wires, and there exists an encoded register  $R$  in  $C$  such that the wires in  $B$  are all set in  $R$ .
2. For a set of faults  $\mathcal{E}$ , there is a Pauli gate  $u'$  that acts only on  $B$  such that the action of  $u$  in  $C[\mathcal{E}]$  equals  $u'$ . Put differently, the quantum circuit obtained from  $C[\mathcal{E}]$  by replacing  $u$  with  $u'$  equals  $C[\mathcal{E}]$ .

**Definition 3.8** (Unitary Decoding Representation). Let  $C = \{u_i\}_{i=1}$  be a quantum circuit, and let  $\mathcal{U} = \{u'_i\}_{i=1} \subset C$  be a subset of the gates in  $C$  that are decoding gadgets. For a set of faults  $\mathcal{E}$ , we define the *unitary decoding representation* of  $C[\mathcal{E}]$  to be the quantum circuit obtained by replacing each of the gates in  $\mathcal{U}$  with their corresponding Pauli gate according to  $\mathcal{E}$ .

**Claim 3.9.** The local gates performing Toom's rule are decoding gadgets.

*Proof.* [prove that](#). □

**Definition 3.10** (Registers Coordinates System). Let  $C = \{u_i\}_{i=1}$  be a quantum circuit, with width  $w$  and registers  $\mathcal{R} = R_1, \dots, R_k$ , each of length at most  $n$ . For register  $R_i$ , we define the register identifier  $\mathbf{1}_{R_i} = \mathbf{1}_i$ . Then, we define the map  $\phi_{\mathcal{R}}: \mathbb{Z}_w \rightarrow \mathbb{Z}_R \times \mathbb{Z}_n$  to return, for a space-location coordinate  $x \in \mathbb{Z}_w$ , the tuple consisting of the identifier of the register containing  $x$ , and its relative coordinate within the register. Namely,

$$\psi_{\mathcal{R}}(x) = (\mathbf{1}_{R_j}, x')$$

Where  $R_j$  is the register containing  $x$ , and  $x'$  is the index of the qubit, set at the  $x$  location, in  $R_j$ . The presentation of  $C$  in the *registers coordinate system* is the encoding by a set of gates  $\{u'_j\}_{j=1}$ , which is the result of replacing each space-location of the original gates with the applications of  $\psi_{\mathcal{R}}$  on them. Namely, for each  $u_i = (g_i, l_i) \in C$ , we map its space-location coordinate  $l_{i2} = (l_{i2}^{(1)}, \dots, l_{i2}^{(\Delta)})$  to:

$$l_{i2} \mapsto (\psi_{\mathcal{R}}(l_{i2}^{(1)}), \dots, \psi_{\mathcal{R}}(l_{i2}^{(\Delta)}))$$

For brevity, we name the first and second coordinates in the result of  $\psi_{\mathcal{R}}$  the *register coordinate* and the *inner coordinate*.

## 4 Infinite Torics

**Definition 4.1** (Shifting in Registers System). Consider the register coordinate system induced by a quantum circuit of width  $w$ , with  $R$  registers, each of length at most  $n$ . For an integer  $r \in \mathbb{Z}$ , we define the  $r$ -shifting function  $\text{Sh}_r: (\mathbb{Z}_R \times \mathbb{Z}_n)^{\times \Delta} \rightarrow (\mathbb{Z}_R \times \mathbb{Z})^{\times \Delta}$  to act on space-location coordinates as follows:

$$\text{Sh}_r(l_1, \dots, l_k) = (\text{Sh}_r(l_1), \dots, \text{Sh}_r(l_k))$$

$$\text{Where } \text{Sh}_r(l_i) = \begin{cases} l_i + r & \text{if } l_i \text{ is an inner coordinate} \\ l_i & \text{else} \end{cases}$$

**Definition 4.2.** Let  $n$  be an integer, and let  $C = \{u_i\}_{i=1}$  be a quantum circuit with registers  $\mathcal{R} = R_1, \dots, R_k$  such that each register  $R_i$  is a toric encoding with a side length  $n$ . In addition, all the decoding gadgets of  $C$  are sweep rule gadgets; let us denote them by  $\mathcal{U}$ . We define the  $\infty$ -extension of  $C$ , denoted by  $C^\infty$ , as follows:

1. For a register  $R_i \in \mathcal{R}$ , denote by  $d_i, d'_i$  the dimension of the toric complex that induces the code, and the dimension of the faces that are associated with the bits. Each register  $R_i$  in  $\mathcal{R}$  is mapped to a register  $R'_i$ , which is a  $(d_i, d'_i)$ -dimensional, infinite side length, toric encoding. Since the mapping between registers is one-to-one, we use the identifier  $\mathbf{1}_{R_i}$  to identify also  $R'_i$ .
2. Each gate  $u_i \in C/\mathcal{U}$  is mapped to:

$$u_i = (g_i, l_i) \mapsto \bigcup_{j=-\infty}^{\infty} \{(g_i, (\text{Sh}_{jn}(l_i)))\}$$

3. Each gate  $u \in \mathcal{U}$  is mapped to the collection of sweep rule gadgets  $\{v_j\}_{j=-\infty}^{\infty}$  such that each of them has the same time coordinate as  $u$ , is in the register that corresponds to the image of  $u$ 's register, and is rooted at  $\text{root}(v) + jn$ .

For a set of faults  $\mathcal{E} = \{f_i\}_{i=1}$ , we denote by  $C^\infty[\mathcal{E}^\infty]$  the  $\infty$ -extension of  $C[\mathcal{E}]$ .

**Claim 4.3.** Let  $u = (g, l)$  and  $v_0, v_{\pm 1}$  plantation of  $u$  at locations  $\text{root}(u)$  and  $\text{root}(u) \pm n$ . Then  $v_0 v_{\pm 1}|_{[w]} = u$ .

*Proof.* [Prove it](#) □

**Definition 4.4.** For a quantum circuit  $C$ , of the form defined in Definition 4.2, and a set of faults  $\mathcal{E}$ , denote:

$$\begin{aligned} \rho &:= C[\mathcal{E}] |0\rangle\langle 0| C[\mathcal{E}]^\dagger \\ \rho_\infty &:= C^\infty[\mathcal{E}^\infty] |0\rangle\langle 0| (C^\infty[\mathcal{E}^\infty])^\dagger \end{aligned}$$

We say that a quantum circuit  $C$  is  $\infty$ -compatible if, for any set of faults  $\mathcal{E} = \{f_i\}_{i=1}$ , it holds that:

$$\rho = \mathbf{Tr}_{/[w]}(\rho_\infty)$$

**Claim 4.5.** Let  $C$  be quantum circuit of the form defined in Definition 4.2. Then  $C$  is  $\infty$ -compatible.

*Proof.* Its enough to prove the claim for circuits in their unitary decoding representation. We prove it by induction on the number of the gates.

1. Base. For an empty circuit,  $\rho_\infty$  is just the infinite string of zeros; thus, the restriction of it to  $[w]$  locations, namely  $\mathbf{Tr}_{/[w]}(\rho_\infty)$ , yields a single matrix element corresponds to the  $w$ -length string of zeros, which is exactly  $\rho$ .

2. Step. Assume the correction of the claim for any circuit, with less than  $\tau$  steps, and consider a circuit  $C$  with  $\tau$  steps. Denote by  $C'$  the circuit obtained from  $C$  by erasing the last gate in  $C$ , observes that  $C'$  is a circuit with  $\tau - 1$  steps. Denote by  $\rho'$  and  $\rho'_\infty$  the: [Enter it](#). By the induction assumption we have:

$$\rho' = \mathbf{Tr}_{/[w]}(\rho'_\infty)$$

Denote the last gate in  $C$  by  $u$ , and by  $I = \{v_j\}_{j=-\infty}^\infty$  the collection of gates in  $C^\infty$  to which  $u$  is mapped. Let us split into cases, depending on whether  $u$  is or is not a decoding gate.

- (a) Suppose that  $u$  is not a decoding gate. Let  $g$  and  $l$  be the gate element and the location of  $u$ . By the definition of  $\infty$ -extension,  $I$  equals:

$$I = \bigcup_{j=-\infty}^{\infty} \{(g, (\text{Sh}_{jn}(l)))\}$$

So there is only one gate in  $I$ , corresponding to  $j = 0$ , that has support on qubits with spatial location in the range  $[w]$ . Denote that gate by  $v_0$ .

$$\begin{aligned} \mathbf{Tr}_{/[w]}(\rho_\infty) &= \mathbf{Tr}_{/[w]} \left( \prod_{j=-\infty}^{\infty} v_j \rho'_\infty \prod_{j=-\infty}^{\infty} v_j^\dagger \right) \\ &= v_0 \mathbf{Tr}_{/[w]} \left( \prod_{j=-\infty, j \neq 0}^{\infty} v_j \rho'_\infty \prod_{j=-\infty, j \neq 0}^{\infty} v_j^\dagger \right) v_0^\dagger \\ &= v_0 \mathbf{Tr}_{/[w]} \left( \prod_{j=-\infty, j \neq 0}^{\infty} v_j^\dagger \prod_{j=-\infty, j \neq 0}^{\infty} v_j \rho'_\infty \right) v_0^\dagger \\ &= v_0 \mathbf{Tr}_{/[w]}(\rho'_\infty) v_0^\dagger \\ &= v_0 \rho' v_0^\dagger \\ &= \rho \end{aligned}$$

- (b) In the case that  $u$  is a decoding gate, observe that there are at most two gates in  $I$  that have support on the qubits with space-location in  $[w]$ . If there is exactly one, then repeating the non-decoding gate case while setting this gate as  $v_0$  gives the desired result. Else, in the case where two gates act non-trivially on qubits at  $[w]$ , denote those gates by  $v_0, v_1$ . Since we assumed that  $C$  is given in its unitary decoding representation, it holds that  $v_0, v_1$  are Paulis, in particular a product operator. Denote their product decompositions by  $v_0 = \prod_k v_0^{(k)}$  and  $v_1 = \prod_k v_1^{(k)}$ . Thus, by repeating the non-decoding case, but extracting from the infinite product only the terms in  $v_0$  and  $v_1$  that act non-trivially at  $[w]$ , we get:

$$\mathbf{Tr}_{/[w]}(\rho_\infty) = v_0 v_1|_{[w]} \rho' (v_0 v_1|_{[w]})^\dagger$$

Moreover, since  $v_0$  and  $v_1$  are a plantation of  $g$  in locations  $\text{root}(u)$  and  $\text{root}(u) \pm n$ , then by Claim 4.3, we have that  $u = v_0 v_1|_{[w]}$ ; therefore:

$$\begin{aligned}\text{Tr}_{/[w]}(\rho_\infty) &= v_0 v_1|_{[w]} \rho' (v_0 v_1|_{[w]})^\dagger \\ &= u \rho' u^\dagger \\ &= \rho\end{aligned}$$

□

Add that the computation graph is lifted naturally to the  $\infty$ -extension, since the image of a mapping are overlapped gates, we can execute them simultaneously. Same for the base graph  $G_o \rightarrow G_o^\infty$ .

## 5 Shrinking

consider changing the name to something like hight lines / contour in order to hint on the similarity to topography map. In addition, the embedding inside  $\mathbb{R}^d$  should appears before that definition.

**Definition 5.1** (Potential Walls). We define the potential walls to be a functions collection from the vertices of  $\mathcal{G}_n^d$  to the reals as follow:

1. For any coordinate  $i \in [d]$ , we define  $L_i: \mathcal{G}_n^d \rightarrow \mathbb{R}$  to act as  $L_i(v) = v_i$ .
2. For any subset of coordinates  $I \subset [d]$  we define  $L_I: \mathcal{G}_n^d \rightarrow \mathbb{R}$  to act as  $L_I(v) = -\sum_{i \in I} v_i$ .

For a subset of vertices  $\Gamma \subset \mathcal{G}_n^d$ , a vertex  $v \in \Gamma$ , and a potential wall  $l: \mathcal{G}_n^d \rightarrow \mathbb{R}$ , we say that  $v$  is a *local maximum* of  $l$  in  $\Gamma$  if for any neighbor  $u$  of  $v$ , such that  $u \in \Gamma$ , we have  $l(v) \geq l(u)$ . We say that  $v$  is a strong local maximum if the inequality is strict, namely  $l(v) > l(u)$ . When the subset  $\Gamma$  is clear from the context, we might omit it and briefly say that  $v$  is a local maximum of  $l$ .

What we want is that if a bit contains both  $v, gv$  then there will be a check that contains both  $v, gv$ , that is true if  $d' > 1$ , then it means that we can erase from the bit a vertex which it is not  $gv$ .

**Claim 5.2.** Let  $z$  be an assignment, and let  $\partial z$  be the set of vertices adjacent to unsatisfied checks induced by  $z$ . Let  $v$  be a vertex in  $\partial z$ , and let  $g$  be a coordinate in  $S$  such that  $v$  is a local maximum of  $L_g$ . Then  $gv \notin \partial z$ .

*Proof.* Suppose  $gv \in \partial z$ , then

$$L_g(gv) = (gv)_g = 1 + v_g = 1 + L_g(v)$$

In contradiction to  $v$  being a local maximum of  $L_g$  in  $\partial z$ . □

**Claim 5.3.** Let  $z$  be an assignment, and let  $\partial z$  be the set of vertices adjacent to unsatisfied checks induced by  $z$ . Let  $v$  be a vertex in  $\partial z$ , and let  $g$  be a coordinate in  $S$  such that  $v$  is a local maximum of  $L_g$ . Then for any check associated with a  $d' - 1$  face rooted at  $v$  that contains  $gv$ , it is satisfied.

*Proof.* Assume, through contradiction, the existence of an unsatisfied check associated with a  $d' - 1$  positive face rooted at  $v$ , which contains  $gv$ . This implies  $gv \in \partial z$ , contradicting Claim 5.2.  $\square$

**Definition 5.4** (Small Code at Vertex  $v$ ). For a vertex  $v$ , we define the  $X$ -small code at vertex  $v$  to be the binary assignments over the bits ( $d'$  faces) rooted at  $v$  that satisfy  $\mathcal{C}_X$ 's restrictions rooted at  $v$  ( $d' - 1$  faces). Similarly, we define the  $Z$ -small code at vertex  $v$  to be the binary assignments over the bits ( $d'$  faces) rooted at  $v$  that satisfy  $\mathcal{C}_Z$ 's restrictions rooted at  $v$  ( $d' + 1$  faces). We denote by  $\mathcal{C}_v^-$  and  $\mathcal{C}_v^+$  the  $X$ -small code at  $v$  and  $Z$ -small code at  $v$ .

**Definition 5.5.** *Should rewrite it better.* We define  $\text{Stab}_v^-$  and  $\text{Stab}_v^+$  as the following linear codes:

$$\text{Stab}_v^- := \left\{ z : \partial z = 0 \text{ and } z \in \text{Span} \left( \left\{ \theta_{\mathbf{v}, S'/\mathbf{g}}, \theta_{\mathbf{g}\mathbf{v}, S'/\mathbf{g}} \text{ for } S' \subset S, |S'| = d' + 1, g \in S' \right\} \right) \right\}$$

$$\text{Stab}_v^+ := \left\{ z : \partial^* z = 0 \text{ and } z \in \text{Span} \left( \left\{ \theta_{\mathbf{v}, S'/\mathbf{g}}, \theta_{\mathbf{g}\mathbf{v}, S'/\mathbf{g}} \text{ for } S' \subset S, |S'| = d' + 1, g \in S' \right\} \right) \right\}$$

**[COMMENT]** Regarding the beneath, The number of restriction in  $\partial^+$  case was counted in negligently. In addition, we should show that for the  $\partial^-$  case, restrictions of the form  $\theta_{\mathbf{g}_2\mathbf{g}_1\mathbf{v}, \mathbf{S}}$  are degenerate. In the bottom line it's look like:

$$\partial^+ \theta_{\mathbf{g}_2\mathbf{g}_1\mathbf{v}, \mathbf{S}} = \partial^+ \theta_{\mathbf{g}_1\mathbf{v}, (\mathbf{S}/\mathbf{g}_3) \cup \mathbf{g}_2} + \partial^+ \theta_{\mathbf{g}_2\mathbf{v}, (\mathbf{S}/\mathbf{g}_3) \cup \mathbf{g}_1}$$

Note that since we talk on mapping checks  $\rightarrow$  bits, we use the  $\partial^+$ . Eventually, we need to use the poicare duality  $H^{d'} \rightarrow H_{d-d'}$ , the duality also should be used to define the SWEEP rule to  $\mathcal{C}_Z$ . I didn't successes to do it other way.

**Claim 5.6.** Let  $v$  be a vertex. For any  $z \in \mathcal{C}_v^\pm$ , there is a stabilizer  $w \in \text{Stab}_v^\pm$  such that  $z = w|_v$ ; namely,  $z$  and  $w$  agree on the faces rooted at  $v$ .

*Proof.* We will show that codes obtained from  $\text{Stab}_v^\pm$  by forcing the faces rooted at  $v$  to  $z|_v$  have non-zero dimensions. Then it follows that any code word of  $\mathcal{C}_v^\pm$  could be extend to codewords in  $\text{Stab}_v^\pm$ . The number of bits in both  $\text{Stab}_v^\pm$  is:

$$\binom{d}{d'} + d \binom{d-1}{d'}$$

The numbers of checks for  $\text{Stab}_\pm$  are:

$$\binom{d}{d' \pm 1} + d \binom{d-1}{d' \pm 1}$$

The restriction to  $z|_v$  sets the bits rooted at  $v$  and satisfies the checks that are rooted at  $v$ , so the checks of the form  $\theta_{v,S'}$  for  $|S'| = d' \pm 1$  are degenerate, and the dimension that remains is:

$$\dim \geq d \binom{d-1}{d'} - d \binom{d-1}{d' \pm 1}$$

In the case that  $d' = (d-1)/2$ , then for any  $x \in [d-1]$  we have  $\binom{d-1}{d'} > \binom{d-1}{x}$  therefore:

$$\begin{aligned} \dim &\geq d \binom{d-1}{d/2} - d \binom{d-1}{d/2 \pm 1} \\ &> d \end{aligned}$$

So there is at least  $2^d$  stabilizers  $w \in \text{Stab}_v$  that agree with  $z$ . □

**Claim 5.7.** Let  $z$  be a finite assignment. Suppose that

$$\max \{L_{d+1}(u) : u \in z\} > \max \{L_{d+1}(u) : u \in \partial^\pm z\}$$

Then there is  $\tilde{z} \sim_{\mathcal{C}_X} z$  such that:

$$\max \{L_{d+1}(u) : u \in \tilde{z}\} \leq \max \{L_{d+1}(u) : u \in z\} - 1$$

*Proof.* Let  $v_1, v_2, \dots, v_k$  be the vertices in  $z$  that achieve the maximum for  $L_{d+1}$ , namely  $v_1, v_2, \dots, v_k = \arg \max_{v \in z} L_{d+1}(v)$ , and assume that  $v_i \notin \partial^\pm z$  for any  $i \in [k]$ .

Consider the  $i$ th vertex  $v_i$ . By definition, since  $v_i \notin \partial^\pm z$ , we have  $\partial^\pm z|_{v_i} = 0$ , and since  $v_i$  is a maximum point of  $L_{d+1}$  in  $z$ , it is also a local maximum, and therefore  $z|_{v_i+} = z|_{v_i}$ . Combining the two, we get that  $z|_{v_i}$  is a codeword of the small code at  $v_i$ , namely  $z|_{v_i} \in \mathcal{C}_{v_i}^\pm$ . Thus, by Claim 5.6, there is a stabilizer  $w_i \in \text{Stab}_{v_i}^\pm$  such that  $w_i$  and  $z$  agree on the positive faces of  $v_i$ , namely  $w_i|_{v_i} = z|_{v_i}$ .

Set  $\tilde{z} \leftarrow z + \sum_i w_i$ . Since  $\sum_i w_i$  is a stabilizer, we have  $\tilde{z} \sim_{\mathcal{C}_X} z$ . In addition, for any  $j \neq i$ , it follows that  $v_j \notin w_i$ . Otherwise, there exists  $S' \subset S$  such that  $v_j \in \theta_{v_i, S'}$ , and therefore  $L_{d+1}(v_j) > L_{d+1}(v_i)$ , in contradiction to  $v_i$  and  $v_j$  being both maximum points for  $L_{d+1}$  in  $\partial^\pm z$ . Thus, we have:

$$\tilde{z}|_{v_i} = (z + \sum_i w_i)|_{v_i} = z|_{v_i} + (z + w_i)|_{v_i} = 0$$

Namely, for any  $i$ , the  $i$ th vertex  $v_i$  is not in  $\tilde{z}$ . Hence, the vertex domain of  $\tilde{z}$  is contained in the union of the vertices of  $z$  with the vertices of  $w_1, w_2, \dots, w_k$ , substituting  $v_1, v_2, \dots, v_k$ . Overall, we get:

$$\begin{aligned} \max \{L_{d+1}(u) : u \in \tilde{z}\} &\leq \max \{L_{d+1}(u) : u \in z \bigcup_i w_i / \bigcup_i v_i\} \\ &\leq \max \{L_{d+1}(u) : u \in z\} - 1 \end{aligned}$$

□



**Claim 5.8.** Let  $z$  be a finite assignment. Then there is  $\tilde{z} \sim_{C_X} z$  such that

$$\max \{L_{d+1}(u) : u \in \tilde{z}\} = \max \{L_{d+1}(u) : u \in \partial^\pm z\}$$

*Proof.* Let  $z$  be a finite assignment, and let  $k$  be an integer such that  $k \geq 1$  and

$$\max \{L_{d+1}(u) : u \in z\} = \max \{L_{d+1}(u) : u \in \partial^\pm z\} + k$$

For an integer  $l \leq k$ , consider the assignments  $\tilde{z}_0, \tilde{z}_1, \dots, \tilde{z}_l$  defined as follow:

- The first element,  $\tilde{z}_0 = z$ .
- If the  $i - 1$ th element satisfies

$$(\star) \quad \max \{L_{d+1}(u) : u \in \tilde{z}_{i-1}\} > \max \{L_{d+1}(u) : u \in \partial^\pm \tilde{z}_{i-1}\}$$

Then define the  $i$ th element,  $\tilde{z}_i$ , to be the result of applying the construction, given in Claim 5.7, on  $\tilde{z}_{i-1}$ , namely  $\partial \tilde{z}_{i+1} = \partial \tilde{z}_i$  and

$$\max \{L_{d+1}(u) : u \in \tilde{z}_i\} \leq \max \{L_{d+1}(u) : u \in \tilde{z}_{i-1}\} - 1$$

If the  $i - 1$ th element doesn't satisfies  $\star$ , then set  $l = i - 1$ .

- If the  $k$ th element was defined, set  $l = k$ .

Since  $\sim_{C_X}$  is a transitive relation, we have that for any  $i \in [l]$ ,  $z \sim_{C_X} \tilde{z}_i$ . If  $l \neq k$ , then there exists  $\tilde{z}_i$  such that  $\tilde{z}_i \sim_{C_X} z$  and  $\tilde{z}_i$  satisfies  $\star$ , namely by setting  $\tilde{z} = \tilde{z}_i$ , we get the desired. So, it is left to handle the case where  $l = k$ . In that case, set  $\tilde{z} = \tilde{z}_k$ . By Claim 5.7:

$$\max \{L_{d+1}(u) : u \in \tilde{z}\} \leq \max \{L_{d+1}(u) : u \in \partial^\pm z\}$$

Yet, since  $\tilde{z} \sim_{C_X} z$ , it follows that  $\partial \tilde{z} = \partial z$ , So:

$$\max \{L_{d+1}(u) : u \in \partial^\pm z\} = \max \{L_{d+1}(u) : u \in \partial^\pm \tilde{z}\} \leq \max \{L_{d+1}(u) : u \in z\}$$

Therefore, we have the equality:

$$\max \{L_{d+1}(u) : u \in \tilde{z}\} = \max \{L_{d+1}(u) : u \in \partial^\pm z\}$$

□

**Claim 5.9.** Let  $z$  be a finite assignment and let  $\xi$  be the result of the application of the Sweep rule on it. Then:

1. For any  $i \in [d]$ :  $\max \{L_i(v) : v \in \partial z\} = \max \{L_i(v) : v \in \partial \xi\}$
2. And for  $i = d + 1$  :  $\max \{L_{d+1}(v) : v \in \partial z\} > \max \{L_{d+1}(v) : v \in \partial \xi\} + 1$

*Proof.* First, since the Sweep rule is invariant to addition by stabilizers, for any  $\tilde{z}$  such that  $\tilde{z} \sim_{C_X} z$ , the correction applied by the Sweep rule on  $\tilde{z}$  is the same as the application on  $z$ . Denote by  $\phi$  the correction, by  $\tilde{\xi}$  the result of applying the Sweep rule on  $\tilde{z}$ , and by  $w$  the stabilizer which is the difference between  $z$  and  $\tilde{z}$ , namely:

$$\begin{aligned}\xi &= z + \phi \\ \tilde{\xi} &= \tilde{z} + \phi \\ \tilde{z} &= z + w\end{aligned}$$

The  $\partial^\pm$  boundary of  $\tilde{\xi}$  equals:

$$\partial\tilde{\xi} = \partial(\tilde{z} + \phi) = \partial(z + \phi) = \partial\xi$$

So, in total, we have that for any  $\tilde{z}$  such that  $\tilde{z} \sim_{C_X} z$ , it holds that  $\partial\tilde{z} = \partial z$  and  $\partial\tilde{\xi} = \partial\xi$ . Therefore, showing that there exists  $\tilde{z} \sim_{C_X} z$  for which the claim holds proves the claim for  $z$ . By Claim 5.8, there is  $\tilde{z} \sim_{C_X} z$  such that:

$$\max \{L_{d+1}(u) : u \in \tilde{z}\} = \max \{L_{d+1}(u) : u \in \partial^\pm \tilde{z}\}$$

For such  $\tilde{z}$ , any maximum of  $L_{d+1}$  in  $\partial\tilde{z}$ , is also a maximum in  $\tilde{z}$ . Denote by  $v_1, \dots, v_k$  the maximum points in  $\partial\tilde{z}$ . Since they are maximum points in  $\tilde{z}$ , it holds that for each  $i \in [k]$ ,  $\tilde{z}|_{v_i} = \tilde{z}|_{v_i+}$ , and hence  $\partial\tilde{z}|_{v_i} = \partial\tilde{z}|_{v_i+}$ . Since  $\tilde{z}|_{v_i+}$  is an assignment rooted at  $v_i$ , then  $v_i$  can suggest  $\tilde{z}|_{v_i+}$ , namely the condition in Sweep rule is satisfied for vertex  $v_i$ . Thus:

$$\partial\tilde{\xi}|_{v_i} = \partial\tilde{z}|_{v_i} + \partial\phi|_{v_i} = 0$$

Namely, for any  $i$ , the  $i$ th vertex  $v_i$  is not in  $\tilde{\xi}$ . For any vertex  $v$  denote by  $\phi_v$  the correction made by the vertex  $v$ , namely  $\phi = \sum_{v \in \partial\tilde{z}} \phi_v$ . Hence, the vertex domain of  $\tilde{\xi}$  is contained in the union of the vertices of  $\tilde{z}$  with the vertices of  $\cup_{v \in V} \phi_v$ , substituting maximum points  $v_1, v_2, \dots, v_k$ . Overall, we get:

$$\begin{aligned}\max \{L_{d+1}(u) : u \in \partial\tilde{\xi}\} &\leq \max \{L_{d+1}(u) : u \in \partial\tilde{z} \cup \bigcup_{v \in \partial\tilde{z}} \phi_v / \bigcup_i v_i\} \\ &\leq \max \{L_{d+1}(u) : u \in \partial\tilde{z}\} - 1\end{aligned}$$

Let  $v$  be a vertex that is a maximum of  $L_g$  in  $\partial\tilde{z}$ , maximum points of  $L_g$  do not see syndrome at the  $g$  direction. If and:

$$\phi_v = \sum \theta_{\mathbf{v}, \mathbf{S}'_i}$$

By Claim 5.3  $\partial\tilde{z}|_{g^v} = 0$ , Hence:

$$\begin{aligned}\partial\phi_v|_{g^v} &= \left( \partial \sum \theta_{\mathbf{v}, \mathbf{S}'_i} \right) |_{g^v} \\ &= \left( \sum_{\mathbf{S}'_i} \sum_{g' \in \mathbf{S}'_i} \theta_{\mathbf{v}, \mathbf{S}'_i / g'} + \theta_{g' \mathbf{v}, \mathbf{S}'_i / g'} \right) |_{g^v} \\ &= 0\end{aligned}$$

Thus if  $g \in \cup S'_i$ , Then:

$$\theta_{\mathbf{g}\mathbf{v}, \mathbf{S}'_i/\mathbf{g}} \in \partial\phi_v|_{gv}$$

So for  $\theta_{\mathbf{g}\mathbf{v}, \mathbf{S}'_i/\mathbf{g}} = 0$ , it must appear at least twice in the sum, yet the left terms in the sum cannot contain it (they differ by the root vertex), and if there is a right term  $\theta_{\mathbf{g}\mathbf{v}, \mathbf{S}'_j/\mathbf{g}} = \theta_{\mathbf{g}\mathbf{v}, \mathbf{S}'_i/\mathbf{g}}$ , then it implies that  $S_j = S_i$ . Which is a contradiction, therefore:  $g \notin \cup S'_i$  and  $gv \notin \phi_v$ .

$$\begin{aligned} \max \left\{ L_g(u) : u \in \partial\tilde{\xi} \right\} &\leq \max \left\{ L_g(u) : u \in \partial\tilde{z} \bigcup_{v \in \partial\tilde{z}} \phi_v / \bigcup_i v_i \right\} \\ &\leq \max \left\{ L_g(u) : u \in \partial\tilde{z} \right\} + 1 - 1 \end{aligned}$$

□

## 6 Toric Encoded Circuits

[COMMENT] Choose one of the following definitions for the base graph  $G_o$ . Pitori Berman and Peter Gacs chose the second, namely the projected version, yet if I had to guess, I would say that the first gives an advantage. Furthermore, We might want to fold computation stages, such that we have GATE  $\rightarrow$  NOISE  $\rightarrow$  DECODING, and after each such cycle freeze the state of the graphs. Finally, we might want that the gates will be SIZE-invariant, like rotating the Torus.

**Definition 6.1** (Troic Encoded Circuit). Let  $C$  be a quantum circuit. We say that  $C$  is *Toric encoded* if the following holds:

1. There is a Toric code  $\mathcal{C}$  such that the qubits of  $\mathcal{C}$  can be partitioned into registers  $R_1, \dots, R_k$ , and each of the registers is encoded by  $\mathcal{C}$ .
2. The gates in  $C$  with even time coordinates are sweep decoding gates.

For a set of faults  $\mathcal{E}$  the gates in  $C[\mathcal{E}]$  can be decomposed to cycles, of the following form:

$$C = D_l P_l \Lambda_l, \dots, D_1 P_1 \Lambda_1$$

**Definition 6.2** (Base graph  $G_o$ ). Let  $C$  be a quantum circuit, at depth  $d$ , with registers  $R_1, \dots, R_k$ , all of which are encoded by Toric codes with the same parameters (i.e., dimension, length, etc.). Let's denote by  $\mathcal{G}_1, \dots, \mathcal{G}_k$  the Toric complexes that induce the codes. We define the base graph of  $C$ , denoted by  $G_o = (V_o, E_o)$  as follows:

1. The vertices  $V_o$  are tuples, where the first coordinates correspond to a vertex in the union over the vertices in  $\mathcal{G}_1, \dots, \mathcal{G}_k$ , and the second coordinate is associated with a time. Namely:

$$V_o := \left\{ (v, t) : t \in [d], v \in \bigcup_i \mathcal{G}_i \right\}$$

2. The edges  $E_o$  are derived from the original edges in  $\mathcal{G}_1, \dots, \mathcal{G}_k$ , with the addition that any two vertices in preceding times  $t$  and  $t+1$  that support qubits (faces) such that  $C$  acts non-trivially on those qubits at time  $t$  are also connected by an edge. Namely:

$$E_o := \{ \{(v, t), (u, t')\} : \text{if} \\ \text{either there exists } i \text{ such } \{v, u\} \in \mathcal{G}_i \text{ and } t' = t, t+1 \\ \text{or there exists } u_i = (g_i, l_i) \in C \text{ such } v, u \in l_{i,2} \text{ and } t = l_{i,1}, t' = l_{i,1} + 1 \}$$

**[COMMENT]** Here we should add that a vertex  $v$  belongs to location  $l$  if there is a qubit  $f$  that  $v$  belongs to its associated face.

**Definition 6.3** (Time Graph of  $C(\mathcal{E})$ ). For a quantum circuit  $C$  and set of faults  $\mathcal{E}$ , let us denote the base graph of  $C[\mathcal{E}]$  by  $G_o$ . For each time  $t$ , denote the deviation induced by  $\mathcal{E}$  at time  $t$  by  $\mathcal{D}_t$ , so  $\partial\mathcal{D}_t$  is the syndrome that the qubits in the deviation admit. We define the history graph  $\mathcal{H} = (V_{\mathcal{H}}, E_{\mathcal{H}})$  as follows:

1. The vertices  $V_{\mathcal{H}}$  are taken to be the vertices of  $G_o$ , namely tuples  $(v, t) \in V_o$ , that satisfy that  $v$  belongs to a face associated with a check, which, at time  $t$ , belongs to the  $\partial\mathcal{D}_t$ . Namely:

$$\{v_t : v_t = (v, t) \in V_o \text{ and there exists a check } f \in C \text{ such } v \in f \in \partial\mathcal{D}_t\}$$

2. The edges  $E_{\mathcal{H}}$  is partitioned into two types  $A_{\mathcal{H}}$  and  $F_{\mathcal{H}}$ , where  $A_{\mathcal{H}}$  are the edges in  $E_o$  restricted to  $V_{\mathcal{H}}$  and  $F_{\mathcal{H}}$  are defined to be the . Namely:

$$A_{\mathcal{H}} = \{e = \{u, v\} \in E_o : u, v \in V_{\mathcal{H}}\} \\ F_{\mathcal{H}} = \{e = \{u, v\} : \text{exists } w \in V_{\mathcal{H}} \text{ such } \{v, w\}, \{u, w\} \in A_{\mathcal{H}}\}$$

By definition of the construction,  $G_{\mathcal{H}}$  is a time-graph.

**Definition 6.4** (**Size()**-Function). Let  $S$  be a subset of vertices, we define the *size* of  $S$  to be:

$$\mathbf{Size}(S) := \sum_i^{d+1} \max_{v \in S} L_i(v)$$

**Claim 6.5.** Let  $\mathbf{S} = \{S_1, S_2, \dots, S_k\}$  where  $S_i \subset \mathbb{R}^d$ , introduce an edge between  $S_i$  and  $S_j$  iff  $S_i \cap S_j \neq \emptyset$ . Assume that  $\mathbf{S}$  is connected. Then  $\mathbf{Size}(\cup_{i=1}^k S_i) \leq \sum_{i=1}^k \mathbf{Size}(S_i)$ .

*Proof.* It suffices to prove that if  $R_1 \cap R_2 \neq \emptyset$ , then  $\mathbf{Size}(R_1 \cup R_2) \leq \mathbf{Size}(R_1) + \mathbf{Size}(R_2)$ . Let  $a_{i,j} = \max_{v \in R_j} L_i(v)$ . Then for any  $w \in R_1 \cap R_2$  we have:

$$\begin{aligned} \mathbf{Size}(R_1 \cup R_2) &= \sum_i^{d+1} \max a_{i,1}, a_{i,2} = \sum_i^{d+1} a_{i,1} + a_{i,2} - \min a_{i,1}, a_{i,2} \\ &\leq \sum_i^{d+1} a_{i,1} + a_{i,2} - L_i(w) = \mathbf{Size}(R_1) + \mathbf{Size}(R_2) - \mathbf{Size}(w) \\ &= \mathbf{Size}(R_1) + \mathbf{Size}(R_2) \end{aligned}$$

□

**Definition 6.6** (Size Invariant Action). [\[COMMENT\]](#) Here we classify both transversal gates, and rotation on the Toric. An action  $g$  is said size invariant action if, for any assignment  $z$ , it holds that  $\mathbf{Size}(gz) = \mathbf{Size}(z)$ .

**Claim 6.7.** The sweep rule is Size  $\xi$ -shrinking.

*Proof.* □

[\[COMMENT\]](#) Eventually, we might not use the projected graph, and then we will split to history of the slices lemma for 3 parts. The first is when time =  $3t$ , in that we apply an invariant action so.

## 7 Space-Time Graph

**Definition 7.1.** Consider a graph  $G = (V, A, F)$ , where  $V$  is a set of vertices, and  $A, F$  are sets of undirected edges. For  $v \in V$ , denote by  $\text{pre}(v) = \{u \in V : \{u, v\} \in A\}$ , we extend the notion for subsets of vertices  $X \subset V$  by  $\text{pre}(X) = \bigcup_{v \in X} \text{pre}(v)$ . Let  $T: V \rightarrow \mathbb{N}$ , We say that  $(G, T)$  is a time graph if and only if:

1.  $\{x, y\} \in A \Rightarrow T(y) = T(x) \pm 1$
2. For  $x, y \in V$   $\{x, y\} \in F$  if and only if there exists  $z \in V$  such both  $\{x, z\}, \{y, z\} \in A$ .

We will call to  $T$  time-function, to  $A$  arrows, and to  $F$  forks. Observes that from the second requirement it follows that forks can connect only vertices that have the same time value (set by  $T$ ).

**Definition 7.2** (History, slice.). For a time graph  $(G, T)$  and a vertex  $u \in V$ , let  $G_u$  be the subgraph of  $(V, A)$  induced by  $\{v \in V : T(v) \leq T(u)\}$ . We will denote by  $H_u$  the 'History of  $u$ ', which is the set of vertices that are connected to  $u$  in  $G_u$ . A slice is an equivalence class of the relation  $u \equiv v$  if and only if  $H_u = H_v$ . Notice that the time function has to be constant on a slice, to see this consider vertices  $x, y$  with  $T(x) > T(y)$ , thus  $x \notin G_y$  and therefore  $x \in H_x/H_y$ . So they are belong to different classes. Thus, we can extend the time function and the history to slices and write for a slice  $K$ ,  $T(K)$  and  $H_K$ .

**Claim 7.3.** Consider slices  $K_1, K_2$  such that  $T(K_1) = T(K_2)$ , then either  $K_1 = K_2$  or  $H_{K_1}, H_{K_2}$  are disjoint.

*Proof.* Assume a non-empty intersection, so there exists  $z \in H_{K_1} \cap H_{K_2}$  such that for any  $x \in K_1$  and  $y \in K_2$ , there exist arrow-paths  $z \rightarrow x$  and  $z \rightarrow y$ . Thus, for any  $w \in H_{K_1}$ , one can concatenate a path  $w \rightarrow x \rightarrow z \rightarrow y$  and conclude that  $y \in H_{K_2}$ . □

**Definition 7.4** (The graph of slice). Let  $K$  be a slice. The graph of  $K$ , denoted by  $G_K = (V_K, E_K)$ , is the graph whose vertices are slices  $R \subset H_K$  with  $T(R) = T(K) - 1$ . Two vertices are connected by an edge if and only if there is a fork whose ends belong to the slices associated with them. Namely,  $S, R \in V_K$  are connected in  $G_K$  if and only if there is  $\{s, r\} = f \in F$  such that  $s \in S$  and  $r \in R$ .

The connection property is crucial, In our construction, there are not neutral forks, and we will have to find way to overcome it, one way is to take  $\arg \min_{u \in \sigma_1, v \in \sigma_2} d(u, v)$  and connect the by fork. More thought should be invested here.

**Claim 7.5.**  $G_K$  is connected.

*Proof.* Let  $R, S \in V_K$  and consider  $r, s \in V$  such that  $r \in R, s \in S$ . Recall that  $R, S \subset H_K$  and therefore  $r, s \in H_K$ , Thus, by definition of the 'History' there exists a path, passing trough arrows only, from  $r$  to  $s$  in  $H_K$ . Consider such a path that is minimal in the number of points  $x$  belong to it, with  $T(x) = T(K)$ . By induction on the number  $k$  of such points we prove that  $R$  and  $S$  are connected in  $G_K$ .

1. Base.  $k = 0$ , In this case, the path from  $r$  to  $s$  lies within  $G_r (= G_s)$ . Hence  $H_r = H_s$ , So  $R = S$ .
2. Induction step. There exists  $y$  on the path from  $r$  to  $s$  such that  $T(y) = T(K)$ . Let  $x$  be the point which precedes  $y$  on the path, and  $z$  the point that follows  $y$ . Since the values of  $T$  at the two ends of an arrow differ by 1, and since  $y$  gets the maximal  $T$  value in  $H_K$  it follows that:
  - (a)  $\{x, z\} \subset \text{pre}(y) \Rightarrow \{x, z\} \in F$ .
  - (b) The slices of  $x$  and  $z$ , say  $K_x$  and  $K_z$  has time value  $T(K_x) = T(K_z) = T(K) - 1$ , Namely they are both members of  $V_K$ .

Since the path connecting  $r, x$  in  $H_K$  has  $k - 1$  vertices with  $T(\cdot) = T(K)$ , then by the induction hypothesis,  $R$  and  $K_x$  are connected in  $G_K$ , and by similar argument so are  $K_z$  and  $S$ . On the other hand,  $\{x, z\} \Rightarrow \{K_x, K_z\} \in E_K$ . Thus,  $R$  and  $S$  are connected in  $G_K$ .

□

**Definition 7.6.** Let  $f: G \hookrightarrow \mathbb{R}^d$  be an embedding that maps each element of  $V \cup A \cup F$  to a point in  $\mathbb{R}^d$ . We say that  $(G, \mathbb{R}^d)$  is a space-time graph if and only if:

- 1.
2. The point set by  $f$  for an edge is the mean of its endpoints' images; that is, for  $e = \{x, y\} \in A \cup F$ , we have  $f(e) = \frac{1}{2} (f(x) + f(y))$ .

**Definition 7.7.** Define  $\text{pre}_i(v) = \arg \max_{u \in \text{pre}(v)} L_i(f(u))$

**Definition 7.8** (Spanned Set). Consider an embedding  $f: \star \hookrightarrow \mathbb{R}^d$ , a subset of  $f$ 's domain  $S \subset \star$ , and  $d + 1$  points  $x_1, \dots, x_{d+1}$ . We say that  $\langle S, x_1, \dots, x_{d+1} \rangle$  is a spanned set of  $S$  if for any  $i \in [d + 1]$  and  $s \in S$ , it holds that  $L(f(s)) \leq L_i(x_i)$  and in addition there are  $v_1, \dots, v_{d+1} \in V$ , not necessary different, that achieve the equalities  $L_i(x_i) = L_i(f(x_i))$ . We name  $x_1, \dots, x_{d+1}$  the poles of  $S$ . Furthermore, we extend the notation to a spanned slice when  $S$  is a slice of some time-graph. Notice that saying  $\langle S, x_1, \dots, x_{d+1} \rangle$  is a spanned set means that  $\mathbf{Size}(S) = \mathbf{Size}(\{x_1, \dots, x_{d+1}\})$ .

Observe that in the case where the subset  $S$  is a slice. Each of these poles can be associated with a vertex of  $G$  that gives the equality. If there are multiple vertices whose images by  $f$  give equality, associate the point with one of them arbitrarily. We will use the notion of applying functions originally defined over vertices to the poles, referring to the application over the vertices associated with them. For example  $\text{pre}(x_i) = \text{pre}(v)$  for the vertex  $v \in S$  which satisfies  $L_i(f(v)) = L_i(x_i)$  and therefore is associated with the pole  $x_i$ . Another example is referring to  $x_i$  as a vertex, so when saying 'For  $x_i \in V_k$ ' or 'For  $\{x_i, u\} \in F$ ', we refer to the associated vertex or the edge connecting it to  $u$ .

**Claim 7.9.** Consider a time graph  $(G, T)$ , an embedding  $f: G \rightarrow \mathbb{R}^d$ , such that.. , Let  $K$  be a slice in  $G$ , and let  $\hat{K} = \langle K, x_1, \dots, x_{d+1} \rangle$  be a spanned slice of  $K$  such that  $K \neq H_K$ . Then for some integer  $k$  there exist spanned slices  $\hat{K}_0 = \langle K_0, \cdot \rangle, \dots, \hat{K}_k = \langle K_k, \cdot \rangle$  and Forks  $F_1, \dots, F_k$  such that:

1. For  $i \in [d+1]$   $\text{pre}_i(x_i)$  belongs to the poles  $\hat{K}_0, \dots, \hat{K}_k$ .
2. Introduce an edge between  $\hat{K}_i$  and  $\hat{K}_j$  if and only if one of the Forks  $F_1, \dots, F_k$  consists of a pole of  $\hat{K}_i$  and a pole of  $\hat{K}_j$ . Then the graph formed from  $\hat{K}_0, \dots, \hat{K}_k$  is connected.
3.  $\sum_{i=0}^k \text{Size}(\hat{K}_i) + \sum_{i=1}^k \text{Size}(F_i) \geq \text{Size}(\hat{K}) + 1$

*Proof.* First Notice that  $K$  doesn't contain leaves in  $(V, A)$ , since a leaf is connected only to itself via arrows, and therefore its 'slice' equals to its 'History'. Thus  $\text{pre}_i(x_i)$  is not empty for  $i = 1, \dots, d+1$ . Next consider the graph  $G_K = (V_K, E_K)$  from definition 7.4. Since  $\{x_i, \text{pre}_i(x_i)\}$  is an arrow  $\text{pre}_i(x_i)$  belongs to a slice in  $V_K$ . for  $i = 1, \dots, d+1$  denote by  $R_i$  the slice in  $V_K$  that contains  $\text{pre}_i(x_i)$ .

By claim 7.5  $G_K$  is connected and therefore there exist a minimal collection  $\{K_0, \dots, K_k\}$  of slices from  $V_K$  which contains  $R_1, R_2, \dots, R_{d+1}$ , and which is connected in  $G_K$ . We pick a set of forks  $\mathbf{F} = \{F_1, \dots, F_k\}$  which realizes a spanning tree for this collection of slices. Later we will identify edges from this spanning tree with the forks from  $\mathbf{F}$ .

For  $i = 1, \dots, d+1$  we set  $\text{pre}_i(v_i)$  to be the  $i$ th pole of  $R_i$ , This assures (1). The set of remaining poles for  $\hat{K}_0, \dots, \hat{K}_k$  will be equal to  $\cup_{i=k}^k F_i$ . This will assure (ii), We will also select poles for  $F_1, \dots, F_k$  to form 'spanned forks'  $\hat{F}_1, \dots, \hat{F}_k$  in such way that:

$$\sum_{i=0}^k \text{Size}(\hat{K}_i) + \sum_{i=1}^k \text{Size}(\hat{F}_i) \geq \sum_{i=1}^{d+1} L_i(\text{pre}_i(v_i)) \geq \text{Size}(\hat{K}) + 1$$

This will assure (3).

Let us denote by  $J_0, \dots, J_{2k}$  the enumerated union  $K_0, \dots, K_k, F_1, \dots, F_k$ . Observes that for any choice of  $(d+1)(2k+1)$  elements  $z_0^1, z_0^2, z_0^3, \dots, z_0^{d+1}, \dots, z_{2k}^{d+1}$  in  $\mathbb{R}^d$ , such that  $z_j^i \in J_j$  it holds that:

$$\sum_{i=0}^k \text{Size}(\hat{K}_i) + \sum_{i=1}^k \text{Size}(\hat{F}_i) \geq \sum_{j,i} L_i(z_j^i) \quad (2)$$

Later, we are going to take the points  $\{z_j^i\}$  from the intersection of  $J_j$ 's. For convenience, let us assume that for the index subset  $X \subset [2k] \cup \{0\}$ , the intersection  $\bigcap_{j \in X} J_j$  returns a single point. If there is more than one, then decide on the point in a way that all the chosen points simultaneously satisfy that for any index subsets  $Y, X$ , such that  $X \cap Y$  and they both introduce a non-trivial intersection  $\bigcap_{j \in Z} J_j : Z \in \{X, Y\}$ , then:

$$\bigcap_{j \in X} J_j = \bigcap_{j \in Y} J_j$$

Now consider a non-trivial maximal intersection  $\bigcap_{j \in X} J_j$ , the intersection is maximal in the sense that for any  $j' \notin X$  it holds that  $\bigcap_{j \in X \cup \{j'\}} J_j = \emptyset$ . We claim that for any  $i$  and for any such  $X$ , there is  $t \in X$  such that  $J_t$  is the successor of  $J_j$  on the path to  $R_i$  for any  $j \in X/\{t\}$ . Assume not. Let  $x, y \in X$ , and denote the paths from  $J_x, J_y$  to the  $i$ th pole by  $\langle J_x, J_{x_2}, \dots, R_i \rangle$ ,  $\langle J_y, J_{y_2}, \dots, R_i \rangle$ , such that  $J_y \neq J_{x_2}$ . If  $y = x_2$ ,  $y_2 = x$ , or  $x_2 = y_2 \in X$ , then pick other  $x, y$  (if there is no such, we get an immediate contradiction). In that case,  $\langle J_x, J_y, J_{y_2}, \dots, R_i \rangle$  is a path from  $J_x$  to  $R_i$  which is different from  $\langle J_x, J_{x_2}, \dots, R_i \rangle$ . Therefore, there are at least two paths from  $J_x$  to the  $i$ th pole, namely, there is a cycle in contradiction to the minimality of  $K_0, \dots, K_k$ .

We will choose the  $z_j^i$  by the following process. Pick a  $z_j^i$  that has not been set yet. If  $J_j = R_i$ , then set  $z_j^i \leftarrow \text{pre}_i(x_i)$  - we call this a type-1 assignment. Otherwise, consider the path from  $J_j$  to  $R_i$  (by connectivity, it has to be such a one, and by minimality, there is exactly one). Let  $J_t$  be the successor on the path, then pick  $z_j^i \in J_j \cap J_t$  - similarly, we call this assignment a type-2 assignment. Since any type-2 assignment is associated with a non-trivial intersection, and any non-trivial intersection induces a type-2 assignment for all  $i$ 's (since for all  $i$ 's, one of the  $J_j$  on its support is a successor of the rest on their path to the  $i$ th pole), we have that:

$$\begin{aligned} \sum_{j,i} L_i(z_j^i) &= \sum_{j,i,z_j^i \in \text{type-1}} L_i(z_j^i) + \sum_{j,i,z_j^i \in \text{type-2}} L_i(z_j^i) \\ &= \sum_i^{d+1} L_i(\text{pre}_i(x_i)) + \sum_{X: \bigcap_{j \in X} J_j \neq \emptyset} (|X| - 1) \sum_i^{d+1} L_i \left( \bigcap_{j \in X} J_j \right) \\ &= \sum_i^{d+1} L_i(\text{pre}_i(x_i)) \end{aligned}$$

Plugging it into eq. (2) gives the desired.  $\square$

**[COMMENT]** In the above, we should formulate the result of the claim as: 'there exists  $K_1, \dots, K_k$  such that:

$$\sum_{i=0}^k \text{Size}(\hat{K}_i) + \sum_{i=1}^k \text{Size}(F_i) \geq \text{Size}(\hat{K}) + \sum_i^{d+1} L_i(\text{pre}_i(x_i))$$

For a decoding phase, we get shrinking. For transversal gate phase and noise phase, we get that the sizes remain the same.



## 8 Explantation of a Spanned Slice

[COMMENT] Here we follow the original proof, but change the triminology,  $m$  will be the number of leaves in the explanation, then in the end we conclude that the number of faults is at least  $m/\Delta$ . So  $|R| \leq f(\frac{m}{\Delta})$ .

**Definition 8.1.** Consider a time graph  $(G, T)$ , an embedding  $f: G \rightarrow \mathbb{R}^d$ , such that.. Let  $\beta = \frac{d+1}{\xi}$  and  $\alpha = \beta \mathbf{Size}(\text{fork})$ . Given sets  $W$  of points,  $R$  of arrows and  $F$  of forks, we define  $U$  as the set of vertices of the graph induced by the edges of  $R \cup F$ . The sets  $W, R$  and  $F$  form an explanation of a spanned slice  $\hat{K} = \langle K, v_1, \dots, v_{d+1} \rangle$  if and only if:

1.  $W \cup \{v_1, \dots, v_{d+1}\} \subset U \subset H_K$  and the graph  $(U, R \cup F)$  is connected.
2. All elements of  $W$  are leaves.
3. For  $m = |W|$  and  $s = \mathbf{Size}(K)$  we have  $|R| \leq \alpha(m - 1) - s$  and  $|F| < m - 1$ .

**Claim 8.2.** Every spanned slice  $K = \langle K, v_1, \dots, v_{d+1} \rangle$  has an explanation.

*Proof.* Let  $h = T(K) - \min_{u \in H_K} T(u)$ . The proof is by induction on  $h$ .

1. Base:  $h = 0$ , namely  $T$  is constant on  $H_K$ , so no arrows connect points of  $H_K$ . Thus,  $H_K$ , as well as  $\mathbf{K}$ , consists only of leaves. In that case,  $U = W = \{u\}$ ,  $m = 1$ ,  $s = 0$ , and  $|R|, |F| = 0$  satisfy the requirement.
2. Induction Step:  $h \neq 0$  implies  $K \neq H_K$ . Thus, by claim 7.9, we know that for some  $k$  there exist spanned slices  $\hat{K}_0, \dots, \hat{K}_k$  and forks  $F_1, \dots, F_k$  that satisfy claim 7.9.

Consider  $i \in 0, \dots, k$ , denote by  $s_i = \mathbf{Size}(\hat{K}_i)$ . By the inductive hypothesis and since  $T(K_i) = T(K) - 1$ , we know that  $\hat{K}_i$  has an explanation formed by a set of leaves  $\mathbf{W}_i$ , a set of arrows  $\mathbf{R}_i$  and a set of forks  $\mathbf{F}_i$ . Let  $m_i = |\mathbf{W}_i|$ , So the sets  $R_i$  and  $F_i$  have at most  $\alpha m_i - s_i$  and  $m_i - 1$  leaves. We define:

- (a)  $\mathbf{W} = \cup_{i=0}^k \mathbf{W}_i$ , Notice that  $\mathbf{W}_i \subset K_i$  and that  $T(K_i) = T(K_j)$  for all  $i, j$ , thus by claim 7.3  $\mathbf{W}$  is a union of disjoint set, Hence  $W$  contains  $m = \sum_{i=0}^k m_i$  leaves.
- (b)  $\mathbf{F} = \{F_1, \dots, F_k\} \cup_{i=0}^k \mathbf{F}_i$  with at most  $k + \sum_{i=0}^k m_i - 1 = m - 1$  elements.
- (c)  $\mathbf{R} = \{\{v_i, \text{pre}_i(v_i)\}\} \cup_{i=0}^k \mathbf{R}_i$  with at most

$$\begin{aligned}
 & d + 1 + \sum_{i=0}^k \alpha(m_i - 1) - \beta s_i \\
 & \leq d + 1 + \alpha m - \alpha(k + 1) - \beta(s + \xi - k \cdot \mathbf{Size}(\text{fork})) \\
 & = \alpha(m - 1) - \beta s + (\alpha + d + 1 - (k + 1)\alpha - \beta\xi + \beta k \cdot \mathbf{Size}(\text{fork}))
 \end{aligned}$$

So its enough to show that for  $k \geq 0$  the term in the closure is negative, We do that by showing that the slop is negative, and then the value of the term at zero is negative:

$$\begin{aligned} -\alpha + \beta \mathbf{Size}(\text{fork}) &\leq 0 \Rightarrow \mathbf{Size}(\text{fork}) \leq \frac{\alpha}{\beta} \\ d + 1 - \beta \xi &\leq 0 \Rightarrow \frac{d+1}{\beta} \leq \xi \end{aligned}$$

And those inequalities hold for  $\beta = \frac{d+1}{\xi}$  and  $\alpha = \beta \mathbf{Size}(\text{fork})$ . Finally, by claim 7.9, the graphs induced by  $\mathbf{R}_i \cup \mathbf{F}_i$  are connected, thus the graph induced by  $\mathbf{R} \cup \mathbf{F}$  is also connected. Therefore,  $\mathbf{W}, \mathbf{R}, \mathbf{F}$  form an explanation of  $K$ .

□

**[COMMENT]** Consider the proof above, when we run two sequences steps, the first is not shrinking, namely  $\xi = 0$ , and the second is shrinking  $\xi \neq 0$ . Each of the proceeding slices  $K_0, \dots, K_k$  has its own  $k_i$  proceeding slices  $K_i^0, \dots, K_i^{k_i}$ . We count the number of arrows in  $K$  by adding the arrows resulting in those two steps  $d + 1 + (k + 1) \cdot (d + 1)$  to the arrows in each of the explanation of the  $K_i^j$  slices. So:

$$\begin{aligned} &(k + 2)(d + 1) + \sum_{i,j=0} \alpha(m_i^j - 1) - \beta s_i^j \\ &\leq (k + 2)(d + 1) + \alpha m - \alpha \sum_{i=0}^k k_i - \beta \left( s + \xi - \left( k + \sum_{i=0}^k k_i - 1 \right) \cdot \mathbf{Size}(\text{fork}) \right) \\ &\leq \alpha(m - 1) - \beta s + \left( \alpha + (k + 2)(d + 1) - \alpha \sum_{i=0}^k k_i - \beta \xi + \beta \left( \sum_{i=0}^k k_i \cdot \mathbf{Size}(\text{fork}) \right) \right) \end{aligned}$$

Now, we want to show that the left term, the linear line, is always negative. For this we look for parameters such when  $k = 0, \sum k_i = 0$  it results with negative value and that the 'slope' of it is negative. For showing negative slope, we use the fact that  $\sum_i k_i \geq k$ .

$$\begin{aligned} k = 0 &\Rightarrow \alpha - \beta \xi \leq 0 \Rightarrow \xi \geq \frac{\alpha}{\beta} \\ &\alpha + (k + 2)(d + 1) - \beta \xi - k(\alpha - \beta \cdot \mathbf{Size}(\text{fork})) \\ &\Rightarrow d + 1 - (\alpha - \beta \cdot \mathbf{Size}(\text{fork})) \leq 0 \\ &\Rightarrow \alpha \geq \beta \mathbf{Size}(\text{fork}) + d + 1 \end{aligned}$$

That is not satisfying the inequalities, one way to try to bypass it is treating to  $O(d)$  SWEEP RULE operations as a constant depth operation, then  $\xi = O(d)$ . For that we need the follow property, for every subset of qubits  $S$ , there is subset of qubits  $S'$  at size at least  $(1 - \gamma)|S|$  such that for  $S$  to be deviated, all the qubits in  $S'$  have to absorb fault. Now suppose that the gate is at depth at most  $\Delta$ , and the fanning is at most  $c_f$ , then we get:

$$\begin{aligned} \Pr[S \in \mathcal{D}] &\leq (c_f^\Delta p)^{|S'|} \leq (c_f^\Delta p)^{(1-\gamma)|S|} \\ &\Rightarrow p \rightarrow (c_f^\Delta p)^{1-\gamma} \end{aligned}$$

A trivial lower bound for  $1 - \gamma$  is  $\frac{1}{c_f^\Delta}$ .

## 9 Counting Subtrees.

**Claim 9.1.** Let  $G = (V, E)$  be a  $\Delta + 1$ -regular graph, and let  $k$  be an integer less than  $|E|$ . Fix a vertex  $r \in V$ . The number of subgraphs rooted at  $r$  with less than  $k$  edges is at most  $\alpha^k$ .

*Proof.* Denote by  $T_\Delta$  and  $T_2$  the infinity  $\Delta$  and binary tress. There is an injection between the subtrees of  $G$ , rooted at  $v$ , with at most  $k$  edges, to the subtrees of  $T_\Delta$  rooted at  $v$ , with at most  $k$  edges. Moreover, there is an injection from the subtrees of  $T_\Delta$  rooted at  $v$ , with at most  $k$  edges to the subtrees of  $T_2$ , rooted at  $v$ , with at most  $k \log \Delta$  edges.

[COMMENT] In the second map, we think on replacing each of the vertex with  $\Delta - 1$ -leaves binary tree, each leaf is associated with a outgoing edge. .

Thus, it's enough to bound the number of of subtrees in  $T_2$  rooted at  $v$ , with at most  $k \log \Delta$  edges, that number is less than the number of binary tress with  $k \log \Delta + 1$  leaves. For exact leaves number we get the  $k \log \Delta$  Catalan number, which its limit is:

$$\sim \frac{4^{k \log \Delta}}{(k \log \Delta)^{\frac{3}{2}} \sqrt{\pi}} \leq \frac{\Delta^{2k}}{(k \log \Delta)^{\frac{3}{2}} \sqrt{\pi}}$$

Therefore we can bound by:

$$\sum_{j=1}^k \frac{\Delta^{2j}}{(j \log \Delta)^{\frac{3}{2}} \sqrt{\pi}} \leq \frac{\Delta^{2k}}{\sqrt{\pi} \log^{\frac{3}{2}} \Delta} \sum_{j=1}^k \frac{1}{j^{\frac{3}{2}}} \leq \zeta\left(\frac{3}{2}\right) \cdot \frac{\Delta^{2k}}{\sqrt{\pi} \log^{\frac{3}{2}} \Delta}$$

□

Now, from here we get an upper bound on the number of subgraphs that connected  $l$  vertices at the top level. We bound it from above by counting combination of rooted subtrees, such the vertices are their root and the overall edges number is summed up to  $k$ . Thus:

$$\leq \zeta\left(\frac{3}{2}\right) \binom{k-1}{l-1} \frac{\Delta^{2k}}{\sqrt{\pi} \log^{\frac{3}{2}} \Delta}$$

## 10 Tree Separator Lemma

For a tree  $T = (V, E)$  and a vertex  $v \in V$ , we call the subtree obtained by taking the descendants of  $v$  the subtree rooted at  $v$ .

**Claim 10.1.** Let  $T = (V, E)$  be a tree with  $m > 3$  leaves and  $\tau$  edges, then it has a subtree with  $m' \in (\frac{1}{3}m, \frac{2}{3}m)$  leaves and at most  $\frac{m'}{m} \tau$  edges.

*Proof.* Order the vertices according to their heights. Denote by  $v \in V$  the lowest vertex for which the subtree rooted at  $v$  has at least  $\frac{1}{3}m$  leaves. Now, denote by  $T_1$  the subtree obtained through the following iterative process over  $v$ 's children:

1. Initialize  $T_1$  as the subtree rooted at  $v$ .
2. For a child  $u$  of  $v$ :
  - (a) Check if erasing from  $T_1$  the subtree rooted at  $u$  doesn't decrease the number of leaves below  $\frac{1}{3}m$ . If so, then we update  $T_1$  to be the result of the erasure.

Clearly, the algorithm is defined such that  $T_1$  has to have at least  $\frac{m}{3}$  leaves. Furthermore, since any child of  $v$  is a root of a subtree with fewer than  $\frac{m}{3}$  leaves (otherwise we get a contradiction to the minimality of  $v$ 's height), we have that if  $T_1$  would have more than  $\frac{2}{3}m$  leaves, then the subtraction of any of its children would yield a tree with at least  $\frac{2}{3}m - \frac{1}{3}m = \frac{1}{3}m$  leaves. Namely,  $v$  has in  $T_1$  a child  $u$  such that the subtraction of the subtree rooted at  $u$  from  $T_1$  results in a subtree with more than  $\frac{1}{3}m$  leaves. But if indeed there were such a child, then the algorithm would erase it, so there is not.

Now let  $T_2 = T/T_1 \cup v$ , namely the tree induced by the subtraction of  $T_1$ 's edges from  $T$ . Observe that since the summation of leaves in  $T_1$  and  $T_2$  is the leaves in  $T$ , and  $T_1$  has no more than  $\frac{2}{3}m$  leaves, and no less than  $\frac{1}{3}m$  leaves, it holds that the number of leaves in  $T_2$  is also in the range  $(\frac{1}{3}m, \frac{2}{3}m)$ . We claim that at least one of the subtrees has fewer than  $\frac{m'}{m}\tau$  edges.

To see this, denote by  $m_1, m_2$  and by  $\tau_1, \tau_2$  the leaves and the edges in  $T_1, T_2$  respectively. Now:

$$\frac{\tau}{m} = \frac{\tau_1 + \tau_2}{m_1 + m_2} = \frac{m_1}{m_1 + m_2} \frac{\tau_1}{m_1} + \frac{m_2}{m_1 + m_2} \frac{\tau_2}{m_2}$$

Namely,  $\frac{\tau}{m}$  is a weighted average of  $\frac{\tau_1}{m_1}$  and  $\frac{\tau_2}{m_2}$ , and therefore it is greater than their minimum. Peak the subtree with the minimal quotient.  $\square$

By repeating recursively on the above construction, we get the following corollary:

**Corollary 10.2.** Let  $T = (V, E)$  be a tree with  $m$  leaves and  $\tau$  edges. Then, for any  $\alpha > 0$ , it has a subtree with  $m' \in (\frac{1}{3}\alpha m, \frac{2}{3}\alpha m)$  leaves and at most  $\frac{m'}{m}\tau$  edges.

**Claim 10.3.** Let  $\gamma, \beta \in \mathbb{R}_{>0}$  and  $n$  be a positive integer. For any tree  $T$  with  $\tau \geq \beta n$  edges and  $m$  leaves such that  $\tau \leq \gamma m$ , there is a subtree  $T'$  of  $T$  with a number of edges  $\tau'$  less than  $\frac{\beta}{2}n$  and a number of leaves greater than  $\frac{\beta}{4\gamma}n$ .

*Proof.* Since the construction of claim 10.1 preserves the ratio of edges to leaves, we have that for a  $T'$  obtained using the construction, the number of edges satisfies  $\tau' \leq \gamma m'$ . So, it's enough to look for the maximal  $\alpha$  such that:

$$\gamma m' \leq \gamma \frac{2}{3}\alpha m \leq \frac{\beta}{2}n \Rightarrow \alpha = \frac{3\beta}{4\gamma} \frac{n}{m}$$

Which gives  $m' \geq \frac{1}{3}\alpha m = \frac{\beta}{4\gamma}n$ .  $\square$

Move this definition to the beginning of the paper.

**Definition 10.4.** We model the qubits on the finite toric with a side length  $n$  by the projection:

$$(x_1, x_2, \dots, x_d) \mapsto (x_1 \bmod n, x_2 \bmod n, \dots, x_d \bmod n)$$

Rewrite when we talk about the first time we have deviation clusters of size  $\in (\alpha d, d)$ . The reason is that having those assumptions means all the droplets in the past shrank; thus, we can use the ratio of edges to time-leaves in the big subgraph.

**Claim 10.5.** Assume that until step  $\tau$ , there was no deviation cluster of size greater than  $d$ . Let  $\hat{K}$  be a spanned slice induced by a deviation of a cluster state at step  $\tau$ , and let  $\omega$  be an explanation for  $\hat{K}$  such that the number of vertices in  $\omega$  is greater than  $\beta n$ . Then there is a subgraph (subtree)  $\omega'$  of  $\omega$  such that  $\omega'$  has fewer than  $\frac{\beta}{2}n$  edges and at least  $\frac{\beta}{4\gamma}n$  leaves.

The connection between the size of the clusters and whether they shrink or not has not been written yet. Moreover, it seems that there might be confusion since two disjoint clusters might be time-connected. It seems that might be a problem.

*Proof.* Let  $\omega$  be an explanation of  $\hat{K}$  with more than  $\beta n$  vertices. Since there is no deviation cluster of size greater than  $d$  until step  $\tau$ , we have that every vertex of  $\omega$  is a spanned slice induced by a boundary of a deviation cluster of size at most  $d$ . Thus,  $\omega$  satisfies the shrink property in Claim 16.6, and by Claim 8.2, the ratio of edges to leaves of  $\omega$  is at most  $\gamma$ . Observe that a spanning tree of  $\omega$  has a ratio of edges to leaves that is at most  $\omega$ 's ratio, and the number of edges equals  $\omega$ 's vertices minus 1. Therefore, by applying Claim 10.3 on  $\omega$ 's spanning tree,  $\omega$  has a subtree with less than  $\frac{\beta}{2}n$  edges and at least  $\frac{\beta}{4\gamma}n$  leaves.  $\square$

State a claim, and notice that in the new version we are not only interested in trees whose root is a single vertex.

*Proof.* Now, since any two connected vertices are at a space-distance of at most 1, it follows that all the leaves in the subtree are at a space-distance of at most  $\beta n$  from each other. Thus, if  $\beta < 1$ , no two of them have the same space-coordinate modulo  $n$ . And therefore the probability to have such subtree is at most  $p^{\frac{\beta}{4\gamma}n}$ .

Because any  $\omega$  with more than  $\beta n$  vertices implies the existence of a subtree with size less than  $\frac{\beta}{2}n$ , it's enough to bound the probability of having such a subtree. Considering that the trees are invariant to shifting by  $n \cdot \mathbf{1}_i$ , we have that it's sufficient to show that there are no such trees which are rooted in the finite cube  $\mathbb{Z}_n^d \times \mathbb{Z}_t$ . Finally, we use claim 9.1 to bound the number of such trees given a fixed root. Overall, we get:

$$\Pr [\text{No } \omega \text{ s with more than } \beta n \text{ vertices}] \leq n^d t \left( \xi^2 p^{\frac{\beta}{4\gamma}} \right)^n$$

$\square$

## 11 Tanner - History Connected Equivalency

Imagine the following process, we apply  $t$  noisy iterations of the SWEEP rule, and then a single ideal (without noise) iteration. Denote by  $\xi$  a deviated configuration (can be either bits or checks), and by  $\xi'$  the result of applying an ideal SWEEP rule on  $\xi$ . Notice that if  $\xi$  is connected in the Tanner graph then  $\xi'$  is connected in the History graph [\[COMMENT\] Require proof, For the classical case enumerate all the options can be done by hand.](#) Thus:

$$\begin{aligned}
\Pr[\xi \text{ connected in Tanner}] &\leq \Pr[\xi' \text{ connected in Time} \mid \text{no noise at the } t+1 \text{ iteration}] \\
&\leq \sum_{m=0}^{\infty} |\{\text{explanation to } \xi' \text{ over } m \text{ leaves restricted to no noise at the final iteration}\}| p^{\frac{m}{\Delta}} \\
&\leq \sum_{m=0}^{\infty} |\{\text{explanation to } \xi' \text{ over } m \text{ leaves}\}| p^{\frac{m}{\Delta}} \\
&\leq \Pr[\xi' \text{ connected in Time} - \text{Upper Bound}] \leq q^{\text{Span}(\xi')} \leq q^{\text{Span}(\xi)-1}
\end{aligned}$$

### 11.1 Another Idea.

Reconsider the construction of the explanation. We could add at the top an imaginary layer in which all the vertices at time  $t$  are connected by arrows to all the vertices (at time  $t+1$ ) that are adjacent to them in the Tanner graph. For the last layer, we might find that the span of a slice expands instead of shrinking. Namely, claim 7.9 for the final slices would change to:

$$\sum_{i=0}^k \text{Size}(\hat{K}_i) + \sum_{i=1}^k \text{Size}(F_i) \geq \text{Size}(\hat{K}) - \xi$$

Thus, following the computation at the induction stage in claim 8.2, the number of arrows is bounded by:

$$\begin{aligned}
&\alpha(m-1) - \beta s + d + 1 + \alpha + \beta \xi \\
&\alpha(m-1) - \beta s + O(1)
\end{aligned}$$

## 12 Intermediate Step Lemma

[\[COMMENT\]](#) The idea of this step is also to bound the case that all the bits are flipped at once, So checks are satisfied (and the argue above is not applied directly).

**Definition 12.1** (Connected Cluster With Span  $s$ ). Let  $G = (B, C, E)$  be a Tanner graph of a code, and let  $f: G \rightarrow \mathbb{R}^{d+1}$  be an embedding of the graph into  $\mathbb{R}^{d+1}$ . For  $s \in \mathbb{R}$ , we say that  $X \subset B$  is a connected cluster of  $G$  with span  $s$  regarding  $f$  if  $X$  is connected in  $G$  (where two bits are connected if there is a check that connects both of them in  $E$ ) and  $f(X) \subset \mathbb{R}^{d+1}$  has  $\text{Size}(f(X)) \leq s$ . When it's clear from the context which  $G$  and  $f$  are, we will call  $X$  in brief a connected cluster with span  $s$ .

**Definition 12.2.** The model, computation, and noise are tossed, and then a correction is made.

**Claim 12.3.** Let  $\xi$  be the event in which, at the **beginning** of time  $t$ , the error contains a connected cluster with a span greater than  $\frac{d}{4}$ , and for  $i \in [t]$ , let  $\sigma_i$  be the event in which, at the **end** of time  $i$ , there is a connected cluster with a span of at least  $\sqrt{d}$  that belongs to the error. Then:

$$\mathbf{Pr} [\xi] \leq t\mathbf{Pr} [\sigma_1] + p^{\frac{1}{8}\sqrt{d}}$$

*Proof.* By the law of total probability it holds:

$$\mathbf{Pr} [\xi] = \mathbf{Pr} \left[ \xi \cap \bigcup_i^t \sigma_i \right] + \mathbf{Pr} \left[ \xi \cap \bigcap_i^t \bar{\sigma}_i \right]$$

First, let us bound from above the probability of the right intersection, In that case, at the **end** of time  $t - 1$ , the error consists of clusters each at span less than  $\sqrt{d}$ . Let  $m$  be the number the clusters and denote them by  $X_1, \dots, X_m$ . Now since the clusters are disjoint, the sum of their **Size**( $X_i$ ) is lower than the total number of the bits, namely:

$$\sum_{i=1}^m \mathbf{Size}(X_i) \leq n \Rightarrow m \leq \frac{n}{\sqrt{d}}$$

$$p \frac{n}{\sqrt{d}} \leq \sqrt{d}$$

, Thus, . So denote by  $\tau$  the number of bits that require to be flipped in order to merge the flipped connected component to a one with a span greater than  $d/4$ , So  $\tau$  has to be at least:

$$\tau \left( \sqrt{d} + 1 \right) \geq \frac{1}{4}d \Rightarrow \tau \geq \frac{1}{8}\sqrt{d}$$

Therefore:

$$\begin{aligned} \mathbf{Pr} [\xi \cap \bigcap_i^t \bar{\sigma}_i] &\leq \sum_{\xi^{(t-1)}} \mathbf{Pr} [\xi | \xi^{(t-1)}] \mathbf{Pr} [\xi^{(t-1)} \cap \bigcap_i^t \bar{\sigma}_i] \\ &\leq \sum_{\xi^{(t-1)}} p^{\frac{1}{8}\sqrt{d}} \mathbf{Pr} [\xi^{(t-1)} \cap \bigcap_i^t \bar{\sigma}_i] \leq p^{\frac{1}{8}\sqrt{d}} \end{aligned}$$

$$\begin{aligned} \mathbf{Pr} [\xi] &= \mathbf{Pr} [\xi \cap \bigcup_i^t \sigma_i] + \mathbf{Pr} [\xi \cap \bigcap_i^t \bar{\sigma}_i] \\ &\leq t\mathbf{Pr} [\sigma_1] + p^{\frac{1}{8}\sqrt{d}} \end{aligned}$$

□

**Definition 12.4.** The atomic events are the sets of space-time locations in which faults occur. A run  $\omega$  is the sequence of states (assignment of all the bits) induced by an atomic event (and the transition function of the machine). We say that a run  $\omega$  closes a non-trivial cycle if it contains a state in which there is a connected cluster  $A$  such that its boundary is not a co-boundary.

Define the bits-cluster graphs  $G = (V, E)$  such that the vertices are connected bit-clusters at different times  $\{A_i^t\}$ , and there is an edge between  $A_i^t$  and  $A_j^{t-1}$  only if  $\Pi_x A_i^t \cap \Pi_x A_j^{t-1} \neq \emptyset$ . Given a state at time  $t - 1$  such that no connected component in  $G$  forms a closed non-trivial circle, consider  $\tilde{A}_i^t$  constructed as follows: perform the application of either noise or Toom's correction in an arbitrary order and stop right before  $\tilde{A}_i^t$  forms a closed non-trivial circle.

**Claim 12.5.** There is  $\phi \in \text{Aut}(\mathbb{T}_n^d)$  for which  $\text{Size}(\phi(\tilde{A}_i^t)) \geq n - 1$ .

**Claim 12.6.** Consider a running in which no non-trivial circle was closed, then there is  $\psi: \mathbb{T}_n^d \rightarrow \mathbb{Z}_{2n}^d$  such:

1. For any slice  $K$ ,  $\psi(K)$  is connected in  $\mathbb{Z}_{2n}^d$ .
2. For any  $t$ , denote by  $S_t$  the state at time  $t$ . Then, for any  $t > 1$ , we have that  $\psi(S_t)$  is the image of the application of the Tooms rule over  $\psi(S_{t-1})$ .

*Proof.* We prove it by induction on time  $t$ .

1. Base case,  $t = 1$ . We will construct  $\psi$  in the following iterative manner. First, let  $\psi$  be the identity. Then, while there are slices that are not connected on  $\mathbb{Z}_{2n}^d$ , do the following: Let  $K$  be a slice that is not connected on  $\mathbb{Z}_{2n}^d$ . Then it can be decomposed into a disjoint union  $K = K_1 \cup K_2$  such that  $K_1$  and  $K_2$  are connected on  $\mathbb{Z}_{2n}^d$ . Assume, without loss of generality, that for a coordinate  $x \in [d]$ , it holds that for any  $a \in K_1$  and  $b \in K_2$ , we have  $b_x - a_x > 1$ .

Observe that there has to be at least one such coordinate; otherwise, it is a contradiction to  $K_1$  and  $K_2$  not being connected in  $\mathbb{Z}_{2n}^d$ . Then define:

$$\psi(z) \leftarrow \begin{cases} \psi(z) & z \cap K_1 = \emptyset \\ z + n\mathbf{1}_x & \text{else} \end{cases}$$

Repeat the above for other coordinates if  $\psi(K)$  is still not connected in  $\mathbb{Z}_{2n}^d$ . The proof that  $\psi(K)$  has to be connected (that the process ends) is omitted.

2. Induction step. We repeat the construction in the base case. To prove the second property, we show that  $\psi$  maps (un)satisfied edges to (un)satisfied edges. [\[COMMENT\] That is wrong.](#)

□

**Claim 12.7.** Denote by  $\omega$  running of the above type, then there is a running  $\omega'$  on  $\mathbb{Z}_\infty^d$  such that  $\omega$  and  $\omega'$  agree.



**Claim 12.8.** Consider a  $t$ -steps running, conditined on the event that no trivial-cycle was closed in the  $t - 1$  first steps, the probability that a non-trivial cycle will be closed at the  $t$ th step is less than:

$$\Pr \left[ \Lambda^t \mid \bigcap_{t' < t} \bar{\Lambda}^{t'} \right] < n^{2d} \sum_{i=n-1}^{\infty} p^i \mathcal{J}_i \leq n^{2d} q^{n-1}$$

*Proof.* Consider the event in which there is a droplet that at time  $t$  closes a non-trivial cycle. Let  $\tilde{A}$  be the subset of that droplet obtained by the process in claim 12.5. By the claim,  $\tilde{A}$  has a boundary  $\sigma$  and two vertices on that boundary  $M = \{u, v\}$  such that  $\text{Size}(M) > n - 1$ , up to isomorphism.

Now, since no slice in the investigation of  $M$  closes a non-trivial cycle then by claim 12.6 .. so by claim 12.7 ..  $\square$

## 13 Configuration up to $C_z^\perp$

## 14 Drafts:

In addition since, in a tree, the number of leavs is at least greater than the maximal degree, then we have that each  $J_j$  intersects with at most  $d + 1$  elements.

$$\begin{aligned} L + \Delta_{\max} + 2(n - L - 1) &\leq \sum_{v \in V} \deg(v) = 2(n - 1) \\ \Rightarrow L &\geq \Delta_{\max} \end{aligned}$$

## 15 Decoder Phase

In the general picture, I would like to use the 4D toric code to encode in any target code. We use the original fault tolerance construction (the concatenation construction) to encode into and decode from the 4D toric code. So, for the decoding phase, we need to have a decoder which, in the ideal setting, runs in less than logarithmic time.

What we know so far:

1. We can assume that two unsatisfied checks are belong to the same history with probability less than  $\sim q^{|u-v|}$ .
2. There is a  $\log^2(n)$  depth algorithm to find the perfect/max matching in a general graph. From first check, the known algorithms to deal with the weight matching are based on LP.
3. Applying Toom's/Sweep rule will require at least linear depth.
4. Having an algorithm that succeeds with constant probability  $\frac{1}{2} + \epsilon$  might be enough (Anyway the qubit at the result of the decoding is going to absorb  $p$ -noise).

## 16 Appendix

[Do something with that.](#) And clearly  $\text{Stab}_v$  is collection of stabilizers since for any  $S' \subset S$  such  $|S'| = d' + 1$  we have:

$$\begin{aligned} \partial(\partial\theta_{\mathbf{v}, S'}) &= \partial \left( \sum_{g \in S'} \theta_{\mathbf{v}, S'/g} + \sum_{g \in S'} \theta_{\mathbf{g}\mathbf{v}, S'/g} \right) \\ &= \sum_{\substack{g', g \in S' \\ g' \neq g}} (\theta_{\mathbf{v}, S'/\{\mathbf{g}, \mathbf{g}'\}} + \theta_{\mathbf{g}'\mathbf{v}, S'/\{\mathbf{g}, \mathbf{g}'\}} + \theta_{\mathbf{g}\mathbf{v}, S'/\{\mathbf{g}, \mathbf{g}'\}} + \theta_{\mathbf{g}'\mathbf{g}\mathbf{v}, S'/\{\mathbf{g}, \mathbf{g}'\}}) \\ &= 0 \end{aligned}$$

Where we used that the term in the closure is invariant to sweeping between  $g$  and  $g'$ , and therefore any term in the sum appears twice.

**Definition 16.1.** Let  $z$  be a connected assignment, and let  $\Gamma$  be the set of vertices adjacent to unsatisfied checks induced by  $z$ . For a generator  $g \in S$ , we say that  $v$  is a  $g$ -pole if  $v \in \Gamma$  and  $gv \notin \Gamma$ . Furthermore, if there is a vertex  $v \in \Gamma$  such that for any  $g \in S$ , we have  $g^{-1}v \notin \Gamma$ , then we say that  $v$  is a local minimum of  $z$ .

Let  $x_1, \dots, x_k \subset \{\pm\}^k$  and let  $p = \prod_i g_i^{x_i} v$  we say that  $p$  is a forward path in  $z$  if it holds that:

1. Any partial product of the first  $j$  elements  $\prod_i^j g_i^{x_i} v$  is a vertex in  $z$ .
2. For any  $g \in S$ , we have that the sign of the sum of the exponents of the powers of the elements in the product associated with  $g$  is positive. Namely:

$$\sum_{i: g_i = g} x_i > 0$$

Similarly, we say that  $p$  is a backward path if the above holds, but the sign in the second condition is either negative or zero. The infimums of  $z$  are defined to be the set of vertices  $U \subset z$  such that for any  $u \in U$  and  $v \in z$ , other vertices in  $z$ , there is a forward path starting at  $u$  and ending at  $v$ .

### 16.1 Alternative Proof For $d' = 2$ .

**Definition 16.2.** We denote by  $\phi^t$  the subsets of bits which are flipped as a result of applying the Toom's rule at time  $t$ , and by  $\phi_v$  the bits which are flipped by vertex  $v$ . We say that  $u$  is added to  $\Gamma^{t+1}$  by  $v$  at time  $t$ , if  $u \in \Gamma^{t+1}$  and  $u \in V(\phi_v)$ . When it clear from the context, we will omit the time, and say that  $u$  is added by  $v$ .

**Claim 16.3.** Let  $v \in \Gamma^t$  for any  $u$  that is added by  $v$ , and for any  $x \in [d+1]$ , we have  $L_x(u) \leq L_x(v) + 1$ .

*Proof.* By definition, any  $u$  that is added by  $v$  is of the form  $v + \sum \alpha_j 1_j$ , where  $\alpha_j \in \{0, 1\}$  and  $\sum \alpha_j \leq 2$ . Thus, for any  $x \in [d]$ :

$$L_x(u) = \langle 1_x, v + \sum \alpha_j 1_j \rangle = \langle 1_x, v \rangle + \alpha_x \leq L_x(v) + 1$$

For  $x = d + 1$  we have:

$$L_x(u) = \langle -\mathbf{1}, v + \sum \alpha_j 1_j \rangle = \langle -\mathbf{1}, v \rangle - \sum_j \alpha_j \leq L_x(v)$$

□

**Claim 16.4.** Let  $v \in \Gamma^t$  such that  $L_{d+1}(v)$  is a (local) maxima in  $\Gamma^t$  then for any  $u$  which is added by  $v$  we have  $L_{d+1}(u) \leq L_{d+1}(v) - 1$ .

*Proof.* We will prove that  $v \notin \Gamma^{t+1}$ , and then by repeating on claim 16.3 where  $\sum \alpha_j \geq 1$ , we get the desired result. Assume, toward contradiction, that  $v \in \Gamma^{t+1}$ , and denote by  $w$  the vertex that adds  $v$ . If  $w \neq v$ , then there is at least one coordinate  $i$  for which  $w_i < v_i$ , thus  $L_{d+1}(w) > L_{d+1}(v)$ , in contradiction to  $L_{d+1}(v)$  being a maximum in  $\Gamma^t$ . In the other case, where  $v$  adds itself, it follows that  $\sigma_v$  can't be decomposed into pairs of unsatisfied checks; namely,  $|\sigma_v|$  is odd. We show by induction that impossible, namely that  $|\sigma_v|$  is always even. The induction is over the size of the plaquettes subset that induce the boundary  $\sigma$ .

1. Base. Consider a boundary that induced by a single plaquette, in that case it obvious that  $|\sigma_v|$  is either zero (where  $v$  is not belong to the plaquette) or 2.
2. Assume for any boundary that induced by at most  $k$  plaquettes, and consider a boundary  $\sigma$  which is induced by  $k + 1$  plaquettes, name this subset  $F$  and observes that  $F$  can be decomposed to a sum of two disjoint subsets  $A$  and  $B$ , each with less than  $k + 1$  plaquettes, Now:

$$\begin{aligned} \sigma_v &= (\partial A)_v \cup (\partial B)_v \\ \Rightarrow |\sigma_v| &= |(\partial A)_v| + |(\partial B)_v| - 2|(\partial A)_v \cap (\partial B)_v| = \text{even} \end{aligned}$$

Where we used the induction assumption, which tells that  $|(\partial A)_v|$  and  $|(\partial B)_v|$  are even numbers.

□

**Claim 16.5.** Let  $x \in [d]$  and let  $v \in \Gamma^t$  such that  $L_x(v)$  is a (local) maxima in  $\Gamma^t$  then for any  $u$  which is added by  $v$  we have  $L_x(u) \leq L_x(v)$ .

*Proof.* Since  $L_x(v)$  is maximal,  $v$  doesn't see edges that goes from it at the positive  $x$ -direction, otherwise denote it by  $e$  and denote by  $w$  the second vertex at the end of  $e$ ,  $w$  has an  $x$ -coordinate which is greater than  $v$ 's  $x$ -coordinate by 1 and in addition  $w \in \Gamma^t$  (since it lays on the support of  $e$ ). Therefore  $L_x(w) > L_x(v)$  in contradiction to  $L_x(v)$  be maximal in  $\Gamma^t$ . Hence, any plaquette that is being flipped by  $v$  lies on

edges associated with  $1_j$  such that  $j \neq x$ . Thus, any vertex  $u$  that is added by  $v$  has the form  $v + \sum_{j \in [d] \setminus \{x\}} \alpha_j 1_j$ , where  $\alpha_j \in \{0, 1\}$  and  $\sum \alpha_j \leq 2$ . So:

$$L_x(u) = \langle 1_x, v + \sum_{j \in [d] \setminus \{x\}} \alpha_j 1_j \rangle = \langle 1_x, v \rangle = L_x(v)$$

□

**Claim 16.6.** Let  $\Gamma^t$  be the vertices in the support of the domain wall  $\sigma^t$ . Denote  $v_i = \arg \max_{v \in \Gamma^t} L_i(v)$ . Then for any  $i \in [d]$ , one can find  $u \in \Gamma^{t-1}$  such that  $u$  adds  $v_i$  and  $L_i(u) \geq L_i(v_i)$ . In addition, one can find  $u_{d+1} \in \Gamma^{t-1}$  such that  $u_{d+1}$  adds  $v_{d+1}$  and  $L_{d+1}(u_{d+1}) = L_{d+1}(v_{d+1}) + 1$ .

*Proof.* Just take  $u_i$  to be any vertex that adds  $v_i$ . By claims claim 16.3, claim 16.4, and claim 16.5, we get the claim. □

[COMMENT]  $V(\phi_v)$  was not defined.  $v \leq w$  also not defined.

**Definition 16.7** ( $\sigma_1$  Precedes  $\sigma_0$ ). We say that  $\sigma_1$  precedes  $\sigma_0$  if, for any  $v \in \sigma_1$ , it holds that  $\phi(v) \subset \sigma_0$ .

**Definition 16.8** (Explanation.). Let  $\sigma_0$  be a boundary at time  $t$ , and let  $\sigma_1, \dots, \sigma_k$  be boundaries at time  $t - 1$ , such that for  $i \in [k]$ , it holds that  $\sigma_i$  precedes  $\sigma_0$ . Then the explanation for  $\sigma_0$  is the subgraph induced by the vertices of  $\sigma_0, \sigma_1, \dots, \sigma_k$ , the edges  $\{u, v\} \in \sigma_i$ , and the arrows  $\{v, u\}$  for  $v \in \cup \sigma_i$  and  $u \in \phi(v)$ .

The number of edges above is not linearly bounded by the leaves. Yet it also seems not minimal. So what we would like to do is to keep the poles, then to connect the islands by fork. So what is it a fork?

**Definition 16.9** (Fork). Let  $u, v$  be such that  $u \in \sigma_1$  and  $v \in \sigma_2 \neq \sigma_1$ . If there exist  $w \in \phi(v)$  and  $z \in \phi(u)$  such that  $\{w, z\} \in \mathcal{F}_1$ , then there is a fork between  $v$  and  $u$ .

*Proof.* Idea: Suppose not, then  $\sigma_0 = \phi(\sigma_1) \cup \phi(\sigma_2)$  is not connected. □

**Claim 16.10.** Assume that  $\sigma_1, \sigma_2 \rightarrow \sigma_0$ , then there is a fork between  $\sigma_1$  and  $\sigma_2$ . So if we keep to extent the history in a recursive manner, then we get a connected graph.

## 17 Garbage

*Proof.*

$$L_{d+1}(u) \leq \min_{v \in \partial z} L_{d+1}(v)$$

Observes that if the *termination-condition* holds then any local maximum of  $L_{d+1}$  in  $\partial z$  is also a local maximum of  $L_{d+1}$  in  $\tilde{z}$  and we finish. Consider the other case,

namely there is at least one vertex  $u \in \tilde{z}/\partial z$  such that  $u$  is a local maximum for  $L_{d+1}$  and

$$L_{d+1}(u) > \min_{v \in \partial z} L_{d+1}(v)$$

In that case, we peak such  $u$ , and looking for stabilizer  $w$  such  $u \notin \tilde{z} + w$  by repeating over the above procedure where  $u$  plays the role of  $v$ . We keep repeating on the procedure until the *termination-condition* holds.

We argue that the process indeed terminates. Consider a vertex  $u$  such that  $u$  is a local maximum of  $L_{d+1}$  in  $\tilde{z}$  and  $u$  is not a local maximum of  $L_{d+1}$  at  $z$ . Put differently,  $u$  becomes a local maximum as a result of adding  $w$  to  $z$ . Thus,  $u$  has to belong to  $w|_v \setminus v$ , which implies that there is a non-empty subset of generators  $I \subset S$  such that  $u = (\prod_{g \in I} g)v$ , and therefore  $L_{d+1}(u) \leq -1 + L_{d+1}(v)$ .

Hence, since  $z$  is finite.

We repeat the process while possible; namely, we stop when all the local maxima of  $L_{d+1}$  are in  $\partial$ .  $\square$

We prove that  $\dim \mathcal{C}_v = \dim \text{Stab}_v$  below, for now assume it is correct. We continue by using the fact that if, for a local maximum  $v$  in  $\partial$ , it holds that  $z|_v$  is rooted at  $v$ , then the decoder will find bits rooted at  $v$  such that their  $\partial$  agrees with  $\partial$ .

To generalize, one might think that it has to complete the local view on  $v$  such that all the checks are satisfied. For  $C_X$ , it sums up to show that the following dimension is positive. Yet the trick fails for  $C_Z$ , where we need to think of a different idea.

$$d \binom{d-1}{d'} - \binom{d}{d'-1} - d \binom{d-1}{d'-1}$$

**Claim 17.1.** Let  $z$  an assignment rooted at vertex  $v$ , such  $\partial z|_{v+} = 0$ . Then there is a stabilizer  $w$ , rooted at  $v$ , such that  $w|_{v+} = z$ .

*Proof.* Denote by  $F$  the faces in  $z$ , and by  $J$  the collection of generators that support  $F$ . We claim that  $|J| \geq d' + 1$ .

Clearly,  $|J| \geq d'$ , since each face is defined by a root vertex and  $d'$  generators. Yet, observe that if  $|J|$  equals exactly  $d'$ , then  $z$  contains a single  $d'$ -face, denote it by  $f$ . Then, any check associated with a  $d' - 1$  face of  $f$  is unsatisfied, in particular the  $d' - 1$  faces that are rooted at  $v$ , in contradiction to  $\partial z|_{v+} = 0$ .

Now, one can choose  $J' \subset J$  of size  $d' + 1$ , and define the stabilizer  $w = \theta_{v,J'}$ .  $\square$

**Claim 17.2.** Suppose  $v \in z$  and for any  $g \in S$ , we have  $g^{-1}v \notin z$ , then there is  $\theta_{v,J} \in \partial z$ .

*Proof.*

$$\partial z = \sum_{S'} \theta_{\mathbf{v}, S'} + ..$$

$\square$

**Claim 17.3.** Let  $z$  be a connected assignment on the  $d'$ -faces at finite size, and let  $\Gamma$  be the set of vertices which are adjacent to unsatisfied checks induced by  $z$ . Let  $v$  be a vertex in  $\mathcal{G}_\infty^d$  such that  $v$  is a strong global maximum of  $L_{d+1}$ , at  $\Gamma$ . Then there is a rooted assignment  $\tilde{z}$  at  $v$  such that  $z_v = \tilde{z}_v$ .

*Proof.* Let  $\Theta = \theta_{v_1, S'_1}, \theta_{v_2, S'_2}, \dots, \theta_{v_k, S'_k}$  be the faces forms  $z$ .

Since  $z$  is finite, it has a maximum for  $L_{d+1}$ . Denote by  $u$  the vertex that gets the maximum, and denote by  $\theta_{u, J_1}, \dots, \theta_{u, J_l}$  the  $d'$ -faces in  $z$  that are rooted at  $u$ . We claim that  $u = v$ .

Suppose not, namely  $u \notin \Gamma$ . Then it means that any  $d' - 1$  face containing  $u$  is satisfied. Observe that  $\sum \theta_{u, J_i} \neq 0$ ; otherwise, it would contradict the non linearity independence of  $\theta_{v_1, S'_1}, \theta_{v_2, S'_2}, \dots, \theta_{v_k, S'_k}$ . So, by Claim 17.2 there exists a  $d' - 1$  face rooted at  $u$ , contained in  $\sum \theta_{u, J_i}$ . Let us denote it by  $\theta_{u, J'}$ .

Since  $\theta_{u, J'}$  is satisfied, it has to be adjacent to a turned bit, namely a  $d'$  face in  $z$ , not in  $\theta_{u, J_1}, \dots, \theta_{u, J_l}$ . Denote that  $d'$  face by  $f$ . Yet, since  $\theta_{u, J'}$  is rooted at  $u$ , it contains  $u$ , and therefore  $f$  contains  $u$ . Now, if  $f$  is rooted at  $u$ , then it is a contradiction that  $\theta_{u, J_1}, \dots, \theta_{u, J_l}$  are all the  $d'$  faces in  $z$  rooted at  $u$ . If  $f$  is not rooted at  $u$ , denote its root by  $w$ , so there is  $g \in S$  such that  $gw = u$ , and  $w \in z$ , in contradiction to  $u$  being maximum for  $L_{d+1}$ .

So, we find that  $v$  is also the maximum of  $L_{d+1}$  in  $z$ . Therefore, we have that  $\tilde{z} = \sum \theta_{u, J_i}$  zeros out  $z_v$ .  $\square$

**Claim 17.4.** Let  $z$  be a connected assignment on the  $d'$ -faces at finite size, and let the collection  $\Theta = \theta_{v_1, S'_1}, \theta_{v_2, S'_2}, \dots, \theta_{v_k, S'_k}$  be the decomposition forms  $z$ . Denote by  $\Gamma$  the set of vertices adjacent to unsatisfied check induced by  $z$ . Then  $\Theta$  is acyclic, if and only if  $z$  has at least one Infimum.

*Proof.* First, suppose that  $z$  has at least one Infimum. Let  $u$  be an infimum in  $z$ . Second, suppose that  $\Theta$  is acyclic.  $\square$

**Claim 17.5.** Suppose that  $d' > 1$  and a vertex  $v \in \mathcal{G}_\infty^d$  is also adjacent to the domain wall, namely  $v \in \Gamma$ . It holds that  $v$  cannot be simultaneously both a (strong) local max of  $L_{d+1}$  and a local min of  $L_{d+1}$ .

*Proof.* Let  $v \in \Gamma$  be a local maximum point of  $L_{d+1}$ , It means that:

1. The vertex  $v$  is adjacent to a  $d'^{-1}$ -face which is unsatisfied.
2. That face is rooted in  $v$ ; otherwise, the root of the face is a vertex in  $\Gamma$  with a higher value of  $L_{d+1}$ , in contradiction to  $v$  being a local maximum.

Thus, there is a subset of generators  $S' \subset S$ , of size  $d' - 1 > 0$ , such that the unsatisfied check set is on the face  $\theta_{v, S'}$ . Given  $d' > 1$ , we have that  $S'$  contains at least one more element besides the empty set, denoted by  $g$ . Now, consider the vertex  $gv$ . Since  $gv$  is adjacent to  $\theta_{v, S'}$ , then  $gv \in \Gamma$ . On the other hand, on the infinite toric  $\mathcal{G}_\infty^d$ , moving along the grid generator decreases  $L_{d+1}$  by one, so  $L_{d+1}(gv) = L_{d+1}(v) - 1$ . In other words,  $gv$  is a neighbor of  $v$  with a lower value of  $L_{d+1}$ , and thus  $v$  cannot be a local minimum of  $L_{d+1}$ .  $\square$

**Remark 17.6.** That claim alone gives a decoder in the clean setting. Write a remark indicating that when  $d' = 1$ , meaning qubits are on the edges (2D toric), the claim is not true.

For  $d = 4, d' = 2$ . For  $i, j \in [d]$ , such that  $i \neq j$ , denote the bit rooted at  $v$  and spanned by generators  $g_i, g_j \in S$ , namely  $\theta_{v, \{g_i, g_j\}}$ , by  $g_{ij} = g_{ji}$ . The restrictions:

$$r_i: \sum_{j \neq i} g_{ij}$$

Thus we get

$$\begin{aligned} \sum_i^n r_i &= \sum_i^n \sum_{j \neq i}^{n+1} g_{ij} \\ &= \sum_i^n \sum_{j < i}^n g_{i,j} + g_{j,i} + \sum_i^n g_{i,n+1} \\ &= r_{n+1} \end{aligned}$$

Thus the dimension of the small code at  $v$  is at least:

$$\binom{d}{2} - (d-1) = 3$$

We have  $\binom{4}{3} = 4$  stabilizers rooted at  $v$  (each for each cube). Yet:

$$\begin{aligned} s_1 &= g_{1,2} + g_{2,3} + g_{1,3} \\ s_2 &= g_{1,2} + g_{2,4} + g_{1,4} \\ s_3 &= g_{1,3} + g_{3,4} + g_{1,4} \\ s_4 &= g_{2,3} + g_{3,4} + g_{2,4} \end{aligned}$$

And clearly

$$s_4 = s_1 + s_2 + s_3$$

**Definition 17.7** (Reduced subset with boundrey at  $v$ ).

**Definition 17.8** (Base graph  $G_o$ ). Let  $C = \{u_i\}_{i=1}^n$  be a quantum circuit, at depth  $d$ , with registers  $R_1, \dots, R_k$ , all of which are encoded by Toric codes with the same parameters (i.e., dimension, length, etc.). Let's denote by  $\mathcal{G}$  the Toric complex that induces the codes, and suppose that the locations of the gates in  $C$  are given in registers' coordinate systems. We define the base graph of  $C$ , denoted by  $G_o = (V_o, E_o)$  as follows:

1. The vertices  $V_o$  are tuples, where the first coordinates correspond to a vertex in  $\mathcal{G}$ , and the second coordinate is associated with a time. Namely:

$$V_o := \bigcup_i V(\mathcal{G}_i) \times [d]$$

2. The edges  $E_o$  are derived from the original edges in  $\mathcal{G}$ , with the addition of edges of the following form: For two vertices at preceding times  $t$  and  $t + 1$ , let's denote them by  $(v_1, t)$  and  $(v_2, t + 1)$ , for which there exist registers  $R'_1, R'_2$ , and faces  $f_1, f_2 \in \mathcal{G}$ , such that  $v_1 \in f_1, v_2 \in f_2$  and  $C$  contains a gate that acts non-trivially on the locations  $(t, f_1, \mathbf{1}_{R'_1})$  and  $(t, f_2, \mathbf{1}_{R'_2})$  at time  $t$ , are also connected by an edge. Namely:

$$\begin{aligned}
E_o := \{ \{ (v, t), (u, t') \} : & \text{ if} \\
& \text{either } \{v, u\} \in \mathcal{G} \text{ and } t' = t, t + 1 \\
& \text{or there exists } u_i = (g_i, l_i) \in C, f_1, f_2 \in \mathcal{G}, \text{ and } R'_1 R'_2 \\
& \text{such } v \in f_1, u \in f_2, t = l_{i,1}, t' = l_{i,1} + 1 \\
& \text{and } (f_1, \mathbf{1}_{R'_1}), (f_2, \mathbf{1}_{R'_2}) \in l_{i,2} \}
\end{aligned}$$