

# SWEEP rule on the KD-Toric code

February 9, 2026

## 1 SWEEP Rule Decoder on the Toric

The decoder works as follows: each vertex  $v$  looks for a suggestion  $\hat{v}$  on  $\uparrow(v) \cap lk_2$  that matches  $\sigma_v^+$ . If there are more than one, then pick the one which has the minimal weight.




Figure 1: SWEEP rule

## 2 Span Expansion

Define the potentials functions  $L_i : \mathbb{R}^d \rightarrow \mathbb{R}$  as follow<sup>1</sup>:

$$L_i(\mathbf{x}) = \begin{cases} \mathbf{x}_i & i < d + 1 \\ -\langle \mathbf{1}, \mathbf{x} \rangle & i = d + 1 \end{cases}$$

The **Size** of a subset  $S$  is defined as:

$$\text{Size}(S) = \sum_i \max_{\mathbf{x} \in S} L_i(\mathbf{x})$$

Numerically, we show that the span property holds for the checks. Namely, let  $e$  be an unsatisfied check at time  $t$ , and let  $H_e$  be the unsatisfied checks at time  $t - 1$ . Then there is a set  $H'_e \subset H_e$  such that:

1. For any  $i < d + 1$  there exists  $e' \in H'_e$  such  $L_i(e') \geq L_i(e)$ .
2. There exists  $e' \in H'_e$  such that  $L_{d+1}(e') \leq L_{d+1}(e) + 1$ .

*Proof.* Consider a vertex  $v$  which at time  $t$  adjoins to an unsatisfied edge  $e$ , let  $u, w$  be vertices which have a positive faces that adjoin to  $e$ .

<sup>1</sup>Notice that we use opposite signs relative to Berman and Peter Gacs, yet we agree with Perskil.

1. If  $\hat{u}|_e \neq \emptyset$ . Idea: At least two (tree in 4D) directions are taken by faces on  $u$ , then we track over a faces paths that is positive in the remain direction, we know that we always can go forward, otherwise we close a stabilizer what contradicts .. (Require refine, cause we can go backward yet don't expect to back to the starting point).

Let  $G \subset \mathcal{F}$  be a connected component of faces that contains  $b_u$ . We say that  $G$  is  $d$  far from  $u$  if

$$\max_{f \in G} |f_x - u_x| \leq d$$

Now, we will prove by induction on  $d$  that there is a face  $f \in G$  with  $f_x \geq v_x$  such one of its edge is not satisfied.

- (a) **Base.** The edge from  $v$  which point at the  $\hat{z}$  direction must not be satisfied.
  - (b) **Assumption.** Assume that for any  $G$  which is at most  $d$  far from  $u$  there is a face, with adjoins edge that is both has a  $x$ -coordinate that is greater than  $u_x$  and is not satisfied.
  - (c) **Induction Step.** Notice that if by substitute a stabilizer from  $G$  we get  $G'$  that is at most  $d'$  far from  $u$  for some  $d' < d$  then we done.
2. Else, if for any  $u$  it holds that  $\hat{u}|_e = \emptyset$ , then it must be that  $e \in H_e$  because  $\hat{v}$  can only decrease the syndrome on  $v$ . Moreover, it follows that  $\hat{v} = \emptyset$  (otherwise, the suggestion would zero out the syndrome on  $\uparrow(v) \cap lk_2$ ). Thus,  $\sigma_v \not\subset \uparrow(v) \cap lk_2 \Rightarrow$  there is also an edge  $e' \in H_e$  where  $e' \in \downarrow(v)$ . In that case, take  $e \in H_e$  as the edge satisfies the first  $d$  equalities and  $e'$  for satisfying the  $d + 1$ th inequality.

□




Figure 2: SWEEP rule