

Final Recitation – Information Theory, Application for Quantum Fault Tolerance.

David Ponarovsky

June 25, 2025

Introduction

Today:

- ▶ Noisy circuit and noisy computation.

Introduction

Today:

- ▶ Noisy circuit and noisy computation.
- ▶ Classical no-go for reversible noisy computation at logarithmic depth (and polynomial space).

Introduction

Today:

- ▶ Noisy circuit and noisy computation.
- ▶ Classical no-go for reversible noisy computation at logarithmic depth (and polynomial space).
- ▶ Quantum case.

Introduction

Today:

- ▶ Noisy circuit and noisy computation.
- ▶ Classical no-go for reversible noisy computation at logarithmic depth (and polynomial space).
- ▶ Quantum case.
- ▶ Trading (local) entropy for space.

Nosiy Circuit.



Nosiy Circuit.

Definition

p - Depolarizing Channel. The qubit depolarizing channel with parameter $p \in [0, 1]$ is the quantum channel \mathcal{D}_p defined by:

$$\mathcal{D}_p(\rho) = (1 - p)\rho + p \cdot \frac{I}{2}$$

where ρ is a single-qubit density matrix and I is the identity matrix.

Nosiy Circuit.

Definition

p - Depolarizing Channel. The qubit depolarizing channel with parameter $p \in [0, 1]$ is the quantum channel \mathcal{D}_p defined by:

$$\mathcal{D}_p(\rho) = (1 - p)\rho + p \cdot \frac{I}{2}$$

where ρ is a single-qubit density matrix and I is the identity matrix.

Definition

p -Noisy Circuit. Given a circuit C (regardless of the model), its p -noisy version \tilde{C} is the circuit obtained by alternately taking layers from C and then passing each (qu)bit through a p -Depolarizing channel.

Classical no-go.

Claim

Let Y be a bit given by moving X through $\text{BSC}(p)$, Then there is $\gamma_p < 1$ such :

$$1 - H(Y) \leq \gamma(1 - H(X))$$

Classical no-go.

Denote by δ the parameter for which X distributed as $\sim \text{Bin}(\frac{1+\delta}{2})$.
First notice that:

$$\Pr(Y = 1) = \frac{1+\delta}{2}(1-p) + \frac{1-\delta}{2}p = \frac{1+\delta-2p\delta}{2}$$

So $Y \sim \text{Bin}(\frac{1+\delta-2p\delta}{2})$, Or $\delta \mapsto 1 - 2p\delta$.

Classical no-go.

Now expand $1 - H(X)$ to it's Taylor Series at δ gives:

$$1 - H(X) = 1 - \frac{1}{2} \left((1 + \delta) \log \left(\frac{1 + \delta}{2} \right) + (1 - \delta) \log \left(\frac{1 - \delta}{2} \right) \right)$$

Classical no-go.

Now expand $1 - H(X)$ to it's Taylor Seryias at δ gives:

$$\begin{aligned} 1 - H(X) &= 1 - \frac{1}{2} \left((1 + \delta) \log \left(\frac{1 + \delta}{2} \right) + (1 - \delta) \log \left(\frac{1 - \delta}{2} \right) \right) \\ &= -\frac{1}{2} \left((1 + \delta) \log \left(\frac{1 + \delta}{2} \right) + (1 - \delta) \log \left(\frac{1 - \delta}{2} \right) \right) \end{aligned}$$

Classical no-go.

Now expand $1 - H(X)$ to it's Taylor Series at δ gives:

$$\begin{aligned}1 - H(X) &= 1 - \frac{1}{2} \left((1 + \delta) \log \left(\frac{1 + \delta}{2} \right) + (1 - \delta) \log \left(\frac{1 - \delta}{2} \right) \right) \\&= -\frac{1}{2} \left((1 + \delta) \log \left(\frac{1 + \delta}{2} \right) + (1 - \delta) \log \left(\frac{1 - \delta}{2} \right) \right) \\&= -\frac{1}{2} \cdot (1 + \delta) \sum_{i=1}^{\infty} \frac{(-1)^{i+1} \delta^i}{i} + (1 - \delta) \sum_{i=1}^{\infty} \frac{(-1)^{i+1} (-\delta)^i}{i}\end{aligned}$$

Classical no-go.

Now expand $1 - H(X)$ to it's Taylor Series at δ gives:

$$\begin{aligned}1 - H(X) &= 1 - \frac{1}{2} \left((1 + \delta) \log \left(\frac{1 + \delta}{2} \right) + (1 - \delta) \log \left(\frac{1 - \delta}{2} \right) \right) \\&= -\frac{1}{2} \left((1 + \delta) \log \left(\frac{1 + \delta}{2} \right) + (1 - \delta) \log \left(\frac{1 - \delta}{2} \right) \right) \\&= -\frac{1}{2} \cdot (1 + \delta) \sum_{i=1}^{\infty} \frac{(-1)^{n+1} \delta^n}{n} + (1 - \delta) \sum_{i=1}^{\infty} \frac{(-1)^{n+1} (-\delta)^n}{n} \\&= -\frac{1}{2} \cdot \sum_{i=1}^{\infty} 2 \frac{\delta^{2n}}{2n} - \sum_{i=1}^{\infty} 2 \frac{\delta^{2n}}{2n - 1}\end{aligned}$$

Classical no-go.

Now expand $1 - H(X)$ to it's Taylor Series at δ gives:

$$\begin{aligned}1 - H(X) &= 1 - \frac{1}{2} \left((1 + \delta) \log \left(\frac{1 + \delta}{2} \right) + (1 - \delta) \log \left(\frac{1 - \delta}{2} \right) \right) \\&= -\frac{1}{2} \left((1 + \delta) \log \left(\frac{1 + \delta}{2} \right) + (1 - \delta) \log \left(\frac{1 - \delta}{2} \right) \right) \\&= -\frac{1}{2} \cdot (1 + \delta) \sum_{i=1}^{\infty} \frac{(-1)^{i+1} \delta^i}{i} + (1 - \delta) \sum_{i=1}^{\infty} \frac{(-1)^{i+1} (-\delta)^i}{i} \\&= -\frac{1}{2} \cdot \sum_{i=1}^{\infty} 2 \frac{\delta^{2i}}{2i} - \sum_{i=1}^{\infty} 2 \frac{\delta^{2i}}{2i - 1} \\&= \sum_{i=1}^{\infty} \frac{\delta^{2i}}{2i(2i - 1)}\end{aligned}$$

Classical no-go.

Now expand $1 - H(X)$ to it's Taylor Series at δ gives:

$$\begin{aligned}1 - H(X) &= 1 - \frac{1}{2} \left((1 + \delta) \log \left(\frac{1 + \delta}{2} \right) + (1 - \delta) \log \left(\frac{1 - \delta}{2} \right) \right) \\&= -\frac{1}{2} \left((1 + \delta) \log \left(\frac{1 + \delta}{2} \right) + (1 - \delta) \log \left(\frac{1 - \delta}{2} \right) \right) \\&= -\frac{1}{2} \cdot (1 + \delta) \sum_{i=1}^{\infty} \frac{(-1)^{n+1} \delta^n}{n} + (1 - \delta) \sum_{i=1}^{\infty} \frac{(-1)^{n+1} (-\delta)^n}{n} \\&= -\frac{1}{2} \cdot \sum_{i=1}^{\infty} 2 \frac{\delta^{2n}}{2n} - \sum_{i=1}^{\infty} 2 \frac{\delta^{2n}}{2n - 1} \\&= \sum_{i=1}^{\infty} \frac{\delta^{2n}}{2n(2n - 1)}\end{aligned}$$

Denote the above by $K(\delta)$