

# Quantum Information Theory - 67749

## Recitation 2, May 7, 2025

### 1 Overview - Quantum States as Computational Resources.

In the last lectures, we saw that quantum states can be considered as resources. In particular, we saw that shared **EPR** pair ( $\mathbf{Bell}_{00}$ ) enables one:

1. Transmit two classical bits by sending a single qubit, via the superdense-coding.
2. 'Teleoperate' a qubit by sending two classical bits. From an engineering point of view, it means that for having a complete quantum internet, it's enough to provide a mechanism to distribute **EPR** pairs.

### 2 Dense Encoding.

)

### 3 Quantum Teleportation.



Figure 1: Measuring the single-qubit state  $|\psi\rangle$  at the  $\{|+\rangle, |-\rangle\}$  base.

## 4 Gate Teleportation.

Gate teleportation is a method to 'encode' operations by states. At the high level, given a precomputed state, it allows one to apply an operation (gate) by using (probably) simpler gates. The precomputed states are called **Magic States**.

### 4.1 Leading Example: $T$ -Teleportation.

Recall that the Clifford<sup>1</sup> +  $T$  is a universal quantum gate set. The Clifford group alone is considered from the computer science point of view a simple/weak computational class since it can be classically simulated<sup>2</sup>. Yet, we will see that given access to the magic  $|T\rangle = T|+\rangle$ , one can simulate the  $T$  gate using only Clifford gates and measurements.




Figure 2: Measuring the single-qubit state  $|\psi\rangle$  at the  $\{|+\rangle, |-\rangle\}$  base.

$$\begin{aligned}
 \left( \sum_x \alpha_x |x\rangle \right) \otimes \frac{1}{\sqrt{2}} (|0\rangle + e^{i\frac{\pi}{4}} |1\rangle) &\xrightarrow{\text{CX}} \sum_{x,y} \frac{1}{\sqrt{2}} \alpha_x |x\rangle |x \oplus y\rangle e^{i\frac{\pi}{4}y} \\
 &\mapsto \begin{cases} \sum_x \alpha_x |x\rangle e^{i\frac{\pi}{4}x} = T|\psi\rangle & \text{measured 0} \\ \sum_x \alpha_x |x\rangle e^{i\frac{\pi}{4}\bar{x}} & \text{measured 1} \end{cases} \\
 &\xrightarrow{\text{CS}} \begin{cases} T|\psi\rangle \\ \sum_x \alpha_x |x\rangle e^{i(\frac{\pi}{4}\bar{x} + \frac{\pi}{2}x)} = \sum_x \alpha_x |x\rangle e^{i\frac{\pi}{4}} e^{i(\frac{\pi}{4}\bar{x} + \frac{\pi}{4}x)} \end{cases} \\
 &= \begin{cases} T|\psi\rangle \\ e^{i\frac{\pi}{4}} \sum_x \alpha_x |x\rangle e^{i\frac{\pi}{4}} = e^{i\frac{\pi}{4}} T|\psi\rangle \end{cases}
 \end{aligned}$$

### 4.2 Extends it.

Let's extend it to a general gate. First create  $|\mathbf{GHZ}_{2n}\rangle$  state, then

Let's split upon the measurement result.

1. If we measured 0, means the states 'agreed' in the computational base.

$$|\psi\rangle \otimes \left( \sum_x |x\rangle \otimes U|x\rangle \right)$$

---

<sup>1</sup>Generated by  $H, S$  and  $CX$

<sup>2</sup>And conjectured to be strictly weaker than  $\mathbf{P}$

## 5 Uhlmann's theorem

**Claim 5.1.**

$$\langle \Omega | A \otimes B | \Omega \rangle = \text{Tr} AB^\dagger$$

*Proof.*

$$\begin{aligned} \langle \Omega | A \otimes B | \Omega \rangle &= \sum_{ij} \langle i; i | AB | j; j \rangle = \sum_{ij} \langle i | A | j \rangle \langle i | B | j \rangle = \sum_{ij} \langle i | A | j \rangle \langle j | B^\dagger | i \rangle \\ &= \sum_i \langle i | AB^\dagger | i \rangle = \text{Tr} AB^\dagger \end{aligned}$$

□

$$|\psi_\rho\rangle = \sum_i \left( \rho^{\frac{1}{2}} |\psi_i\rangle \right) |i\rangle = \sum_i \left( \rho^{\frac{1}{2}} U_\rho |i\rangle \right) |i\rangle = \left( \rho^{\frac{1}{2}} U_\rho \right) \otimes I |\Omega\rangle$$

$$|\psi_\sigma\rangle = \sum_i \left( \sigma^{\frac{1}{2}} |\psi'_i\rangle \right) |i'\rangle = \sum_i \left( \sigma^{\frac{1}{2}} U_\sigma |i\rangle \right) V |i\rangle = \left( \sigma^{\frac{1}{2}} U_\sigma \right) \otimes V |\Omega\rangle$$

**Claim 5.2.** For any square matrix  $A$ :

$$\max_{U \in \mathcal{U}} \text{Tr} AU = \text{Tr} \sqrt{A^\dagger A}$$

$$\begin{aligned} \max |\langle \psi_\rho | \psi_\sigma \rangle|^2 &= \max |\langle \Omega | \left( U_\rho^\dagger \rho^{\frac{1}{2}} \right) \otimes I \left( \sigma^{\frac{1}{2}} U_\sigma \right) \otimes V | \Omega \rangle|^2 \\ &= \max |\text{Tr} \left[ \left( U_\rho^\dagger \rho^{\frac{1}{2}} \sigma^{\frac{1}{2}} U_\sigma \right) V^\dagger \right]|^2 \\ &= \max |\text{Tr} \left[ \rho^{\frac{1}{2}} \sigma^{\frac{1}{2}} V^\dagger \right]|^2 \\ &\leq \left| \text{Tr} \sqrt{\rho^{\frac{1}{2}} \sigma^{\frac{1}{2}} \sigma^{\frac{1}{2}} \rho^{\frac{1}{2}}} \right|^2 = \left| \text{Tr} \sqrt{\rho^{\frac{1}{2}} \sigma \rho^{\frac{1}{2}}} \right|^2 \end{aligned}$$

## 6 Monotonicity of Fidelity.

Let  $\rho_{AB}, \sigma_{AB} \in \mathcal{D}(\mathcal{H}_A \otimes \mathcal{H}_B)$ . Then the fidelity is non-decreasing with respect to the partial trace:

$$F(\rho_{AB}, \sigma_{AB}) \leq F(\rho_A, \sigma_A),$$

where  $\rho_A = \text{Tr}_B \{\rho_{AB}\}$  and  $\sigma_A = \text{Tr}_B \{\sigma_{AB}\}$ .

*Proof.* Consider fixed purifications  $|\psi\rangle_{RAB}$  and  $|\phi\rangle_{RAB}$  of  $\rho_{AB}$  and  $\sigma_{AB}$ , respectively, which also purify  $\rho_A$  and  $\sigma_A$ . By Uhlmann's theorem,

$$F(\rho_{AB}, \sigma_{AB}) = \max_{U_R} |\langle \psi | U_R \otimes I_A \otimes I_B | \phi \rangle|^2.$$

On the other hand, since  $U_R \otimes I_A$  is a subset of the larger class of unitaries  $U_{RB}$  on  $RB$ ,

$$F(\rho_A, \sigma_A) = \max_{U_{RB}} |\langle \psi | U_{RB} \otimes I_A | \phi \rangle|^2 \geq F(\rho_{AB}, \sigma_{AB}).$$

Thus, we conclude that

$$F(\rho_{AB}, \sigma_{AB}) \leq F(\rho_A, \sigma_A).$$

□

Notice that  $|i\rangle\langle j|$  is unitray since.

## 7 $|\text{EPR}\rangle$ Distillation.

Distillation of entanglement is a process in quantum information theory where a mixed entangled state is transformed into a more pure form of entanglement. This is achieved by using local operations and classical communication (LOCC) to extract a smaller number of highly entangled pairs from a larger number of weakly entangled pairs. The goal is to maximize the quality of entanglement, making it more suitable for quantum communication and computation tasks.

The process involves multiple copies of a mixed entangled state, which are manipulated to increase the fidelity of the entanglement. Distillation protocols, such as the Bennett-Brassard protocol, are used to achieve this transformation. The distilled entangled states are more robust and can be used for tasks like quantum teleportation and superdense coding, where high-quality entanglement is crucial for optimal performance.




Figure 3: Measuring the single-qubit state  $|\psi\rangle$  at the  $\{|+\rangle, |-\rangle\}$  base.

$$\rho = p |\beta_{00}\rangle \langle \beta_{00}| + \frac{1-p}{3} \sum_{j \neq 00} |\beta_j\rangle \langle \beta_j|$$

The density matrix over each qubit is  $\frac{1}{2}I$ , so the measurement is equivalent to flipping coins and asking if the first pair and second pair are the same. Thus, the success probability is  $\frac{1}{4}$ . Now the state that is left is the projection of  $\rho$  into the space in which both the bits of Alice and Bob are equal, namely:

$$\begin{aligned}
\frac{1}{4}((|00\rangle \pm |11\rangle) \otimes (|00\rangle \pm |11\rangle)) &\rightarrow (|0000\rangle + |1111\rangle) \\
\frac{1}{4}((|00\rangle \pm |11\rangle) \otimes (|00\rangle \mp |11\rangle)) &\rightarrow (|0000\rangle - |1111\rangle) \\
\frac{1}{4}((|00\rangle \pm |11\rangle) \otimes (|01\rangle \pm |10\rangle)) &\rightarrow \emptyset \\
\frac{1}{4}((|01\rangle \pm |10\rangle) \otimes (|01\rangle \pm |10\rangle)) &\rightarrow (|0101\rangle + |1010\rangle) \\
\frac{1}{4}((|01\rangle \pm |10\rangle) \otimes (|01\rangle \mp |10\rangle)) &\rightarrow (|0101\rangle - |1010\rangle)
\end{aligned}$$

$$p^2 + 2\frac{p(1-p)}{3} + \frac{1}{3}(1-p)^2 + 4\frac{1}{3}(1-p)^2$$

$$p' \leftarrow \frac{p^2 + \frac{1}{3}(1-p)^2}{p^2 + 2\frac{p(1-p)}{3} + \frac{1}{3}(1-p)^2 + 4\frac{1}{3}(1-p)^2}$$

## 8 Magic State Distillation.

**Question.** Can we purify noisy magic states into high-fidelity ones, using only Clifford operations?

Magic state distillation is a procedure that uses many copies of noisy magic states, plus only Clifford gates and measurements, to produce fewer, higher-fidelity magic states.