

Quantum Information Theory - 67749

Recitation 2, May 6, 2025

1 Overview - Quantum States as Computational Resources.

In the last lectures, we saw that quantum states can be considered as resources. In particular, we saw that shared **EPR** pair (Bell_{00}) enables one:

1. Transmit two classical bits by sending a single qubit, via the superdense-coding.
2. 'Teleoperate' a qubit by sending two classical bits. From an engineering point of view, it means that for having a complete quantum internet, it's enough to provide a mechanism to distribute **EPR** pairs.

2 Dense Encoding.

)

3 Quantum Teleportation.

4 Gate Teleportation.

Gate teleportation is a method to 'encode' operations by states. At the high level, given a precomputed state, it allows one to apply an operation (gate) by using (probably) simpler gates. The precomputed states are called **Magic States**.

4.1 Leading Example: T -Teleportation.

Recall that the Clifford¹ + T is a universal quantum gate set. The Clifford group alone is considered from the computer science point of view a simple/weak

¹Generated by H, S and CX

computational class since it can be classically simulated ². Yet, we will see that given access to the magic $|T\rangle = T|+\rangle$, one can simulate the T gate using only Clifford gates and measurements.



Figure 1: Measuring the single-qubit state $|\psi\rangle$ at the $\{|+\rangle, |-\rangle\}$ base.

$$\begin{aligned}
\left(\sum_x \alpha_x |x\rangle\right) \otimes \frac{1}{\sqrt{2}} (|0\rangle + e^{i\frac{\pi}{4}} |1\rangle) &\xrightarrow{\text{CX}} \sum_{x,y} \frac{1}{\sqrt{2}} \alpha_x |x\rangle |x \oplus y\rangle e^{i\frac{\pi}{4}y} \\
&\mapsto \begin{cases} \sum_x \alpha_x |x\rangle e^{i\frac{\pi}{4}x} = T|\psi\rangle & \text{measured 0} \\ \sum_x \alpha_x |x\rangle e^{i\frac{\pi}{4}\bar{x}} & \text{measured 1} \end{cases} \\
&\xrightarrow{\text{CS}} \begin{cases} T|\psi\rangle \\ \sum_x \alpha_x |x\rangle e^{i(\frac{\pi}{4}\bar{x} + \frac{\pi}{2}x)} = \sum_x \alpha_x |x\rangle e^{i\frac{\pi}{4}} e^{i(\frac{\pi}{4}\bar{x} + \frac{\pi}{4}x)} \end{cases} \\
&= \begin{cases} T|\psi\rangle \\ e^{i\frac{\pi}{4}} \sum_x \alpha_x |x\rangle e^{i\frac{\pi}{4}} = e^{i\frac{\pi}{4}} T|\psi\rangle \end{cases}
\end{aligned}$$

4.2 Extends it.

Let's extend it to a general gate. First create $|\mathbf{GHZ}_{2n}\rangle$ state, then

Let's split upon the measurement result.

1. If we measured 0, means the states 'agreed' in the computational base.

$$|\psi\rangle \otimes \left(\sum_x |x\rangle \otimes U|x\rangle\right)$$

5 Magic State Distillation.

Question. Can we purify noisy magic states into high-fidelity ones, using only Clifford operations?

Magic state distillation is a procedure that uses many copies of noisy magic states, plus only Clifford gates and measurements, to produce fewer, higher-fidelity magic states.

²And conjectured to be strictly weaker than \mathbf{P}