Quantum Information Theory - 67749 Exercise 2, May 20, 2025

1 Submission Guidelines.

- Due date May 31, 2025.
- Make sure your submission is clear. Unreadable assignments will get zero score.
- Using any Generative AI tool is forbidden but not enforced. Yet, please keep in mind that we might call you for an interview about your assignment.

2 Fidelity.

Compute the fidelity between ρ and σ in the following cases:

1. Use the construction presented in the proof of Uhlman's to calculate the fidelity between: $\rho = \frac{4}{7} |0\rangle \langle 0| + \frac{3}{7} |+\rangle \langle +|$ and $\sigma = \frac{3}{7} |0\rangle \langle 0| + \frac{4}{7} |+\rangle \langle +|$.

Solution. Observes that $\sigma = H\rho H$, Since H is a unitrary then $\sigma^{\frac{1}{2}} = H\rho^{\frac{1}{2}}H$. Thus:

$$\max \mathbf{Tr} \sigma^{\frac{1}{2}} \rho^{\frac{1}{2}} V^{\dagger} = \max \mathbf{Tr} \sigma^{\frac{1}{2}} H \sigma^{\frac{1}{2}} H V^{\dagger} = \mathbf{Tr} \sigma^{\frac{1}{2}} H \sigma^{\frac{1}{2}} H$$

$$\sigma = \left(1 - \frac{4}{7} + \frac{2}{7}\right) |0\rangle \langle 0| + \frac{2}{7} |0\rangle \langle 1| + \frac{2}{7} |1\rangle \langle 0| + \frac{2}{7} |1\rangle \langle 1| = \begin{bmatrix} \frac{5}{7} & \frac{2}{7} \\ \frac{7}{7} & \frac{2}{7} \end{bmatrix}$$

$$\Rightarrow \left(\frac{5}{7} - \lambda\right) \left(\frac{2}{7} - \lambda\right) - \frac{4}{49} = 0$$

$$\Rightarrow \lambda^2 - \lambda - \frac{6}{49} = 0 \Rightarrow \lambda_{\pm} = \frac{1}{2} \pm \frac{1}{2} \cdot \frac{5}{7}$$

So the eigenvalues are $\lambda_+ = \frac{6}{7}$ and $\lambda_- = \frac{1}{7}$ with eigenvectors:

$$|\omega_{+}\rangle = \sqrt{\frac{2}{5}} \begin{bmatrix} 1 \\ \frac{1}{2} \end{bmatrix}$$
 and $|\omega_{-}\rangle = \frac{1}{\sqrt{5}} \begin{bmatrix} 1 \\ -2 \end{bmatrix}$

Thus $\sigma^{\frac{1}{2}}$ equals:

$$\lambda_{+}^{\frac{1}{2}} \left| \omega_{+} \right\rangle \left\langle \omega_{+} \right| + \lambda_{-}^{\frac{1}{2}} \left| \omega_{-} \right\rangle \left\langle \omega_{-} \right| = \sqrt{\frac{6}{7}} \frac{2}{5} \begin{bmatrix} 1 & \frac{1}{2} \\ \frac{1}{2} & \frac{1}{4} \end{bmatrix} + \sqrt{\frac{1}{7}} \frac{1}{5} \begin{bmatrix} 1 & -2 \\ -2 & 4 \end{bmatrix}$$

$$\rho = \left(1 - \frac{3}{7} + \frac{3}{7 \cdot 2}\right) |0\rangle \langle 0| + \frac{3}{7 \cdot 2} |0\rangle \langle 1| + \frac{3}{7 \cdot 2} |1\rangle \langle 0| + \frac{3}{7 \cdot 2} |1\rangle \langle 1| = \begin{bmatrix} \frac{11}{14} & \frac{3}{14} \\ \frac{3}{14} & \frac{3}{14} \end{bmatrix}$$

$$\Rightarrow \left(\frac{11}{14} - \lambda\right) \left(\frac{3}{14} - \lambda\right) - \frac{9}{196} = 0$$

$$\Rightarrow \lambda^2 - \lambda - \frac{9}{196} = 0 \Rightarrow \lambda_{\pm} = \frac{1}{2} \pm \frac{1}{2} \cdot \frac{4\sqrt{10}}{14} = \frac{7 + 2\sqrt{10}}{14}, \frac{7 - 2\sqrt{10}}{14}$$

W are almost ready to compute $\sigma^{\frac{1}{2}}H$ for that first let's compute:

$$\begin{aligned} |\omega_{+}\rangle \langle \omega_{+}| \, H &= \frac{2}{5 \cdot \sqrt{2}} \begin{bmatrix} 1 & \frac{1}{2} \\ \frac{1}{2} & \frac{1}{4} \end{bmatrix} \begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix} = \frac{2}{5 \cdot \sqrt{2}} \begin{bmatrix} \frac{3}{2} & \frac{1}{2} \\ \frac{3}{4} & \frac{1}{4} \end{bmatrix} = \frac{1}{5 \cdot \sqrt{2}} \begin{bmatrix} 3 & 1 \\ \frac{3}{2} & \frac{1}{2} \end{bmatrix} \\ |\omega_{-}\rangle \langle \omega_{-}| \, H &= \frac{1}{5 \cdot \sqrt{2}} \begin{bmatrix} 1 & -2 \\ -2 & 4 \end{bmatrix} \begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix} = \frac{1}{5 \cdot \sqrt{2}} \begin{bmatrix} -1 & 3 \\ 2 & -6 \end{bmatrix} \end{aligned}$$

Thus:

$$\sigma^{\frac{1}{2}}H = \frac{1}{5 \cdot \sqrt{2}} \begin{bmatrix} 3 \cdot \sqrt{\frac{6}{7}} - 1 \cdot \sqrt{\frac{1}{7}} & 1 \cdot \sqrt{\frac{6}{7}} + 4 \cdot \sqrt{\frac{1}{7}} \\ \frac{3}{2} \cdot \sqrt{\frac{6}{7}} + 2 \cdot \sqrt{\frac{1}{7}} & \frac{1}{2} \cdot \sqrt{\frac{6}{7}} - 6 \cdot \sqrt{\frac{1}{7}} \end{bmatrix}$$

aaa

Let $|\psi_{\rho}\rangle$ and $|\psi_{\sigma}\rangle$ be purifications of ρ and σ .

$$\langle \psi_{\rho} | \psi_{\sigma} \rangle$$