Problem Set 3.4

1 (Recommended) Execute the six steps of Worked Example **3.4 A** to describe the column space and nullspace of A and the complete solution to $A\boldsymbol{x} = \boldsymbol{b}$:

$$A = \begin{bmatrix} 2 & 4 & 6 & 4 \\ 2 & 5 & 7 & 6 \\ 2 & 3 & 5 & 2 \end{bmatrix} \qquad \boldsymbol{b} = \begin{bmatrix} b_1 \\ b_2 \\ b_3 \end{bmatrix} = \begin{bmatrix} 4 \\ 3 \\ 5 \end{bmatrix}$$

2 Carry out the same six steps for this matrix A with rank one. You will find *two* conditions on b_1, b_2, b_3 for $A\boldsymbol{x} = \boldsymbol{b}$ to be solvable. Together these two conditions put $\boldsymbol{b}$ into the _____ space (two planes give a line):

$$A = \begin{bmatrix} 1 \\ 3 \\ 2 \end{bmatrix} \begin{bmatrix} 2 & 1 & 3 \end{bmatrix} = \begin{bmatrix} 2 & 1 & 3 \\ 6 & 3 & 9 \\ 4 & 2 & 6 \end{bmatrix} \qquad \boldsymbol{b} = \begin{bmatrix} b_1 \\ b_2 \\ b_3 \end{bmatrix} = \begin{bmatrix} 10 \\ 30 \\ 20 \end{bmatrix}$$

* Just get the complete solution

20 Reduce A to its echelon form U. Then find a triangular L so that $A = LU$.

$$A = \begin{bmatrix} 3 & 4 & 1 & 0 \\ 6 & 5 & 2 & 1 \end{bmatrix} \quad \text{and} \quad A = \begin{bmatrix} 1 & 0 & 1 & 0 \\ 2 & 2 & 0 & 3 \\ 0 & 6 & 5 & 4 \end{bmatrix}.$$

21 Find the complete solution in the form $\boldsymbol{x}_p + \boldsymbol{x}_n$ to these full rank systems:

(a) $x + y + z = 4$ (b) $\begin{aligned} x + y + z &= 4 \\ x - y + z &= 4. \end{aligned}$

Problem Set 3.5

Questions 1–10 are about linear independence and linear dependence.

1 Show that $\boldsymbol{v}_1, \boldsymbol{v}_2, \boldsymbol{v}_3$ are independent but $\boldsymbol{v}_1, \boldsymbol{v}_2, \boldsymbol{v}_3, \boldsymbol{v}_4$ are dependent:

$$\boldsymbol{v}_1 = \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} \quad \boldsymbol{v}_2 = \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix} \quad \boldsymbol{v}_3 = \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} \quad \boldsymbol{v}_4 = \begin{bmatrix} 2 \\ 3 \\ 4 \end{bmatrix}.$$

Solve $c_1\boldsymbol{v}_1 + c_2\boldsymbol{v}_2 + c_3\boldsymbol{v}_3 + c_4\boldsymbol{v}_4 = \mathbf{0}$ or $A\boldsymbol{x} = \mathbf{0}$. The $\boldsymbol{v}$'s go in the columns of A.