

History of elliptic curves rank records

Let E be an elliptic curve over \mathbf{Q} . By Mordell's theorem, $E(\mathbf{Q})$ is a finitely generated abelian group. This means that $E(\mathbf{Q}) = E(\mathbf{Q})_{\text{tors}} \times \mathbf{Z}^r$. By Mazur's theorem, we know that $E(\mathbf{Q})_{\text{tors}}$ is one of the following 15 groups:

$\mathbf{Z}/n\mathbf{Z}$ with $1 \leq n \leq 10$ or $n = 12$,
 $\mathbf{Z}/2\mathbf{Z} \times \mathbf{Z}/2m\mathbf{Z}$ with $1 \leq m \leq 4$.

On the other hand, it is not known what values of rank r are possible for elliptic curves over \mathbf{Q} . The "folklore" conjecture is that a rank can be arbitrary large. The current record is an example of elliptic curve with **rank ≥ 28 , found by Elkies in 2006** (the previous record was rank ≥ 24 , found by Martin and McMillen in 2000).

The highest rank of an elliptic curve which is known exactly (not only a lower bound for rank) is equal to [19](#), and it was found by Elkies in 2009. It improves previous records due to Kretschmer (rank = [10](#)), Schneiders-Zimmer (rank = [11](#)), Fermigier (rank = [14](#)), Dujella (rank = [15](#)) and Elkies (rank = [17](#), rank = [18](#)).

The following table contains some historical data on elliptic curve rank records.

rank \geq	year	Author(s)
3	1938	Billing
4	1945	Wiman
6	1974	Penney - Pomerance
7	1975	Penney - Pomerance
8	1977	Grunewald - Zimmert
9	1977	Brumer - Kramer
12	1982	Mestre
14	1986	Mestre
15	1992	Mestre
17	1992	Nagao
19	1992	Fermigier
20	1993	Nagao
21	1994	Nagao - Kouya
22	1997	Fermigier
23	1998	Martin - McMillen
24	2000	Martin - McMillen
28	2006	Elkies

Click on rank r to see the corresponding curve(s) and independent points P_1, P_2, \dots, P_r of infinite order.

References:

1. G. Billing, *Beitrage zur arithmetischen Theorie der ebenen kubischen Kurven vom Geschlecht Eins*, Nova Acta Soc. Sci. Upsal. (4) **11** (1938), Nr.1, Diss.165.
2. A. Wiman, *Über den Rang von Kurven $y^2 = x(x+a)(x+b)$* , Acta Math. **76** (1945), 225-251.
3. A. Wiman, *Über rationale Punkte auf Kurven $y^2 = x(x^2-c^2)$* , Acta Math. **77** (1945), 281-320.
4. D.E. Penney and C. Pomerance, *A search for elliptic curves with large rank*, Math. Comp. **28** (1974), 851-853.
5. D.E. Penney and C. Pomerance, *Three elliptic curves with rank at least seven*, Math. Comp. **29** (1975), 965-967.
6. F.J. Grunewald and R. Zimmert, *Über einige rationale elliptische Kurven mit treiem Rang ≥ 8* , J. Reine Angew. Math. **296** (1977), 100-107.
7. A. Brumer and K. Kramer, *The rank of elliptic curves*, Duke Math. J. **44** (1977), 715-743.
8. J.-F. Mestre, *Construction d'une courbe elliptique de rang ≥ 12* , C. R. Acad. Sci. Paris Ser I Math. **295** (1982), 643-644.
9. J.-F. Mestre, *Courbes elliptiques et formules explicites*, Theorie des nombres, Semin. Delange-Pisot-Poitou, Paris 1981/82, Prog. Math. 38, Birkhauser, Boston, 1983, pp. 179-187.
10. J.-F. Mestre, *Formules explicites et minorations de conducteurs de varietes algebriques*, Compositio Math. **58** (1986), 209-232.
11. J.-F. Mestre, *Un exemple de courbe elliptique sur \mathbf{Q} de rang ≥ 15* , C. R. Acad. Sci. Paris Ser I Math. **314** (1992), 453-455.
12. K. Nagao, *Examples of elliptic curves over \mathbf{Q} with rank ≥ 17* , Proc. Japan Acad. Ser. A Math. Sci. **68** (1992), 287-289.
13. S. Fermigier, *Un exemple de courbe elliptique definie sur $\mathbf{Q}(t)$ de rang ≥ 19* , C. R. Acad. Sci. Paris Ser. I **315** (1992), 719-722.
14. K. Nagao, *An example of elliptic curve over \mathbf{Q} with rank ≥ 20* , Proc. Japan Acad. Ser. A Math. Sci. **69** (1993), 291-293.
15. K. Nagao and T. Kouya, *An example of elliptic curve over \mathbf{Q} with rank ≥ 21* , Proc. Japan Acad. Ser. A Math. Sci. **70** (1994), 104-105.
16. K. Nagao, *Construction of high-rank elliptic curves*, Kobe J. Math. **11** (1994), 211-219.
17. S. Fermigier, *Une courbe elliptique definie sur $\mathbf{Q}(t)$ de rang ≥ 22* , Acta Arith. (1997), 359-363.

18. R. Martin and W. McMillen, *An elliptic curve over \mathbf{Q} with rank at least 23*, Number Theory Listserver, March 1998.
 19. R. Martin and W. McMillen, *An elliptic curve over \mathbf{Q} with rank at least 24*, Number Theory Listserver, May 2000.
 20. N. D. Elkies, \mathbf{Z}^{28} in $E(\mathbf{Q})$, etc., Number Theory Listserver, May 2006.
 21. N. D. Elkies, *Three lectures on elliptic surfaces and curves of high rank*, Lecture notes, Oberwolfach, 2007, arXiv:0709.2908.
-
22. T.J. Kretschmer, *Construction of elliptic curves with large rank*, Math. Comp. **46** (1986), 627-635.
 23. U. Schneiders and H.G. Zimmer, *The rank of elliptic curves upon quadratic extensions*, in: Computational Number Theory (A. Petho, H.C. Williams, H.G. Zimmer, eds.), de Gruyter, Berlin, 1991, pp. 239-260.
 24. S. Fermigier, *Exemples de courbes elliptiques de grand rang sur $\mathbf{Q}(t)$ et sur \mathbf{Q} possédant des points d'ordre 2*, C. R. Acad. Sci. Paris Ser. I **322** (1996), 949-952.
 25. A. Dujella, *An example of elliptic curve over \mathbf{Q} with rank equal to 15*, Proc. Japan Acad. Ser. A Math. Sci. **78** (2002), 109-111.
 26. N. D. Elkies, $E(\mathbf{Q}) = (\mathbf{Z}/2\mathbf{Z}) * \mathbf{Z}^{17}$, Number Theory Listserver, Jun 2005.
 27. N. D. Elkies, *Some more rank records: $E(\mathbf{Q}) = (\mathbf{Z}/2\mathbf{Z}) * \mathbf{Z}^{18}$, $(\mathbf{Z}/4\mathbf{Z}) * \mathbf{Z}^{12}$, $(\mathbf{Z}/8\mathbf{Z}) * \mathbf{Z}^6$, $(\mathbf{Z}/2\mathbf{Z}) * (\mathbf{Z}/6\mathbf{Z}) * \mathbf{Z}^6$* , Number Theory Listserver, Jun 2006.
 28. N. D. Elkies, Personal communication, 2009.

[High rank elliptic curves with prescribed torsion](#)

[Infinite families of elliptic curves with high rank and prescribed torsion](#)

[High rank elliptic curves with prescribed torsion over quadratic fields](#)

[Andrej Dujella home page](#)