

**RAD HRVATSKE AKADEMIJE ZNANOSTI I UMJETNOSTI  
MATEMATIČKE ZNANOSTI**

A. Al-Kateeb and A. Dagher

*On the number of terms of some families of the ternary cyclotomic  
polynomials  $\Phi_{3p_2p_3}$*

**Manuscript accepted for publication**

This is a preliminary PDF of the author-produced manuscript that has been  
peer-reviewed and accepted for publication. It has not been copy-edited,  
proofread, or finalized by Rad HAZU Production staff.

# ON THE NUMBER OF TERMS OF SOME FAMILIES OF THE TERNARY CYCLOTOMIC POLYNOMIALS $\Phi_{3p_2p_3}$

ALA'A AL-KATEEB AND AFNAN DAGHER

ABSTRACT. We study the number of non-zero terms in two specific families of ternary cyclotomic polynomials, we find formulas for the number of terms by writing the cyclotomic polynomial as a sum of smaller sub-polynomials and study the properties of these polynomials.

## 1. INTRODUCTION

The  $n$ -th cyclotomic polynomial  $\Phi_n$  is defined as the monic polynomial in  $\mathbb{Z}[x]$  whose complex roots are the primitive  $n$ -th roots of unity. Due to its importance in many branches of mathematics, there have been extensive investigation on its properties. Recently, Sanna in [20], write a concise survey attempts to collect the main results regarding the coefficients of the cyclotomic polynomials and to provide all the relevant references to their proofs.

The investigation on height (maximum absolute value of coefficients) was initiated by the finding that the height can be bigger than 1. It has produced numerous results, to list a few [2, 6, 5, 7, 13, 11, 12, 19, 10, 15, 16, 22].

The investigation on maximum gap (maximum difference between the consecutive exponents) became a problem on its own because it could be viewed as a first step toward understanding of sparsity structure of cyclotomic polynomials. In 2012, Hong, et al. [14] proved that the maximum gap for binary cyclotomic polynomial  $\Phi_{p_1p_2}$  is  $p_1 - 1$ , that is,  $g(\Phi_{p_1p_2}) = p_1 - 1$ . In 2014, Moree [18] revisited the result and provided an inspiring conceptual proof by making a connection to numerical semigroups of embedding dimension two. In 2016, Zhang [21] gave a simpler proof, along with the result on the number of occurrences of the maximum gaps. In 2021, Al-Kateeb, et al. [3] proved that  $g(\Phi_{mp}) = \varphi(m)$ , where  $m$  is a square free odd integer and  $p > m$  is prime number.

The investigation of the number of non-zero terms in  $\Phi_n(x)$  (also called the hamming weight  $\text{hw}(f)$ ) was initiated in 1965 by Carlitz [9], who gave an explicit formula for  $\text{hw}(\Phi_{pq})$ , where  $p < q$  are two distinct prime numbers. Hence, a natural question is whether there is a formula for the number of nonzero terms of a ternary cyclotomic polynomials  $\Phi_{pqr}$ , where  $p < q < r$  are three odd prime numbers, and ultimately, arbitrary cyclotomic polynomials? In 2014 Bezdega [8] prove that the hamming weight of the cyclotomic polynomial  $\Phi_n(x)$  is greater than or equal to  $n^{\frac{1}{3}}$ . In 2016 A. Al-Kateeb [1] investigated the number of terms for a ternary cyclotomic polynomial and the following theorem was given (Theorem 7.1).

**THEOREM 1.1.** *Let  $p_1 < p_2 < p_3$  be odd prime numbers such that  $p_2 \equiv \pm 1 \pmod{p_1}$  and  $p_3 \equiv \pm 1 \pmod{p_1p_2}$  and  $p_3 > p_1p_2$ . Then*

$$\text{hw}(\Phi_{p_1p_2p_3}) = \begin{cases} N(p_3 - 1) + 1, & r_3 = 1 \\ N(p_3 + 1) - 1, & r_3 = p_2 - 1 \end{cases}$$

---

2010 *Mathematics Subject Classification.* 11B83.

*Key words and phrases.* Number of terms, Cyclotomic polynomials, maximum gap.

where  $N = \frac{2}{3} \left( \frac{(p_1-1)((p_1+4)(p_2-1)-(r_2-1))}{p_1 p_2} \right)$  and  $r_2 = p_2 \mod p_1, r_3 = p_3 \mod p_1 p_2$ .

It is natural to think about fixing the smallest prime number  $p_1$  or to consider some small values of  $r_2$  and  $r_3$ , since computations show that the structure of  $\Phi_{p_1 p_2 p_3}$  is simpler for small values of  $p_1, r_2$  or  $r_3$ . In this paper, we tackle the number of terms for of the cyclotomic polynomial  $\Phi_{3p_2 p_3}$  where  $p_3 \equiv \pm 2 \mod 3p_2$ . We come up with a nice formulas (Theorems 2.1 and 2.4) for  $\text{hw}(\Phi_{3p_2 p_3})$  where  $p_3 \equiv \pm 2 \mod 3p_2$ . To prove the results we use proof techniques used for studying cyclotomic polynomials in [1].

This paper is structured as follows: In Section 2, we list the main results of the paper. In Section 3, we give several preliminaries and properties of cyclotomic polynomials needed for the rest of the paper. Then, in Section 4, we prove the results of the paper. In Appendix A, we list some technical proofs and finally in Appendix B, we give some examples that explain the proof methods.

## 2. RESULTS

Let  $p_1 < p_2$  and  $p_3$  be three prime numbers such that  $p_3 > p_1 \cdot p_2$  and  $p_3 \equiv \pm 2 \mod 3p_2$ . Throughout this paper, we denote

$$r_i := \text{rem}(p_i, p_1 \cdots p_{i-1}), \quad q_i := \text{quo}(p_i, p_1 \cdots p_{i-1})$$

where quo, rem of course, stand for quotient and remainder.

**THEOREM 2.1.** *Let  $3 < p_2 < p_3$  be odd prime numbers such that  $p_2 \equiv 1 \mod 3$  and  $p_3 > 3p_2$ . Then*

$$\text{hw}(\Phi_{3p_2 p_3}) = \begin{cases} N(p_3 - 2) + \left(\frac{4p_2-1}{3}\right), & \text{if } r_3 = 2 \\ N(p_3 + 2) - \left(\frac{4p_2-1}{3}\right), & \text{if } r_3 = 3p_2 - 2 \end{cases},$$

where  $N = \frac{7(p_2^2-1)}{9p_2}$ .

**EXAMPLE 2.2 (Toy Example).** In this example, we use small prime numbers  $p_2, p_3$  and use Theorem 2.1 to compute  $\text{hw}(\Phi_{3p_2 p_3})$ . Let  $p_2 = 7$ , here  $N = \frac{7 \cdot 48}{63} = \frac{16}{3}$ .

- Let  $p_3 = 23 = 1 \cdot 3 \cdot 7 + 2 \Rightarrow \text{hw}(\Phi_{3 \cdot 7 \cdot 23}(x)) = \frac{16}{3}(23 - 2) + \frac{27}{3} = 121$ .
- Let  $p_3 = 61 = 2 \cdot 3 \cdot 7 + 19 \Rightarrow \text{hw}(\Phi_{3 \cdot 7 \cdot 61}(x)) = \frac{16}{3}(61 + 2) + \frac{27}{3} = 327$ .

**EXAMPLE 2.3 (Big Example).** In this example, we consider larger values of  $p_2$  and  $p_3$  which needs more time and effort to compute  $\Phi_{3p_2 p_3}(x)$ . Let  $p_2 = 283$ , here  $N = \frac{186872}{849}$ . Let  $p_3 = 84916133 = 100019 \cdot 3 \cdot 283 + 2$ . Then using Theorem 2.1 we have

$$\text{hw}(\Phi_{3 \cdot 283 \cdot 84916133}(x)) = \frac{186872}{849}(84916133 - 2) + \frac{1131}{3} = 18690750945.$$

**THEOREM 2.4.** *Let  $3 < p_2 < p_3$  be odd prime numbers such that  $p_2 \equiv 2 \mod 3$  and  $p_3 > 3p_2$ . Then*

$$\text{hw}(\Phi_{3p_2 p_3}) = \begin{cases} N(p_3 - 2) + \frac{4p_2+1}{3}, & \text{if } r_3 = 2 \\ N(p_3 + 2) - \frac{4p_2+1}{3}, & \text{if } r_3 = 3p_2 - 2 \end{cases},$$

where  $N = \frac{(p_2+1)(7p_2-2)}{9p_2}$ .

**EXAMPLE 2.5 (Toy Example).** In this example, we use small prime numbers  $p_2, p_3$  and use Theorem 2.4 to compute  $\text{hw}(\Phi_{3p_2 p_3})$ . Let  $p_2 = 5$ , here  $N = \frac{(5+1) \cdot (5 \cdot 7 - 2)}{9 \cdot 5} = \frac{22}{5}$ .

- Let  $p_3 = 17 = 3 \cdot 3 \cdot 5 + 2 \Rightarrow \text{hw}(\Phi_{3 \cdot 5 \cdot 17}(x)) = \frac{22}{5}(17 - 2) + \frac{21}{3} = 73$ .
- Let  $p_3 = 43 = 2 \cdot 3 \cdot 5 + 13 \Rightarrow \text{hw}(\Phi_{3 \cdot 5 \cdot 13}(x)) = \frac{22}{5}(43 + 2) - \frac{21}{3} = 191$ .

## 3. PRELIMINARIES

In this section we review some needed properties of cyclotomic polynomials and give some important notation needed in the rest of the paper.

3.1. *Partition of Cyclotomic polynomials.* In [1, 4], a partition of cyclotomic polynomials was introduced, and also the following properties were given. This partition can be used to simplify studying several properties of cyclotomic polynomial

NOTATION 1 (Partition). *Let*

$$\begin{aligned}\Phi_{mp}(x) &= \sum_{i \geq 0} f_{m,p,i}(x) x^{ip} & \text{where } \deg f_{m,p,i}(x) < p \\ f_{m,p,i}(x) &= \sum_{j \geq 0} f_{m,p,i,j}(x) x^{jm} & \text{where } \deg f_{m,p,i,j}(x) < m\end{aligned}$$

NOTATION 2 (Operation). *For a polynomial  $f$  of degree less than  $m$ , let*

1.  $\mathcal{T}_s f = \text{rem}(f, x^s)$  “Truncate”
2.  $\mathcal{F}f = x^{m-1} f(x^{-1})$  “Flip”
3.  $\mathcal{R}_s f = \text{rem}(x^{m-\text{rem}(s,m)} f, x^m - 1)$  “Rotate”
4.  $\mathcal{E}_s f = f(x^{\text{rem}(s,m)})$  “Expand”

Throughout this paper, for an integer  $m$  and a prime  $p$ , we denote

$$r := \text{rem}(p, m), \quad q := \text{quo}(p, m)$$

where quo, of course, stand for quotient. The formula of the sub-polynomials  $f_{m,p,i,j}$  is given by the following theorem.

THEOREM 3.1 (Block). *For  $0 \leq i \leq \varphi(m) - 1$  and  $0 \leq j \leq q$ ,*

$$f_{m,p,i,j} = \begin{cases} -\mathcal{R}_{ir}(\Psi_m \cdot \mathcal{E}_r \mathcal{T}_{i+1} \Phi_m) & 0 \leq j \leq q-1 \\ \mathcal{T}_r f_{m,p,i,0} & j = q \end{cases},$$

where  $\Psi_m(x) = \frac{x^m - 1}{\Phi_m(x)}$ , the  $m$ -th inverse cyclotomic polynomial.

For more information and properties of  $\Psi_m(x)$  see [17].

NOTATION 3. *We will also use the following notation*

$$\Phi_m = \sum_{s \geq 0} a_s x^s \quad \Psi_m = \sum_{t \geq 0} b_t x^t \quad f_{m,p,i,0} = \sum_{k=0}^{m-1} c_k x^k$$

LEMMA 3.2 (Explicit expression for blocks). *Let  $p > m$ . Let  $r = \text{rem}(p, m)$ . Then the blocks  $f_{m,p,i,j}$  can be written explicitly as in*

$$\begin{aligned}1. \quad f_{m,p,i,j} &= \begin{cases} -\sum_{s=0}^i a_s \mathcal{R}_{(i-s)r} \Psi_m & \text{if } j < q \\ \mathcal{T}_r f_{m,p,i,0} & \text{if } j = q \end{cases} \\ 2. \quad c_k &= -\sum_{s=0}^i a_s b_{\text{rem}(k+(i-s)r, m)}.\end{aligned}$$

PROOF.

1. Note

$$\Phi_{mp} = \frac{\Phi_m(x^p)}{\Phi_m(x)} = -\Phi_m(x^p) \Psi_m \frac{1}{1-x^m} = -\Phi_m(x^p) \Psi_m \sum_{u \geq 0} x^{um} = \sum_{s \geq 0} x^{sp} \left( -a_s \Psi_m \sum_{u \geq 0} x^{um} \right)$$

Thus  $\Phi_{mp}$  is the sum of weighted-shifted  $\Psi_m$ , as illustrated by the following diagram.




FIGURE 1. weighted-shifted  $\Psi_m$

The claim is immediate from the  $f_{m,p,i,j}$  slice of the above diagram.

2. Note

$$f_{m,p,i,0} = - \sum_{s=0}^i a_s \mathcal{R}_{(i-s)r} \sum_{k \geq 0} b_k x^k = - \sum_{s=0}^i a_s \sum_{k \geq 0} b_{\text{rem}(k+(i-s)r,m)} x^k = \sum_{k \geq 0} \left( - \sum_{s=0}^i a_s b_{\text{rem}(k+(i-s)r,m)} \right) x^k$$

$$\text{Thus, } c_k = - \sum_{s=0}^i a_s b_{\text{rem}(k+(i-s)r,m)}$$

□

**THEOREM 3.3 (Intra-Structure).** *Within a cyclotomic polynomial, we have*

1. (Repetition)  $f_{m,p,i,0} = \dots = f_{m,p,i,q-1}$ .
2. (Truncation)  $f_{m,p,i,q} = \mathcal{T}_r f_{m,p,i,0}$ .
3. (Symmetry)  $f_{m,p,i',0} = \mathcal{R}_{\varphi(m)-1-r} \mathcal{F} f_{m,p,i,0}$  if  $i' + i = \varphi(m) - 1$ .

**THEOREM 3.4 (Inter-Structure).** *Among cyclotomic polynomials, we have*

1. (Invariance)  $f_{m,\tilde{p},i,0} = f_{m,p,i,0}$  if  $\tilde{p} - p \equiv 0 \pmod{m}$ .
2. (Semi-Invariance)  $f_{m,\tilde{p},i,0} = -\mathcal{R}_{\varphi(m)-1} \mathcal{F} f_{m,p,i,0}$  if  $\tilde{p} + p \equiv 0 \pmod{m}$ .

As an application of the properties above we have the following theorem.

**THEOREM 3.5 (Hamming weight).** *Let  $\text{hw}(f)$  stands for the number of non-zero terms in the polynomial  $f$ . Then we have*

1. [Linear]  $\text{hw}(\Phi_{mp}) = A \cdot p + B$ .
2. [Parallel]  $\text{hw}(\Phi_{m\tilde{p}}) = A \cdot \tilde{p} - B$ ,

where  $p + \tilde{p} \equiv 0 \pmod{m}$ .

In [3], another version of  $f_{m,p,i,0}$  was also given.

**3.2. Properties of  $\Phi_{3p_2p_3}$ .** In this subsection we introduce some properties of  $\Phi_{3p_2p_3}(x)$  to be used in the proofs of the main results.

**LEMMA 3.6.** *Let  $m = 3p_2, p_2 > 3$  be a prime number. Let  $\Phi_m = \sum_s a_s x^s$ . For  $0 \leq i \leq p_2 - 1$ , we have*

$$a_i = \begin{cases} 1, & \text{if } i \equiv 0 \pmod{3} \\ -1, & \text{if } i \equiv 1 \pmod{3} \\ 0, & \text{if } i \equiv 2 \pmod{3} \end{cases},$$

and for  $p_2 \leq i \leq \varphi(3p_2)$ , we have

$$a_i = \begin{cases} 0, & \text{if } i \equiv 0 \pmod{3} \\ -1, & \text{if } i \equiv 1 \pmod{3} \\ 1, & \text{if } i \equiv 2 \pmod{3} \end{cases}.$$

**PROOF.** Immediate from Notation 1, Theorem 3.1 or Lemma 3.2. We moved the proof into Appendix

A

□

LEMMA 3.7. For  $i = 1, \dots, p_2 - 2$ , we have

$$\mathcal{T}_{i+1}\Phi_{3p_2} = \begin{cases} (1-x) \sum_{j=0}^{\lfloor \frac{i-1}{3} \rfloor} x^{3j} + x^i, & \text{if } i \equiv 0 \pmod{3} \\ (1-x) \sum_{j=0}^{\lfloor \frac{i}{3} \rfloor} x^{3j}, & \text{if } i \equiv 1, 2 \pmod{3} \end{cases}.$$

PROOF. Immediate from Lemma 3.6.  $\square$

LEMMA 3.8. For  $i = p_2 - 1, \dots, \varphi(3p_2) - 2$  and  $p_2 \equiv 1 \pmod{3}$ , we have

$$\mathcal{T}_{i+1}\Phi_{3p_2} = (1-x) \sum_{j=0}^{q_2-1} x^{3j} + (1-x^2)x^{p_2-1} \begin{cases} \sum_{j=0}^{\lfloor \frac{i-p_2-1}{3} \rfloor} x^{3j} + x^i, & \text{if } i \equiv 0 \pmod{3} \\ \sum_{j=0}^{\lfloor \frac{i-p_2-1}{3} \rfloor} x^{3j} + x^{i-1}, & \text{if } i \equiv 1, 2 \pmod{3} \end{cases}.$$

PROOF. Immediate from Lemma 3.6.  $\square$

PROPOSITION 3.9. For  $0 \leq i \leq \varphi(3p_2) - 1$  we have

$$\begin{aligned} f_{m,p,i,0} &= \mathcal{N}_{\text{rem}(2i,3p_2)}(\Psi_{3p_2}\mathcal{E}_2\mathcal{T}_{i+1}\Phi_{3p_2}) \\ &= -\mathcal{R}_{\text{rem}(2i,3p_2)}(\Psi_{3p_2}\mathcal{E}_2\mathcal{T}_{i+1}\Phi_{3p_2}) \\ &= -\text{rem}\left(x^{3p_2-\text{rem}(2i,3p_2)}C_i, x^{3p_2}-1\right), \end{aligned}$$

$$\text{where } C_i = \Psi_{3p_2} \begin{cases} (x^2-1) \sum_{j=0}^{\lfloor \frac{i-1}{3} \rfloor} x^{6j} - x^{2i}, & \text{if } i \equiv 0 \pmod{3} \\ (x^2-1) \sum_{j=0}^{\lfloor \frac{i}{3} \rfloor} x^{6j}, & \text{if } i \equiv 1, 2 \pmod{3} \end{cases}$$

if  $i < p_2 - 1 \Rightarrow 2i = 2p_2 - 2 < 3p_2 = m \Rightarrow \text{rem}(2i, 3p_2) = 2i$ .

REMARK 3.10. The number of terms will not be changed with rotation and negation, so in order to study the number of terms of  $f_{m,p,i,0}$  it is enough to study the number of terms of  $C_i$ .

#### 4. PROOFS

In this section we will prove the results of this paper.

4.1. *Proof of Theorem 2* ( $r_2 = 1$ ). In this subsection we assume that  $r_2 = 1$  that is  $p_2 \equiv 1 \pmod{3}$ , and  $r_3 \equiv 2 \pmod{3p_2}$ . In the following six Lemmas we will compute  $\text{hw}(f_{3p_2,p_3,i,0})$  for several values of  $i$ , in order to be used later in proving the main result.

LEMMA 4.1. Let  $i = 3u + v < p_2 - 1$ , where  $v \in \{1, 2\}$ . Then

$$\text{hw}(f_{3p_2,p_3,i,0}) = \begin{cases} 8(u+1), & \text{if } u = 0, 1, \dots, \frac{q_2}{2} - 1 \\ 4(u+1 + \frac{q_2}{2}), & \text{if } u = \frac{q_2}{2}, \dots, q_2 - 1 \end{cases}.$$

PROOF. As stated in Remark 3.10 we have,

$$\begin{aligned}
\text{hw}(f_{3p_2,p,i,0}) &= \text{hw}(C_i) \\
&= \text{hw} \left( \Psi_{3p_2} \cdot (x^2 - 1) \cdot \sum_{j=0}^{\lfloor \frac{i}{3} \rfloor} x^{6j} \right) && \text{by Lemma 3.7} \\
&= \text{hw} \left( \Phi_3(x) \cdot (x^{p_2} - 1)(x^2 - 1) \sum_{j=0}^{\lfloor \frac{i}{3} \rfloor} x^{6j} \right) \\
&= \text{hw} \left( (1+x)(x^{p_2} - 1)(x^3 - 1) \sum_{j=0}^{\lfloor \frac{i}{3} \rfloor} x^{6j} \right) && \Phi_3(x) \cdot (x^2 - 1) = (x+1)(x^3 - 1) \\
&= \text{hw} \left( (1+x)(1-x^3) \left( \sum_{j=0}^u x^{6j} - \sum_{j=0}^u x^{6j+p_2} \right) \right) \\
&= 2 \cdot \text{hw} \left( \sum_{j=0}^u x^{6j} - \sum_{j=0}^u x^{6j+3} - \sum_{j=0}^u x^{6j+p_2} + \sum_{j=0}^u x^{6j+p_2+3} \right).
\end{aligned}$$

If  $u \leq \frac{q_2}{2} - 1$ , then there isn't any cancellation in the above sum, thus

$$\text{hw}(f_{3p_2,p_3,i,0}) = 2 \cdot 4 \cdot (u+1) = 8(u+1)$$

as desired. On the other hand, if  $\frac{q_2}{2} \leq u \leq q_2 - 1$ , then we have

$$\begin{aligned}
\text{hw}(f_{3p_2,p_3,i,0}) &= \text{hw} \left( (x+1)(1-x^{p_2}) \sum_{j=0}^u x^{6j} + (x+1)x^3(x^{p_2} - 1) \sum_{j=0}^u x^{6j} \right) \\
&= \text{hw} \left( (x+1)(1-x^{p_2}) \sum_{j=0}^u x^{6j} \right) + \text{hw} \left( x^3(x+1)(x^{p_2} - 1) \sum_{j=0}^u x^{6j} \right) \\
&= 2 \cdot \text{hw} \left( (x+1)(1-x^{p_2}) \sum_{j=0}^u x^{6j} \right) \\
&= 2 \cdot (\text{hw}(A_i) + \text{hw}(B_i)).
\end{aligned}$$

Where  $A_i \equiv A_u = (1-x^{p_2+1}) \sum_{j=0}^u x^{6j}$  and  $B_i \equiv B_u = (x-x^{p_2}) \sum_{j=0}^u x^{6j}$ . We will study  $A_i$  and  $B_i$  for the case  $r_2 = 1$ . we claim that  $\text{hw}(A_i) = 2(u+1)$  and  $\text{hw}(B_i) = q_2$ . For  $A_u$ , if there is any cancelation in the sum, then  $p_2 + 1 + 6j = 6k$  for some integers  $j$  and  $k$ . Thus,  $p_2 + 1 \equiv 0 \pmod{3}$  which contradicts the fact that  $p_2 \equiv 1 \pmod{3}$ . Now, we claim that

$$B_u = \sum_{j=0}^{\frac{q_2}{2}-1} x^{6j+1} - x^{p_2+6(1+j+u-\frac{q_2}{2})}$$

We prove the claim by induction on  $u$  starting from  $u = \frac{q_2}{2}$ .

- If  $u = \frac{q_2}{2}$ , then

$$\begin{aligned}
B_u &= (x - x^{p_2}) \sum_{j=0}^{\frac{q_2}{2}} x^{6j} \\
&= x + x^7 + \dots + x^{3q_2+1} - x^{p_2} - x^{p_2+6} - \dots - x^{p_2+3q_2} \\
&= \sum_{j=0}^{\frac{q_2}{2}-1} x^{6j+1} - x^{p_2+6(1+j)}
\end{aligned}$$

- Assume that  $B_u = \sum_{j=0}^{\frac{q_2}{2}-1} x^{6j+1} - x^{p_2+6(1+j+u-\frac{q_2}{2})}$
- Consider

$$\begin{aligned}
B_{u+1} &= (x - x^{p_2}) \sum_{j=0}^{u+1} x^{6j} \\
&= (x - x^{p_2}) \sum_{j=0}^u x^{6j} + x^{6u+6}(x - x^{p_2}) \\
&= \sum_{j=0}^{\frac{q_2}{2}-1} (x^{6j+1} - x^{p_2+6(1+j+u-\frac{q_2}{2})}) + (x^{6u+7} - x^{p_2+6u+6}) \\
&= \sum_{j=0}^{\frac{q_2}{2}-1} x^{6j+1} - \sum_{j=1}^{\frac{q_2}{2}-1} x^{p_2+6(1+j+u-\frac{q_2}{2})} - x^{6u+7} + (x^{6u+7} - x^{p_2+6u+6}) \quad \text{since } p_2 - 3q_2 = 1 \\
&= \sum_{j=0}^{\frac{q_2}{2}-1} x^{6j+1} - \sum_{j=1}^{\frac{q_2}{2}-1} x^{p_2+6(1+j+u-\frac{q_2}{2})} - x^{p_2+6u+6} \\
&= \sum_{j=0}^{\frac{q_2}{2}-1} x^{6j+1} - \sum_{j=0}^{\frac{q_2}{2}-1} x^{p_2+6(2+j+u-\frac{q_2}{2})} \quad \text{by reindexing}
\end{aligned}$$

by induction we have  $B_u = \sum_{j=0}^{\frac{q_2}{2}-1} x^{6j+1} - x^{p_2+6(1+j+u-\frac{q_2}{2})}$  as desired. Thus  $\text{hw}(f_{3p_2,p_3,0}) = 4(u+1+\frac{q_2}{2})$ , for  $\frac{q_2}{2} \leq u \leq q_2 - 1$ .  $\square$

LEMMA 4.2. If  $i = 3u + v < p_2 - 1$  where  $v \in \{1, 2\}$ , then

$$\text{hw}(f_{3p_2,p,i,q}) = \begin{cases} 1, & \text{if } u = 0, 1, \dots, \frac{q_2}{2} - 1 \\ 3 - v, & \text{if } \frac{q_2}{2} \leq u \leq q_2 - 1 \end{cases}.$$

PROOF. It is enough to show that **only one** of the terms  $x^0$  and  $x^1$  will appear in  $f_{3p_2,p,i,0}$  when  $0 \leq u \leq \frac{q_2}{2} - 1$ , and **only one** of the terms  $x^0$  and  $x$  will appear in  $f_{3p_2,p,i,0}$  when  $\frac{q_2}{2} \leq u \leq q_2$  if  $v = 2$ , while both of them will appear when  $v = 1$ . In the table below we list the terms that appear in  $C_i = \Psi_{3p_2} \cdot (x^2 - 1) \cdot \sum_{j=0}^u x^{6j}$  and the corresponding terms in  $f_{3p_2,p,i,0} = \text{rem}(x^{-2i}C_i, x^{3p_2} - 1)$ .

$u/\text{Terms in}$	$C_i$ if $v = 1$	$f_{3p_2,p,i,0}$ if $v = 1$	$C_i$ if $v = 2$	$f_{3p_2,p,i,0}$ if $v = 2$
$0 \leq u \leq \frac{q_2}{2} - 1$	$-x^{6u+2v+1}$	$-x$	$-x^{6u+2v}$	$-1$
$\frac{q_2}{2} \leq u \leq q_2 - 1$	$-x^{6u+2v} - x^{6u+2v+1}$	$-1 - x$	$x^{6u+2v+1}$	$x$

For the case  $0 \leq u \leq \frac{q_2}{2} - 1$ , recall the proof of Lemma 4.1, we have

$$C_i = (1 + x)(1 - x^3) \left( \sum_{j=0}^u x^{6j} - \sum_{j=0}^u x^{6j+p_2} \right)$$



So it is clear that  $x^{6u+2v}$  will not appear when  $v = 1$ , while  $x^{6u+2v+1}$  will not appear when  $v = 2$ . Now for the case  $u > \frac{q_2}{2} - 1$ , again from  $C_i = (1+x)(1-x^3) \left( \sum_{j=0}^u x^{6j} - \sum_{j=0}^u x^{6j+p_2} \right)$ . When  $v = 1$  the terms  $-x^{6u+3} - x^{p_2+6(u-\frac{q_2}{2})+1} = -x^{6u+2v} - x^{6u+2v}$  will appear. Finally, when  $v = 2$  we have only the term  $x^{p_2+6(u-\frac{q_2}{2})+4} = x^{6u+5} = x^{6u+2v+1}$ .  $\square$

LEMMA 4.3. *If  $i = 3u < p_2 - 1$ , then*

$$\text{hw}(f_{3p_2, p_3, i, 0}) = \begin{cases} 8u + 6, & \text{if } u = 0, 1, \dots, \frac{q_2}{2} - 1 \\ 4u + 2q_2 + 2, & \text{if } u = \frac{q_2}{2}, \dots, q_2 \end{cases}.$$

PROOF. The case when  $u = 0$  is trivial. Consider  $0 < u \leq q_2$ , in this case the cyclotomic polynomial  $\Phi_{3p_2p_3}$  is flat (see Theorem 38 in [10]), so we will only worry about cancelations. From Remark 3.10 we have

$$\begin{aligned} \text{hw}(f_{3p_2, p_3, i, 0}) &= \text{hw}(C_i) \\ &= \text{hw} \left( \Psi_{3p_2} \cdot \left( (x^2 - 1) \cdot \sum_{j=0}^{\lceil \frac{i-1}{3} \rceil} x^{6j} - x^{2i} \right) \right) \\ &= \text{hw} \left( \Psi_{3p_2} \cdot \left( (x^2 - 1) \cdot \sum_{j=0}^{\lceil u-\frac{1}{3} \rceil} x^{6j} - x^{2i} \right) \right) \\ &= \text{hw} \left( \Psi_{3p_2} \cdot (x^2 - 1) \cdot \sum_{j=0}^{u-1} x^{6j} - x^{6u} \Psi_{3p_2} \right). \end{aligned}$$

We handle two cases:

- $1 \leq u \leq \frac{q_2}{2} - 1$ : In this case there are no cancelations between the sums  $S_1 = \Psi_{3p_2} \cdot (x^2 - 1) \cdot \sum_{j=0}^{u-1} x^{6j}$  and  $S_2 = x^{6u} \Psi_{3p_2}$ . Notice that  $S_1 = (1 - x^{p_2})(1 + x - x^3 - x^4) \sum_{j=0}^{u-1} x^{6j}$  and  $S_2 = \Phi_3 \cdot (x^{p_2+6u} - x^{6u})$ . Thus  $\text{hw}(f_{3p_2, p_3, i, 0}) = 8u + 6$ .
- $u \geq \frac{q_2}{2}$ : Notice that  $\Psi_{3p_2} \cdot (x^2 - 1) = (x+1)(x^3 - 1)(1 - x^{p_2})$ , so we have

$$\text{hw}(f_{3p_2, p_3, i, 0}) = \text{hw}((x+1)A_u + B_u)$$

where  $A_u = (x^3 - 1)(1 - x^{p_2}) \sum_{j=0}^{u-1} x^{6j}$  and  $B_u = x^{6u} \Psi_{3p_2}$ . We will use mathematical induction on  $u$  to prove that

$$(x+1)A_u = (x+1) \sum_{j=0}^{q_2-1} (-1)^{j+1} x^{3j} + (1-x^2) \sum_{j=q_2}^{2u-1} (-1)^{j+1} x^{3j} + (x+1) \sum_{j=2u}^{2u+q_2-1} (-1)^j x^{3j+1}$$

– For  $u = \frac{q_2}{2}$  :

$$\begin{aligned} (x+1)A_{\frac{q_2}{2}} &= (x+1)(x^3 - 1)(1 - x^{p_2}) \sum_{j=0}^{\frac{q_2-2}{2}} x^{6j} \\ &= (x+1)(1 - x^{p_2}) \sum_{j=0}^{q_2-1} (-1)^{j+1} x^{3j} \\ &= (x+1) \left( \sum_{j=0}^{q_2-1} (-1)^{j+1} x^{3j} + \sum_{j=0}^{q_2-1} (-1)^j x^{3j+p_2} \right) \\ &= (x+1) \left( \sum_{j=0}^{q_2-1} (-1)^{j+1} x^{3j} + \sum_{j=q_2}^{2q_2-1} (-1)^j x^{3j+1} \right) \end{aligned}$$

– Assume that

$$(x+1)A_u = (x+1) \sum_{j=0}^{q_2-1} (-1)^{j+1} x^{3j} + (1-x^2) \sum_{j=q_2}^{2u-1} (-1)^{j+1} x^{3j} + (x+1) \sum_{j=2u}^{2u+q_2-1} (-1)^j x^{3j+1}$$

– Consider

$$\begin{aligned} (x+1)A_{u+1} &= (x+1)(x^3-1)(1-x^{p_2}) \sum_{j=0}^u x^{6j} \\ &= (x+1)(x^3-1)(1-x^{p_2}) \sum_{j=0}^{u-1} x^{6j} + (x+1)(x^3-1)(1-x^{p_2})x^{6u} \\ &= (x+1) \sum_{j=0}^{q_2-1} (-1)^{j+1} x^{3j} + (1-x^2) \sum_{j=q_2}^{2u-1} (-1)^{j+1} x^{3j} \\ &\quad + (x+1) \sum_{j=2u}^{2u+q_2-1} (-1)^j x^{3j+1} \quad \text{by induction} \\ &\quad + (-x^{6u} - x^{6u+1} + x^{6u+3} + x^{6u+4} + x^{6u+p_2} + x^{6u+p_2+1} - x^{6u+p_2+3} - x^{6u+p_2+4}) \\ &= (x+1) \sum_{j=0}^{q_2-1} (-1)^{j+1} x^{3j} + (1-x^2) \sum_{j=q_2}^{2u+1} (-1)^{j+1} x^{3j} \\ &\quad + (x+1) \sum_{j=2u+2}^{2u+q_2+1} (-1)^j x^{3j+1} \end{aligned}$$

as desired.

Now, it remains to find  $\text{hw}(f_{3p_2,p,i,0}) = \text{hw}((x+1)A_u + B_u)$ . Notice that there are two cancelations between  $B_u$  and the third sum from  $(x+1)A_u$ , namely  $x^{6u+1}$  and  $x^{6u+2}$ . Thus  $\text{hw}(f_{3p_2,p,i,0}) = \text{hw}((x+1)A_u + B_u)$  equals the sum of hamming weights of the three sums of  $(x+1)A_u$  plus 2. so  $\text{hw}((x+1)A_u + B_u) = 2 + 2q_2 + 2(2u - q_2) + 2q_2 = 4u + 2q_2 + 2$ .

□

LEMMA 4.4. *If  $i = 3u$ , then*

$$\text{hw}(f_{3p_2,p,i,q}) = \begin{cases} 2, & \text{if } u = 0, 1, \dots, \frac{q_2}{2} - 1 \\ 1, & \text{if } u = \frac{q_2}{2}, \dots, q_2 \end{cases}.$$

PROOF. It is enough to prove that both the terms  $x^0$  and  $x$  will appear in  $f_{3p_2,p,i,0}$  when  $0 \leq u \leq \frac{q_2}{2} - 1$  and only one of the terms  $x^0$  and  $x$  will appear in  $f_{3p_2,p,i,0}$  when  $\frac{q_2}{2} \leq u \leq q_2$ . In the table below we list the term appears in  $C_i = \Psi_{3p_2} \cdot (x^2 - 1) \cdot \sum_{j=0}^{u-1} x^{6j} - x^{6u} \Psi_{3p_2}$  and the corresponding term in  $f_{3p_2,p,i,0}$ .

$u$	Terms in $C_i$	Terms in $f_{3p_2,p,i,0}$
$0 \leq u \leq \frac{q_2}{2} - 1$	$x^{6u} + x^{6u+1}$	$1 + x$
$\frac{q_2}{2} \leq u \leq q_2$	$x^{6u}$	$1$

The reasoning is clear from the proof of Lemma 4.3.

□

$$\text{LEMMA 4.5. } \text{hw}(f_{3p_2,p_3,p_2-1,0}) = \begin{cases} 2(p_2 - 1), & \text{if } r_2 = 1 \\ 2p_2 + 1, & \text{if } r_2 = 2 \end{cases}.$$

PROOF. Immediate using Remark 3.10 and analogous arguments in the proofs of other lemmas.

□

LEMMA 4.6. *If  $i \geq p_2$ , then  $\text{hw}(f_{3p_2,p,i,q}) = 0$*

PROOF. From Lemma 3.2-2 we know that

$$c_k = - \sum_{s=0}^i a_s b_{\text{rem}(k+2(i-s), 3p_2)}$$

and for  $\Psi_{3p_2}$

$$b_k = \begin{cases} -1 & k = 0, 1, 2 \\ 1 & k = p_2, p_2 + 1, p_2 + 2 \\ 0 & \text{otherwise} \end{cases}$$

We will compute  $c_0$  and  $c_1$  for each  $i \geq p_2$ . At first

$$\begin{aligned} c_0 &= - \sum_{s=0}^i a_s b_{2(i-s)} \\ &= -(a_{i-\frac{p_2+1}{2}} b_{p_2+1} + a_{i-1} b_2 + a_i b_0) \quad \text{the only possible values for } k \text{ to be even and } b_k \neq 0 \\ &= -(a_{i-\frac{p_2+1}{2}} - a_{i-1} - a_i) \\ &= 0 \end{aligned}$$

we summarise the last conclusion in the next table, we use Lemma 3.6 and the fact that  $i \equiv i - \frac{p_2+1}{2} \pmod{3}$  and  $i > p_2$ .

$i \pmod{3}$	$a_i$	$a_{i-1}$	$a_{i-\frac{p_2+1}{2}}$	$a_{i-\frac{p_2+1}{2}} - a_{i-1} - a_i$
0	0	1	1	0
1	-1	0	-1	0
2	1	-1	0	0

For  $i = p_2$  and  $p_2 \equiv 1 \pmod{3}$  we have  $a_{i-\frac{p_2+1}{2}} - a_{i-1} - a_i = -1 - 0 + 1 = 0$  and if  $p_2 \equiv 2 \pmod{3}$  we have  $a_{i-\frac{p_2+1}{2}} - a_{i-1} - a_i = 0 + 1 - 1 = 0$ . So from the discussion above we see that  $c_0 = 0$ . It remains to prove that  $c_1 = 0$  which can be done using similar ideas so we skip the proof of that part.

□

THEOREM 4.7. *Let  $3 < p_2 < p_3$  be odd prime numbers such that  $p_2 \equiv 1 \pmod{3}$ . Then*

$$\text{hw}(\Phi_{3p_2p_3}) = \begin{cases} N(p_3 - 2) + \left(\frac{4p_2-1}{3}\right), & \text{if } r_3 = 2 \\ N(p_3 + 2) - \left(\frac{4p_2+1}{3}\right), & \text{if } r_3 = 3p_2 - 2 \end{cases},$$

where  $N = \frac{7(p_2^2-1)}{9p_2^2}$ .

PROOF. For  $r_3 = 2$ , note

$$\begin{aligned}
\text{hw}(\Phi_{3p_2p_3}) &= \sum_{0 \leq i \leq \varphi(m)-1} \text{hw}(f_{m,p,i}) x^{ip_3} \\
&= q_3 \sum_{i=0}^{\varphi(m)-1} \text{hw}(f_{m,p,i,0}) + \sum_{i=0}^{\varphi(m)-1} \text{hw}(f_{m,p,i,q}) \\
&= q_3 \sum_{i=0}^{\varphi(m)-1} \text{hw}(f_{m,p,i,0}) + \sum_{i=0}^{p_2-1} \text{hw}(f_{m,p,i,q}) && \text{by Lemma 4.6} \\
&= q_3 \sum_{i=0}^{\varphi(m)-1} \text{hw}(f_{m,p,i,0}) + 4q_2 + 1 && \text{by Lemma 4.2 and Lemma 4.4} \\
&= 2q_3 \sum_{i=0}^{p_2-2} \text{hw}(f_{m,p,i,0}) + 4q_2 + 1 && \text{by Symmetry} \\
&= 2q \left( \sum_{u=0}^{\frac{q_2^2-1}{2}} (24u+22) + \sum_{u=\frac{q_2^2}{2}}^{q_2-1} (12u+10+6q_2) \right) + 4q_2 + 1 && \text{by Lemma 4.3 and Lemma 4.1} \\
&= 2q_3(q_2(6q_2+7+\frac{9}{2}q_2)) + 4q_2 + 1 \\
&= 2q_3(q_2(7+\frac{21}{2}q_2)) + 4q_2 + 1 \\
&= 2q_3\left(\frac{7}{2}q_2(3q_2+2)\right) + 4q_2 + 1 \\
&= \frac{7(p_3-2)}{9p_2}(p_2-1)(p_2+1) + 4\frac{p_2-1}{3} + 1 && q_3 = \frac{p_3-2}{3p_2}, q_2 = \frac{p_2-1}{3} \\
&= (p_3-2)\frac{7(p_2^2-1)}{9p_2} + \frac{4p_2-1}{3} = N(p_3-2) + \frac{4p_2-1}{3}
\end{aligned}$$

Now for  $r_3 = 3p_2 - 2$ , from Theorem 3.5(Theorem 6.1 in [1]),

$$\text{hw}(\Phi_{3p_2p_3}) = Np_3 - \left(\frac{4p_2-1}{3} - 2N\right) = N(p_3+2) - \left(\frac{4p_2-1}{3}\right)$$

as desired. □

#### 4.2. Proof of Theorem 2.4 ( $r_2 = 2$ ).

LEMMA 4.8. *If  $i = 3u + v < p_2 - 1$  where  $v = 1, 2$ , then*

$$\text{hw}(f_{3p_2,p_3,i,0}) = \begin{cases} 8(u+1), & \text{if } 0 \leq u \leq \frac{q_2-1}{2} \\ 4(u+1+\frac{q_2+1}{2}), & \text{if } \frac{q_2+1}{2} \leq u \leq q_2-1 \end{cases}.$$

PROOF. As stated in Remark 3.10 we have

$$\begin{aligned}
\text{hw}(f_{3p_2,p,i,0}) &= \text{hw}(C_i) \\
&= \text{hw} \left( \Psi_{3p_2} \cdot (x^2 - 1) \cdot \sum_{j=0}^{\lfloor \frac{i}{3} \rfloor} x^{6j} \right) && \text{Lemma 3.7} \\
&= \text{hw} \left( \Phi_3(x) \cdot (x^{p_2} - 1)(x^2 - 1) \sum_{j=0}^{\lfloor \frac{i}{3} \rfloor} x^{6j} \right) \\
&= \text{hw} \left( (1+x)(x^{p_2} - 1)(x^3 - 1) \sum_{j=0}^{\lfloor \frac{i}{3} \rfloor} x^{6j} \right) && \Phi_3(x) \cdot (x^2 - 1) = (x+1)(x^3 - 1) \\
&= \text{hw} \left( (1+x)(1-x^3) \left( \sum_{j=0}^u x^{6j} - \sum_{j=0}^u x^{6j+p_2} \right) \right) \\
&= 2 \cdot \text{hw} \left( \sum_{j=0}^u x^{6j} - \sum_{j=0}^u x^{6j+3} - \sum_{j=0}^u x^{6j+p_2} + \sum_{j=0}^u x^{6j+p_2+3} \right)
\end{aligned}$$

If  $u \leq \frac{q_2-1}{2}$ , then there is no cancellation in the sum above, thus  $\text{hw}(f_{3p_2,p_3,i,0}) = 2 \cdot 4 \cdot (u+1) = 8(u+1)$  as desired. On the other hand, if  $u > \lceil \frac{q_2}{2} \rceil$ , then we have

$$\begin{aligned}
\text{hw}(f_{3p_2,p_3,i,0}) &= \text{hw} \left( (x+1)(1-x^{p_2}) \sum_{j=0}^u x^{6j} + (x+1)x^3(x^{p_2} - 1) \sum_{j=0}^u x^{6j} \right) \\
&= \text{hw} \left( (x+1)(1-x^{p_2}) \sum_{j=0}^u x^{6j} \right) + \text{hw} \left( x^3(x+1)(x^{p_2} - 1) \sum_{j=0}^u x^{6j} \right) \\
&= 2 \cdot \text{hw} \left( (x+1)(1-x^{p_2}) \sum_{j=0}^u x^{6j} \right) \\
&= 2 \cdot (\text{hw}(A_i) + \text{hw}(B_i))
\end{aligned}$$

Where  $A_i = A_u = (1-x^{p_2+1}) \sum_{j=0}^u x^{6j}$  and  $B_i = B_u = (x-x^{p_2}) \sum_{j=0}^u x^{6j}$ . For  $B_u = (x-x^{p_2}) \sum_{j=0}^u x^{6j}$  there is no cancelation in the sum and we have  $\text{hw}(B_u) = 2(u+1)$ . Assume by contrary that there is a cancelation, then  $6j+1 = 6k+p_2$  for some  $k$  and  $j$  which gives us  $p_2-1 \equiv 0 \pmod{3}$  a contradiction to the fact that  $p_2 \equiv 2 \pmod{3}$ .

For  $A_u = (1-x^{p_2+1}) \sum_{j=0}^u x^{6j}$  we claim that  $A_u = \sum_{j=0}^{\frac{q_2-1}{2}} x^{6j} - x^{p_2+1+6(u-\frac{q_2-1}{2}+j)}$ . We prove the claim by induction on  $u$  starting from  $u = \frac{q_2+1}{2}$ .

- If  $u = \frac{q_2+1}{2}$ , then

$$\begin{aligned}
A_{\frac{q_2+1}{2}} &= (1-x^{p_2+1}) \sum_{j=0}^{\frac{q_2+1}{2}} x^{6j} \\
&= (1+x^6 + \dots + x^{3q_2+3}) - (x^{p_2+1} + x^{p_2+7} + \dots + x^{2p_2+2}) \\
&= (1+x^6 + \dots + x^{3q_2-3}) - (x^{p_2+7} + \dots + x^{2p_2+2}) \\
&= \sum_{j=0}^{\frac{q_2-1}{2}} x^{6j} - x^{p_2+1+6(1+j)}
\end{aligned}$$

- Assume that  $A_u = \sum_{j=0}^{\frac{q_2-1}{2}} x^{6j} - x^{p_2+1+6(u-\frac{q_2-1}{2}+j)}$
- Consider

$$\begin{aligned}
A_{u+1} &= (1 - x^{p_2+1}) \sum_{j=0}^{u+1} x^{6j} \\
&= \sum_{j=0}^{\frac{q_2-1}{2}} x^{6j} - x^{p_2+1+6(u-\frac{q_2-1}{2}+j)} + x^{6u+6}(1 - x^{p_2+1}) \\
&= \sum_{j=0}^{\frac{q_2-1}{2}} x^{6j} - \sum_{j=1}^{\frac{q_2-1}{2}} x^{p_2+1+6(u-\frac{q_2-1}{2}+j)} - x^{6u+6} + x^{6u+6} - x^{p_2+1+6(u+1)} \\
&= \sum_{j=0}^{\frac{q_2-1}{2}} x^{6j} - \sum_{j=0}^{\frac{q_2+1}{2}} x^{p_2+1+6(u+1-\frac{q_2-1}{2}+j)} \\
&= \sum_{j=0}^{\frac{q_2-1}{2}} x^{6j} - \sum_{j=0}^{\frac{q_2-1}{2}} x^{p_2+1+6(u-\frac{q_2-1}{2}+j)} \quad \text{by reindexing}
\end{aligned}$$

Thus we have  $\text{hw}(f_{3p_2,p_3,0}) = 4(u+1 + \lceil \frac{q_2}{2} \rceil)$ , for  $\lceil \frac{q_2}{2} \rceil \leq u \leq q_2 - 1$ .  $\square$

LEMMA 4.9. If  $i = 3u < p_2 - 1$  and  $p_2 \equiv 2 \pmod{3}$ , then

$$\text{hw}(f_{3p_2,p,i,0}) = \begin{cases} 6 + 8u, & \text{if } 0 \leq u \leq \frac{q_2-1}{2} \\ 4u + 5 + 2q_2, & \text{if } \frac{q_2+1}{2} \leq u \leq q_2 - 1 \end{cases}$$

PROOF. In this case the cyclotomic polynomial  $\Phi_{3p_2p_3}$  may have some non flat coefficients, so we will worry about cancelations and overlapping.

$$\begin{aligned}
\text{hw}(f_{3p_2,p,i,0}) &= \text{hw} \left( \Psi_{3p_2} \cdot ((x^2 - 1) \sum_{j=0}^{\lfloor \frac{i-1}{3} \rfloor} x^{6j} - x^{2i}) \right) \quad \text{from Lemma 3.7} \\
&= \text{hw} \left( (1+x)(x^{p_2} - 1)(x^3 - 1) \sum_{j=0}^{\lfloor \frac{i-1}{3} \rfloor} x^{6j} - x^{2i} \Psi_{3p_2} \right) \\
&= \text{hw} \left( (1+x)(x^{p_2} - 1)(x^3 - 1) \sum_{j=0}^{u-1} x^{6j} - x^{6u} \Psi_{3p_2} \right)
\end{aligned}$$

We have the following cases

1.  $u \leq \frac{q_2-1}{2}$ , in this case there are no cancelations or overlapping in the above sum. Thus  $\text{hw}(f_{3p_2,p,i,0}) = 8u + 6$
2.  $u \geq \frac{q_2+1}{2}$ . Notice that  $\Psi_{3p_2} \cdot (x^2 - 1) = (x+1)(x^3 - 1)(1 - x^{p_2})$ , so we have

$$\text{hw}(f_{3p_2,p,i,0}) = \text{hw}((x+1)A_u + B_u)$$

where  $A_u = (x^3 - 1)(1 - x^{p_2}) \sum_{j=0}^{u-1} x^{6j}$  and  $B_u = x^{6u} \Psi_{3p_2}$ . We claim that

$$(x+1)A_u = (x+1) \sum_{j=0}^{q_2} (-1)^{j+1} x^{3j} + (1-x^2) \sum_{j=q_2}^{2u-2} (-1)^{j+1} x^{3j+2} + (x+1) \sum_{j=2u-1}^{2u+q_2-1} (-1)^j x^{3j+2}$$

The proof of this claim is similar to the one in the proof of Lemma 4.3, so we omit it. Now since there is a cancelation of the term  $x^{6u+2}$  between  $B_u$  and  $(x+1)A_u$  and an overlapping between the term

$x^{6u}$  between  $B_u$  and  $(x+1)A_u$ , we have  $\text{hw}(f_{3p_2,p,i,0}) = \text{hw}((x+1)A_u + B_u) = 2(q_2 + 1) + 2(2u - 1 - q_2) + 2(q_2 + 1) - 1 + 4 = 4u + 2q_2 + 5$ .

□

LEMMA 4.10. *If  $i = 3u + v < p_2 - 1$  where  $v = 1, 2$  and  $p_2 \equiv 2 \pmod{3}$ , then*

$$\text{hw}(f_{3p_2,p,i,q}) = \begin{cases} 1, & \text{if } 0 \leq u \leq \frac{q_2-3}{2} \\ v, & \text{if } \frac{q_2-1}{2} \leq u \end{cases}.$$

PROOF. Similar to the proof of Lemma 4.2 so we move it to the appendix. □

THEOREM 4.11. *Let  $3 < p_2 < p_3$  be odd prime numbers such that  $p_2 \equiv 2 \pmod{3}$ . Then*

$$\text{hw}(\Phi_{3p_2p_3}) = \begin{cases} N(p_3 - 2) + \frac{4(p_2+1)}{3}, & \text{if } r_3 = 2 \\ N(p_3 + 2) - \frac{4(p_2+1)}{3}, & \text{if } r_3 = p_2 - 2 \end{cases}$$

where  $N = \frac{(p_2+1)(7p_2-2)}{9p_2}$ .

PROOF. Note if  $r_2 = r_3 = 2$ , then

$$\begin{aligned} \text{hw}(\Phi_{3p_2p_3}) &= \sum_{0 \leq i \leq \varphi(m)-1} \text{hw}(f_{m,p,i}) x^{ip_3} \\ &= q_3 \sum_{i=0}^{\varphi(m)-1} \text{hw}(f_{m,p,i,0}) + \sum_{i=0}^{\varphi(m)-1} \text{hw}(f_{m,p,i,q}) \\ &= q_3 \sum_{i=0}^{\varphi(m)-1} \text{hw}(f_{m,p,i,0}) + \sum_{i=0}^{p_2-1} \text{hw}(f_{m,p,i,q}) && \text{by Lemma 4.6} \\ &= q_3 \sum_{i=0}^{\varphi(m)-1} \text{hw}(f_{m,p,i,0}) + 4q_2 + 3 && \text{by Lemma 4.10} \\ &= 2q_3 \sum_{i=0}^{p_2-2} \text{hw}(f_{m,p,i,0}) + 4q_2 + 3 && \text{by Symmetry} \\ &= 2q_3 \left( \sum_{u=0}^{\frac{q_2-1}{2}} (24u + 22) \right) + 4q_2 + 3 && \text{by Lemma 4.8 and 4.9} \\ &+ 2q_3 \left( \sum_{u=\frac{q_2+1}{2}}^{q_2-1} (12u + 6q_2 + 17) \right) + 2q_3(2p_2 + 1) && \text{by Lemma 4.5} \\ &= 2q_3 \left( \frac{21}{2}q_2^2 + \frac{21}{2}q_2 + 1 \right) + 2q_3(2p_2 + 1) + 4q_2 + 3 \\ &= 21q_2q_3(q_2 + 1) + 2q_3(2p_2 + 2) + 4q_2 + 3 \\ &= 7 \frac{p_3-2}{p_2} \cdot \frac{p_2+1}{3} \cdot \frac{p_2-2}{3} + 4 \frac{p_3-2}{3p_2} (p_2 + 1) + 4 \frac{p_2-2}{3} + 3 \\ &= \frac{p_3-2}{9p_2} \cdot (p_2 + 1)(7p_2 - 2) + \frac{4p_2+1}{3} \\ &= N(p_3 - 2) + \frac{4p_2+1}{3} \end{aligned}$$

as desired. Now for  $r_3 = 3p_2 - 2$ , from Theorem 3.5 (Theorem 6.1 in [1]),

$$\text{hw}(\Phi_{3p_2p_3}) = Np_3 - \left(\frac{4(p_2 + 1)}{3} - 2N\right) = N(p_3 + 2) - \frac{4(p_2 + 1)}{3},$$

as desired.  $\square$

#### ACKNOWLEDGEMENTS.

We would like to thank Hoon Hong and Eunjeong Lee for the discussions lead to the proof of Lemma 3.2. Also, we would like to We thank the editor and the anonymous referees for helpful feedbacks and detailed suggestions for improving the paper.

#### APPENDIX A. TECHNICAL PROOFS

- **Proof of Lemma 3.6:** We have

$$\begin{aligned} \Phi_{3p_2}(x) &= \sum_{i \geq 0} f_{3,p_2,i}(x) x^{ip_2} & \text{where } \deg f_{m,p_2,i}(x) < p_2 \\ f_{3,p_2,i}(x) &= \sum_{j \geq 0} f_{3,p_2,i,j}(x) x^{3j} & \text{where } \deg f_{3,p_2,i,j}(x) < 3 \end{aligned}$$

It is clear that  $i = 0, 1$ . Using Lemma 3.2-2 and knowing that  $a_0 = a_1 = a_2 = 2$  and  $b_1 = 1, b_0 = -1$ , we can compute the coefficient  $c_k$  in  $f_{3,p_2,i,j}$ :

$i$	$c_0$	$c_1$	$c_2$	$f_{3,p_2,i,0}$
0	1	-1	0	$1 - x$
1	0	-1	1	$x^2 - x$

Thus,

$$\Phi_{3p_2}(x) = \sum_{j=0}^{q_2-1} (1-x)x^{3j} + x^{3q_2}(1-x^{r_2-1}) + x^{p_2} \sum_{j=0}^{q_2-1} (x-x^2)x^{3j} + x^{\varphi(3p_2)}$$

which proves the Lemma.

REMARK A.1. The last proof is a concrete example which illustrate the partition on cyclotomic polynomials which we use in the proofs of this paper.

- **Proof of Lemma 4.10:** It is enough to prove that only one of the terms  $x^0$  and  $x^1$  will appear in  $f_{3p_2,p,i,0}$  when  $0 \leq u \leq \frac{q_2-3}{2}$  and only one of the terms  $x^0$  and  $x$  will appear in  $f_{3p_2,p,i,0}$  when  $\frac{q_2-1}{2} \leq u \leq q_2$  if  $v = 1$  while both of them will appear when  $v = 2$ . In the table below we list the term appears in  $C_i = \Psi_{3p_2} \cdot (x^2 - 1) \cdot \sum_{j=0}^u x^{6j}$  and the corresponding term in  $f_{3p_2,p,i,0}$ .

$u/\text{Terms in}$	$C_i$ if $v = 1$	$f_{3p_2,p,i,0}$ if $v = 1$	$C_i$ if $v = 2$	$f_{3p_2,p,i,0}$ if $v = 2$
$0 \leq u \leq \frac{q_2-1}{2}$	$-x^{6u+2v+1}$	$-x$	$x^{6u+2v}$	1
$\frac{q_2+1}{2} \leq u \leq q_2$	$x^{6u+2v}$	1	$-x^{6u+2v} - x^{6u+2v+1}$	$1 + x$

The reasoning why the term  $x^{6u+1}$  will not appear in  $C_i$  for  $\frac{q_2}{2} \leq u \leq q_2$ , was given in the proof of Lemma 4.3. Now for the case  $u > \frac{q_2}{2} - 1$  since  $i = 3u + v$  we have  $2i > 6(\frac{q_2}{2} - 1) + 2v = 3q_2 - 6 + 2v = p_2 - 7 + v$ . For  $x^0$  the equation  $x^{-2i} \cdot x^w = x^0 \Rightarrow w = 2i = 6u + 2v$  which has two solutions when  $v = 1$  and one solution when  $v = 0$ .

#### APPENDIX B. EXAMPLES

Here we add two examples to illustrate the results in the Sections 4.1 and 4.2.

EXAMPLE B.1. We will illustrate the results in Subsection 4.1 by an example. Let  $p_1 = 3, p_2 = 7$  and  $p_3 = 23$ . Note that  $r_2 = 1$  and  $r_3 = 2$ . The following two figures present the relationship between  $i$ ,  $\text{hw}(f_{21,p_3,i,0})$  and  $\text{hw}(f_{21,p_3,i,q})$ .






FIGURE 2. The relationship between  $i$ ,  $hw(f_{21,p_3,i,0})$  and  $hw(f_{21,p_3,i,q})$

EXAMPLE B.2. We will illustrate the results in Subsection 4.2 by an example. Let  $p_1 = 3, p_2 = 11$  and  $p_3 = 101$ . Note that  $r_2 = 2$  and  $r_3 = 2$ . The following figure presents the relationship between  $i$  and  $hw(f_{33,p_3,i,0})$  and  $hw(f_{33,p_3,i,q})$ .




FIGURE 3. The relationship between  $i$  and  $hw(f_{33,p_3,i,0})$




FIGURE 4. The relationship between  $i$  and  $\text{hw}(f_{33,p_3,i,q})$

Naslov

*Prvi autor, drugi autor i treći autor*

SAŽETAK. Hrvatski prijevod sažetka.

#### REFERENCES

- [1] A. Al-Kateeb. *Structures and properties of cyclotomic polynomials*. PhD thesis, North Carolina state university, 2016.
- [2] A. Al-Kateeb. On the height of cyclotomic polynomials. *Bull. Math. Soc. Sci. Math. Roumanie*, 110(1):101–106, 2019.
- [3] A. Al-Kateeb, M. Ambrosino, H. Hong, and E. Lee. Maximum gap in cyclotomic polynomials. *Journal of Number Theory*, 229:1–15, 2021.
- [4] A. Al-Kateeb, H. Hong, and E. Lee. Block structure of cyclotomic polynomials. *arXiv:1704.04051v2*, 2017.
- [5] G. Bachman. On the coefficients of ternary cyclotomic polynomials. *J. Number Theory*, 100:104–116, 2003.
- [6] M. Beiter. Magnitude of the coefficients of the cyclotomic polynomial  $F_{pqr}$ . *Amer. Math. Monthly*, 75(4):370–372, 1968.
- [7] B. Bzdega. Bounds on ternary cyclotomic coefficients. *Acta Arithmetica*, 144(1):5–16, 2010.
- [8] B. Bzdega. Jumps of ternary cyclotomic coefficients. *Acta Arithmetica*, 163(3):203–213, 2014.
- [9] L. Carlitz. The number of terms in the cyclotomic polynomial  $F_{pq}(x)$ . *Amer. Math. Monthly*, 73(9):979–981, 1966.
- [10] S. Elder. Flat cyclotomic polynomials: A new approach. *arXiv:1207.5811v1 [math.NT]*, 2012.
- [11] Y. Gallot and P. Moree. Neighboring ternary cyclotomic polynomials coefficients differ by at most one. *J. Ramanujan Math Soc*, 24:235–248, 2009.
- [12] Y. Gallot, P. Moree, and H. Hommersom. Value distribution of coefficients of cyclotomic polynomials. *Unif. Distrib. Theory*, 6:177–206, 2011.
- [13] Y. Gallot, P. Moree, and R. Wilms. The family of ternary cyclotomic polynomials with one free prime. *Involve*, 4(4):317–341, 2011.
- [14] H. Hong, E. Lee, H-S. Lee, and C-M. Park. Maximum gap in (inverse) cyclotomic polynomial. *J. Number Theory*, 132(10):2297–2317, 2012.
- [15] N. Kaplan. Flat cyclotomic polynomials of order three. *J. Number Theory*, 127(1):118–126, 2007.
- [16] N. Kaplan. Flat cyclotomic polynomials of order four and higher. *Integers*, 10:357–363, 2010.
- [17] P. Moree. Inverse cyclotomic polynomials. *J. Number Theory*, 129(3):667–680, 2009.
- [18] P. Moree. Numerical semigroups, cyclotomic polynomials, and bernoulli numbers. *Amer. Math. Monthly*, 121(10):890–902, 2014.
- [19] P. Moree and E. Rosu. Non-beiter ternary cyclotomic polynomials with an optimally large set of coefficients. *I. J. Number Theory*, 8(8):1883–1902, 2012.
- [20] C. Sanna. A survey on coefficients of cyclotomic polynomials. *Expo. Math*, 2022.
- [21] B. Zhang. Remarks on the maximum gap in binary cyclotomic polynomials. *Bull. Math. Soc. Sci. Math. Roumanie (N.S.)*, 59:109–115, 2016.
- [22] B. Zhang. Remarks on the flatness of ternary cyclotomic. *I. J. Number theory*, 13(2):529–547, 2017.

First author  
 Department of Mathematics  
 Yarmouk University- Jordan  
 E-mail: alaa.kateeb@yu.edu.jo

Second author  
Department of Mathematics  
Yarmouk University- Jordan  
*E-mail:* `afnand@yu.edu.jo`