High rank elliptic curves with prescribed torsion over quadratic fields

Maintained by Andrej Dujella, University of Zagreb

Let T be one of 26 admissible torsion groups for an elliptic curve over a quadratic field [KM,Ka]. We define B(T) as the supremum of the ranks of elliptic curves defined over any quadratic field and having torsion group T (for the trivial torsion group we put T = 0).

In the following table, in the second column we give the best known lower bounds for B(T), while the third column gives the values d of the corresponding quadratic number fields $\mathbf{K} = \mathbf{Q}(\sqrt{d})$.

Values in the brackets in the second column are conditional, assuming the Parity Conjecture.

T	B(T)>=	d	References
0	30	-3	[Na,E14], [E11,E13,Me]
z /2 z	28	-1	[Wa, ACP, ADJBP]
z /3 z	22	-3	[Na, ER]
Z /4 Z	15 (16)	-25689	[ADJBP], [Na]
z /5 z	10	*	[Le1,Er,Me]
z /6 z	11	3521	[ADJBP]
z /7 z	7	*	[Ku,El2,Er,Le2,Me]
z /8 z	9	-227	[ADJBP]
z /9 z	6	-155	[ADJBP], [Fi,Me]
z /10 z	7	-2495	[ADJBP]
z /11 z	2	-3239	[ADJBP]
z /12 z	7	2014	[ADJBP]
z /13 z	2	193	[Ra]
z /14 z	2	265	[Ra, ADJBP]
z /15 z	1	-7	[BBDN], [ADJBP]
z /16 z	3 (4)	34720105	[Na]
z /18 z	2	26521	[BBDN]
$\mathbf{z}/2\mathbf{z} \times \mathbf{z}/2\mathbf{z}$	19 (20)	d_{22}	[ADJBP,E12]
$\mathbf{z}/2\mathbf{z} \times \mathbf{z}/4\mathbf{z}$	13	-83201	[ADJBP]
$\mathbf{z}/2\mathbf{z} \times \mathbf{z}/6\mathbf{z}$	10	624341	[ADJBP]
$\mathbf{z}/2\mathbf{z} \times \mathbf{z}/8\mathbf{z}$	8	31230597	[ADJBP]
$\mathbf{z}/2\mathbf{z} \times \mathbf{z}/10\mathbf{z}$	4 (5)	1065333545	[BBDN], [ADJBP]
$\mathbf{z}/2\mathbf{z} \times \mathbf{z}/12\mathbf{z}$	4	2947271015	[BBDN]
$\mathbf{z}/3\mathbf{z} \times \mathbf{z}/3\mathbf{z}$	7	-3	[JB]
$\mathbf{z}/3\mathbf{z} \times \mathbf{z}/6\mathbf{z}$	6	-3	[JB]
$\mathbf{Z}/4\mathbf{Z} \times \mathbf{Z}/4\mathbf{Z}$	7	-1	[DJB]

$d_{22} = -3901785498412536920668361993073821511$

The marks * in the table refer to the curves obtained by an application of Mestre's general construction (he proved that for any elliptic curve E over \mathbf{Q} there exist infinitely many quadratic twists with rank ≥ 2) which produces quadratic fields with huge discriminants.

References:

- [ACP] J. Aguirre, F. Castaneda, J. C. Peral, <u>High rank elliptic curves with torsion group Z/(2Z)</u>, Math. Comp. **73** (2004), 323-331.
- [ADJBP] J. Aguirre, A. Dujella, M. Jukic Bokun, J. C. Peral, <u>High rank elliptic curves with prescribed torsion group over quadratic fields</u>, Period. Math. Hungar. **68** (2014), 222-230.
- [BBDN] J. Bosman, P. Bruin, A. Dujella and F. Najman, *Ranks of elliptic curves with prescribed torsion over number fields*, Int. Math. Res. Not. IMRN **2014 (11)** (2014), 2885-2923.
- [DJB] A. Dujella and M. Jukic Bokun, On the rank of elliptic curves over Q(i) with torsion group Z/4Z × Z/4Z, Proc. Japan Acad. Ser. A Math. Sci. 86 (2010), 93-96.
- [DJBS] A. Dujella, M. Jukic Bokun and I. Soldo, <u>On the torsion group of elliptic curves induced by Diophantine triples over quadratic fields</u>, Rev. R. Acad. Cienc. Exactas Fis. Nat. Ser. A Math. RACSAM, to appear.
- [El1] N. D. Elkies, \mathbf{Z}^{28} in $E(\mathbf{Q})$, etc., Number Theory Listserver, May 2006.
- [E12] N. D. Elkies, Personal communication, 2006, 2009.
- [El3] N. D. Elkies, *Three lectures on elliptic surfaces and curves of high rank*, Lecture notes, Oberwolfach, 2007, arXiv:0709.2908.
- [El4] N. Elkies, *j* = 0; rank 15; also 3-rank 6 and 7 in real and imaginary quadratic fields, Number Theory Listserver, Dec 2009.
- [ER] N. D. Elkies and N. F. Rogers, <u>Elliptic curves $x^3 + y^3 = k$ of high rank</u>, Proceedings of ANTS-6 (D. Buell, ed.), Lecture Notes in Comput. Sci. **3076** (2004), 184-193.
- [Er] Y. G. Eroshkin, Personal communication, 2009, 2011.
- [Fi] T. A. Fisher, Personal communication, 2009.
- [JB] M. Jukic Bokun, On the rank of elliptic curves over $Q(\sqrt{-3})$ with torsion groups $\mathbb{Z}/3\mathbb{Z} \times \mathbb{Z}/3\mathbb{Z}$ and $\mathbb{Z}/3\mathbb{Z} \times \mathbb{Z}/6\mathbb{Z}$, Proc. Japan Acad. Ser. A Math. Sci. 87 (2011), 61-64.
- [JB2] M. Jukic Bokun, *Elliptic curves over quadratic elds with fixed torsion subgroup and positive rank*, Glas. Mat. Ser. III 47 (2012), 277-284.
- [Ka] S. Kamienny, *Torsion points on elliptic curves and q-coecients of modular forms*, Invent. Math. 109 (1992), 221-229.
- [KM] M. A. Kenku and F. Momose, *Torsion points on elliptic curves defined over quadratic fields*, Nagoya Math. J. **109** (1988), 125-149.
- [Ku] L. Kulesz, Arithmetique des courbes algebriques de genre au moins deux, These de doctorat, Universite Paris 7, 1998.
- [Le1] O. Lecacheux, Rang de courbes elliptiques sur *Q* avec un groupe de torsion isomorphe a *Z/5Z*, C. R. Acad. Sci. Paris Ser. I Math. 332 (2001), 1-6.
- [Le2] O. Lecacheux, <u>Rang de courbes elliptiques dont le groupe de torsion est non trivial</u>, Ann. Sci. Math. Quebec 28 (2004), 145-151.
- [Me] J.-F. Mestre, Rang des courbes elliptiques d'invariant donné, C. R. Acad. Sci. Paris 314 (1992), 919-922.
- [Na] F. Najman, *Some rank records for elliptic curves with prescribed torsion over quadratic fields*, An. Stiint. Univ. "Ovidius" Constanta Ser. Mat. **22** (2014), 215-220.
- [Ra] F. P. Rabarison, Structure de torsion des courbes elliptiques sur les corps quadratiques, Acta Arith. 144 (2010), 17-52.
- [SZ] U. Schneiders and H.G. Zimmer, *The rank of elliptic curves upon quadratic extensions*, in: Computational Number Theory (A. Petho, H.C. Williams, H.G. Zimmer, eds.), de Gruyter, Berlin, 1991, pp. 239-260.
- [Wa] M. Watkins, Personal communication, 2005.

High rank elliptic curves with prescribed torsion

Infinite families of elliptic curves with high rank and prescribed torsion

History of elliptic curves rank records