

Diophantine m -tuples

Andrej Dujella

Department of Mathematics
University of Zagreb, Croatia

e-mail: duje@math.hr

URL: <http://web.math.hr/~duje/>

Diophantus: Find four numbers such that the product of any two of them, increased by 1, is a perfect square:

$$\left\{ \frac{1}{16}, \frac{33}{16}, \frac{17}{4}, \frac{105}{16} \right\}$$

Fermat: $\{1, 3, 8, 120\}$

$$\begin{aligned} 1 \cdot 3 + 1 &= 2^2, & 3 \cdot 8 + 1 &= 5^2, \\ 1 \cdot 8 + 1 &= 3^2, & 3 \cdot 120 + 1 &= 19^2, \\ 1 \cdot 120 + 1 &= 11^2, & 8 \cdot 120 + 1 &= 31^2. \end{aligned}$$

Euler: $\{1, 3, 8, 120, \frac{777480}{8288641}\}$

$$ab + 1 = r^2 \mapsto \{a, b, a + b + 2r, 4r(a + r)(b + r)\}$$

Gibbs (1999): $\{\frac{11}{192}, \frac{35}{192}, \frac{155}{27}, \frac{512}{27}, \frac{1235}{48}, \frac{180873}{16}\}$

Dujella (2009):

$$\{\frac{27}{35}, -\frac{35}{36}, -\frac{352}{315}, \frac{1007}{1260}, -\frac{5600}{4489}, \frac{72765}{106276}\}$$

Definition: A set $\{a_1, a_2, \dots, a_m\}$ of m non-zero integers (rationals) is called a (*rational*) *Diophantine m -tuple* if $a_i \cdot a_j + 1$ is a perfect square for all $1 \leq i < j \leq n$.

Question: How large such sets can be?

Conjecture 1: There does not exist a Diophantine quintuple.

Baker & Davenport (1969):

$$\{1, 3, 8, d\} \Rightarrow d = 120$$

(problem raised by Gardner (1967), van Lint (1968))

Arkin, Hoggatt & Strauss (1978): Let

$$ab + 1 = r^2, \quad ac + 1 = s^2, \quad bc + 1 = t^2$$

and define

$$d_{+,-} = a + b + c + 2abc \pm 2rst.$$

Then $\{a, b, c, d_{+,-}\}$ is a Diophantine quadruple (if $d_- \neq 0$).

Conjecture 2: If $\{a, b, c, d\}$ is a Diophantine quadruple, then $d = d_+$ or $d = d_-$, i.e. all Diophantine quadruples satisfy

$$(a - b - c + d)^2 = 4(ad + 1)(bc + 1).$$

Such quadruples are called *regular*.

D. & Fuchs (2004): All Diophantine quadruples in $\mathbb{Z}[X]$ are regular.

D. & Jurasić (2010): In $\mathbb{Q}(\sqrt{-3})[X]$, the Diophantine quadruple

$\left\{ \frac{\sqrt{-3}}{2}, -\frac{2\sqrt{-3}}{3}(X^2 - 1), \frac{-3 + \sqrt{-3}}{3}X^2 + \frac{2\sqrt{-3}}{3}, \frac{3 + \sqrt{-3}}{3}X^2 + \frac{2\sqrt{-3}}{3} \right\}$
is not regular.

D. (1997): $\{k-1, k+1, 4k, d\} \Rightarrow d = 16k^3 - 4k$

D. & Pethő (1998): $\{1, 3\}$ cannot be extended to a Diophantine quintuple

Fujita (2008): $\{k-1, k+1\}$ cannot be extended to a Diophantine quintuple

Bugeaud, D. & Mignotte (2007):

$\{k-1, k+1, 16k^3 - 4k, d\} \Rightarrow$
 $d = 4k$ or $d = 64k^5 - 48k^3 + 8k$

D. (2004): There does not exist a Diophantine sextuple.

There are only finitely many Diophantine quintuples.

$$\max\{a, b, c, d, e\} < 10^{10^{26}}$$

Fujita (2009): If $\{a, b, c, d, e\}$, with $a < b < c < d < e$, is a Diophantine quintuple, then $\{a, b, c, d\}$ is a regular Diophantine quadruple.

Extending the Diophantine triple $\{a, b, c\}$, $a < b < c$, to a Diophantine quadruple $\{a, b, c, d\}$:

$$ad + 1 = x^2, \quad bd + 1 = y^2, \quad cd + 1 = z^2.$$

System of simultaneous Pellian equations:

$$cx^2 - az^2 = c - a, \quad cy^2 - bz^2 = c - b.$$

Binary recursive sequences:

finitely many equations of the form $v_m = w_n$.

Linear forms in three logarithms:

$$v_m \approx \alpha\beta^m, \quad w_n \approx \gamma\delta^n \Rightarrow$$

$$m \log \beta - n \log \delta + \log \frac{\alpha}{\gamma} \approx 0$$

Baker's theory gives upper bounds for m, n (logarithmic functions in c).

Simultaneous Diophantine approximations:

$\frac{x}{z}$ and $\frac{y}{z}$ are good rational approximations to $\sqrt{\frac{a}{c}}$ and $\sqrt{\frac{b}{c}}$, resp.

$\frac{bsx}{abz}$ and $\frac{aty}{abz}$ are good rational approximations to $\frac{s}{a}\sqrt{\frac{a}{c}} = \sqrt{1 + \frac{b}{abc}}$ and $\frac{t}{b}\sqrt{\frac{b}{c}} = \sqrt{1 + \frac{a}{abc}}$, resp.

If c is large compared to b (say $c > b^6$), then hypergeometric method gives (very good) upper bounds for x, y, z .

Congruence method (D. & Pethő):

$$v_m \equiv w_n \pmod{c^2}$$

If m, n are small (compared with c), then \equiv can be replaced by $=$, and this (hopefully) leads to a contradiction (if $m, n > 2$).

Therefore, we obtain lower bounds for m, n (small powers of c , e.g. $c^{0.04}$).

Conclusion: Contradiction for large c .

If $\{k - 1, k + 1, c\}$ is a Diophantine triple, then $c = c_\nu$, where

$$c_1 = 4k, \quad c_2 = 16k^3 - 4k, \quad c_3 = 64k^5 - 48k^3 + 8k, \dots$$

For c_ν , $\nu \geq 3$, gap is large enough for the application of results on simultaneous Diophantine approximations – **Fujita (2008)**.

The case c_1 leads to simultaneous approximations to the numbers $\sqrt{1 - \frac{1}{k}}$ and $\sqrt{1 + \frac{1}{k}}$ (a result by **Rickert (1993)**) – **D. (1997)**.

For c_2 – **Bugeaud, D. & Mignotte (2007)**:

Improved congruence method:

Combination of congruences mod $4k(k-1)$ and mod c_2^2 gives $m > 4.9k^{1.5}$ (if $m > 2$).

Recent results on linear forms in three logarithms:

by **Matveev (2000)**: $k < 3.8 \cdot 10^{10}$;

by **Mignotte (2007)**: $k < 5.4 \cdot 10^8$.

Baker-Davenport reduction method:

Starting with $m \leq 3.6 \cdot 10^{16}$, we obtain $m \leq 2$.

Bo He, Togbé, Filipin (2009,2010):

$$\{k, A^2k + 2A, (A+1)^2k + 2(A+1)\}$$

extends uniquely to a Diophantine quadruple if

$$1 \leq A \leq 22 \text{ or } A \geq 51767$$

(using linear forms in *two* logarithms)

Let $\{a, b, c\}$ be a Diophantine triple. Consider the elliptic curve

$$E : y^2 = (ax + 1)(bx + 1)(cx + 1).$$

Rational points $P = [0, 1]$, $Q = [1/abc, rst/abc]$ satisfy $x(P \mp Q) = d_{+,-}$.

Conjecture 3: All integer points on E are: $[0, \pm 1]$, $[d_+, \pm(at + rs)(bs + rt)(cr + st)]$, $[d_-, \pm(at - rs)(bs - rt)(cr - st)]$, and also $[-1, 0]$ if $1 \in \{a, b, c\}$.

D. (2000): Conjecture is true for elliptic curves

$$E_k : y^2 = ((k-1)x+1)((k+1)x+1)(4kx+1),$$

under assumption that $\text{rank } E_k(\mathbb{Q}) = 1$ (also for two subfamilies of rank 2 and one subfamily of rank 3). Furthermore, it is true for all k , $2 \leq k \leq 1000$ (extended to $k \leq 5000$ by **Najman (2010)**). The condition $\text{rank } E_k(\mathbb{Q}) = 1$ is not unrealistic since $\text{rank } E(\mathbb{Q}(k)) = 1$.

D. & Pethő (2000): Conjecture is true for elliptic curves

$$E'_k : y^2 = (x + 1)(3x + 1)(c_k x + 1),$$

where $\{1, 3, c_k\}$ is a Diophantine triple, i.e.

$$c_k = \frac{1}{6} \left((2 + \sqrt{3})(7 + 4\sqrt{3})^k + (2 - \sqrt{3})(7 - 4\sqrt{3})^k - 4 \right),$$

under assumption that $\text{rank } E'_k(\mathbb{Q}) = 2$. Furthermore, it is true for all k , $1 \leq k \leq 40$, with possible exceptions $k = 23$ and $k = 37$ (extended by **Jacobson & Williams (2002)** to $k \leq 100$, with the possible exception of $k = 37$, for which the result holds under the Extended Riemann Hypothesis).

Similar results for other families of Diophantine triples:

D. (2001), Fujita (2007, 2008), Najman (2009, 2010).

Definition: Let n be an integer. A set of m positive integers is called a *Diophantine m -tuple with the property $D(n)$* or simply *$D(n)$ - m -tuple* (or P_n -set of size m), if the product of any two of them, increased by n , is a perfect square.

$$M_n = \sup\{\#D : D \text{ is a } D(n)\text{-tuple}\}$$

Conjecture 4: There exist a constant C such that $M_n < C$ for all non-zero integers n .
In particular, there does not exist a rational C -tuple.

D. (2004): $4 \leq M_1 \leq 5$
(implies directly $4 \leq M_4 \leq 7$)

Filipin (2008): $4 \leq M_4 \leq 5$

D. (2004): $M_n \leq 31$ if $|n| \leq 400$
 $M_n < 15.476 \cdot \log |n|$ if $|n| > 400$

D. & Luca (2005): $M_p < 2^{146}$ if p is a prime

Brown, Gupta & Singh, Mohanty & Ramasamy (1985):

If $n \equiv 2 \pmod{4}$, then $M_n = 3$.

D. (1993): If $n \not\equiv 2 \pmod{4}$ and $n \notin S_1 = \{-4, -3, -1, 3, 5, 8, 12, 20\}$, then $M_n \geq 4$.

Conjecture 5: If $n \in S_1$, then $M_n = 3$.

D. & Fuchs (2005): $3 \leq M_{-1} \leq 4$

Remark: $n \equiv 2 \pmod{4}$ if and only if n is not representable as a difference of the squares of two integers.

D. (1997), Franušić (2004, 2008): Analogous results: strong connection between the existence of $D(n)$ -quadruples and the representability as a difference of two squares also hold for integers in some quadratic fields.

D., Filipin & Fuchs (2007): There are only finitely many $D(-1)$ -quadruples.

If $\{a, b, c, d\}$ is a $D(-1)$ -quadruple, then $\max\{a, b, c, d\} < 10^{10^{23}}$.

Conjecture 6: If n is not a perfect square, then there exist only finitely many $D(n)$ -quadruples.

Euler: There exist infinitely many $D(1)$ -quadruples, and therefore infinitely many $D(k^2)$ -quadruples.

DFF implies that the conjecture is true for $n = -1$ and $n = -4$ (note that all elements of a $D(-4)$ -quadruple are even).

$$\boxed{a_i \cdot a_j + 1 = k\text{-th power}} \quad k \geq 3 \text{ fixed}$$

Such a set is called a *k-th power Diophantine m-tuple*.

$\{2, 171, 25326\}$ is a third power Diophantine triple

$\{1352, 8539880, 9768370\}$ is a fourth power Diophantine triple

$$C(k) = \sup\{\#D : D \text{ is a } k\text{-th power D. tuple}\}$$

Bugeaud & D. (2003): $C(3) \leq 7$, $C(4) \leq 5$,
 $C(k) \leq 4$ for $5 \leq k \leq 176$, $C(k) \leq 3$ for $k \geq 177$

$$a_i \cdot a_j + 1 = \text{perfect power}$$

Such a set is called a *Diophantine powerset*.

$D \subset \{1, 2, \dots, N\}$ such that $ab + 1$ is a perfect power for all $a \neq b$ in D .

Gyarmati, Sárközy & Stewart (2002):

$$\#D \leq 340 \frac{(\log N)^2}{\log \log N}$$

Improvements by **Bugeaud-Gyarmati (2004)**,
Dietmann-Elsholtz-Gyarmati-Simonovits (2005),
Luca (2005), **Gyarmati-Stewart (2007)**

Stewart (2008): $\#D \ll (\log N)^{2/3} (\log \log N)^{1/3}$

Luca (2005): *abc-conjecture* implies that $\#D$ is bounded by an absolute constant.

D., Fuchs & Luca (2008):

In $\mathbb{Z}[X]$, $\#D < 8 \cdot 10^5$.

D. & Jurasić (2010):

In $\mathbb{K}[X]$, where \mathbb{K} is a field of characteristic 0,
 $\#D < 2 \cdot 10^7$.

Let $D_m(N) =$
 $\# \{D \subseteq \{1, 2, \dots, N\} : D \text{ is a Diophantine-}m\text{-tuple} \}.$

D. (2008): $D_2(N) = \frac{6}{\pi^2} N \log N + O(N);$

$ab + 1 = r^2 \rightarrow r^2 \equiv 1 \pmod{b}$

$D_3(N) = \frac{3}{\pi^2} N \log N + O(N);$

almost all triples are of form $\{a, b, a + b + 2r\}$

$0.1608 \sqrt[3]{N} \log N < D_4(N) < 0.5354 \sqrt[3]{N} \log N$

Martin & Sitar (2010):

$D_4(N) = C \sqrt[3]{N} \log N + O(\sqrt[3]{N} (\log N)^{2/3 + \sqrt{2}/6} (\log \log N)^{5/12}),$
 where $C = \frac{2^{4/3}}{3\Gamma(\frac{2}{3})^3} \approx 0.338285.$

almost all quadruples are on the form

$\{a, b, a + b + 2r, 4r(a + r)(b + r)\};$

Erdős-Turán inequality - discrepancy between the number of elements of a sequence that lie in a particular interval modulo 1 and the expected number;

equidistribution of solutions of polynomial congruences

Fujita (2010): $D_5(N) < 10^{276}$

a fixed Diophantine triple $\{a, b, c\}$ has at most 4 extensions to Diophantine quintuple $\{a, b, c, d, e\}$ such that $\max\{a, b, c\} < d < e$