

Mathematical Thinking

Module 1: Introduction to Mathematical Thinking

Instructor: Keith Devlin

Generated by mooc2handout

July 29, 2026

This handout was generated from lecture captions, supplementary materials, and AI-inferred keyframes.

Contents

Knowledge Map

Topic	Core Question
Mathematical Thinking	What is mathematics really about? Why is it more than computation?
Precise Language	Why does math require more precision than everyday language?
Logical Connectives	How do AND, OR, NOT, IF...THEN combine propositions?
Quantifiers	How do we express 'for all' and 'there exists'? What are the negation rules?
Set Theory Basics	How are sets, empty set, subsets, unions, and intersections defined?

0. Overview

This unit introduces the mindset of mathematical thinking. Unlike school math, which focuses on procedures and computation, this course emphasizes **understanding the nature of mathematics**—a precise, rigorous, and abstract way of thinking developed over 3,000 years.

Core Objectives

- Transition from "school math" to "university math" thinking
- Understand that mathematics is the study of **patterns**
- Master the importance of precise language in mathematical reasoning
- Learn to use logical connectives and quantifiers for rigorous argumentation

1. Welcome

Review: [Welcome](#)(8:15)

Keith Devlin opens with an analogy: if mathematics were the automotive world, **school math is learning to drive**, while this course teaches you **how the engine works**.

- Mathematical thinking is a mental ability developed over 3,000 years
- This course teaches no new formulas or computational techniques
- The goal is to help you transition from high school to university-level math
- Key distinction: school math asks **how**; university math asks **why**

Practical advice: Don't expect to solve every assignment problem in one sitting. What matters is **attempting each question**, not meeting a deadline.

Supplementary material: -0- - [_Supplement__Set_Theory_.pdf](#)


Auto-generated reading digest Expanded automatically from the extracted PDF text to surface the main points.

- In general, however, except for finite sets with only a small number of elements, sets are best described by giving the property which defines the set.
- If we have to prove that the sets A and B are equal, we usually split the proof into two parts: (a) Show that every member of A is a member of B .
- This set is denoted by A , and has the formal definition $A = \{x \in U \mid x \in A\}$ [Notice that we write $x \in A$ instead of $\neg(x \notin A)$, for brevity.] For instance, if the universal set is the set N of natural numbers, and E is the set of even (natural) numbers, then E is the set of


Supplementary material: -0- - [_Background_Reading.pdf](#)

Auto-generated reading digest Expanded automatically from the extracted PDF text to surface the main points.

- For example, the commutative law for addition could be written in English as: When two numbers are added, their order is not important.
- Counter-intuitive (and occasionally paradoxical) results made them realize that some of the methods they had developed to solve important, real-world problems had consequences they could not explain.
- This introspection led, in the middle of the nineteenth century, to the adoption of a new and different conception of the mathematics, where the primary focus was no longer on performing a calculation or computing an answer, but formulating and understanding abstract concepts and relationships.



Timestamp 29s - Concept shift



Timestamp 44s - Concept shift

2. Introduction to the Material

Review: [Introduction to the Material](#) (27:59)

This lecture answers a fundamental question: **What is mathematics, really?**

Devlin points out that schools spend years teaching procedures but rarely explain what the subject is about. It's like explaining soccer by saying "it's a series of maneuvers to get the ball into the goal"—not wrong, but missing the big picture.

The Nature of Mathematics Mathematics is the study of **patterns**:

- Counting patterns (arithmetic)
- Measurement patterns (geometry)
- Symmetry patterns (group theory)
- Motion patterns (calculus)
- Change patterns (analysis)
- Randomness patterns (probability)

Mathematical notation is not pretension—it is a **necessary tool** for describing abstract patterns precisely, just as maps are for terrain and musical scores are for sound.

Historical milestone: Around the 6th century BCE, Thales and Pythagoras first proposed that precisely stated assertions could be proved by formal arguments—marking the birth of the **theorem**.



Introduction to Mathematical Thinking




One American dies
of melanoma almost
every hour.

Almost every hour,
an American dies of
melanoma.



Timestamp 671s - Core definition



Timestamp 824s - Core definition

3. Logical Connectives

Review: [Logical Connectives](#) (26:08)


This lecture introduces the core tools of mathematical reasoning: **logical connectives**.

Everyday language uses "and," "or," and "if...then" loosely. Mathematics requires **precise, unambiguous** definitions:


Five Basic Logical Connectives

1. **Conjunction (AND):** $P \wedge Q$ — true iff both P and Q are true
2. **Disjunction (OR):** $P \vee Q$ — true if at least one is true (inclusive OR!)
3. **Negation (NOT):** $\neg P$ — flips the truth value
4. **Conditional (IF...THEN):** $P \Rightarrow Q$ — false only when P is true and Q is false
5. **Biconditional (IFF):** $P \Leftrightarrow Q$ — true when P and Q have the same truth value

Common pitfall: In everyday English, "or" often means "one or the other but not both" (exclusive OR). In mathematics, $\vee$ includes the case where both are true.



Timestamp 106s - Core definition



Timestamp 159s - Core definition

4. Assignment 1 Tutorial


Review: Assignment 1 Tutorial(2:30)

This tutorial covers Assignment 1, focusing on **Euclid's proof that there are infinitely many primes**.


Core argument:

1. Assume primes are finite: $p_1, p_2, \dots, p_n$
2. Construct $N = p_1 \times p_2 \times \dots \times p_n + 1$
3. N leaves remainder 1 when divided by any p_i , so no known prime divides it
4. Therefore N is either a new prime itself, or has a prime factor not in the list
5. Either way, this contradicts "primes are finite"

Assignment tip: Question 8 asks "how to prove $p_1 p_2 \dots p_n + 1$ is not always prime"—find a counterexample (e.g., $2 \times 3 \times 5 \times 7 \times 11 \times 13 + 1 = 30031 = 59 \times 509$).



Timestamp 31s - Concept shift



Timestamp 69s - Concept shift

5. Assignment 2 Tutorial


Review: Assignment 2 Tutorial(9:21)

This tutorial covers Assignment 2: **simplifying logical expressions**.


Core technique: identify **superfluous conjuncts**:

- If $x < 4$, then $x < 6$ is automatically true — the second condition is redundant
- If $y^2 < 9$ (i.e., $-3 < y < 3$), then $y < 4$ is automatic — the first condition is redundant
- $7 \leq p < 12$ can be written concisely as a chained inequality

Method: Draw each condition as an interval on the number line, take the intersection, then check which condition is fully contained by another.



Timestamp 18s - Concept shift



Timestamp 27s - Concept shift

6. Key Takeaways

1. Mathematics is the study of patterns, not just computation.

2. Precise language is the foundation of mathematical reasoning—ambiguity is unacceptable.
3. Logical connectives (AND/OR/NOT/IF...THEN) have strict definitions; don't rely on everyday intuition.
4. Quantifier negation: $\forall$ becomes $\exists$, $\exists$ becomes $\forall$, and the inner proposition is negated.
5. The heart of proof is logical argument, not example verification.

7. Exercises

A. Multiple Choice (10 questions)

1. What does mathematics primarily study?
A. Number crunching B. Patterns C. Physical measurement D. Computer programming
2. $P \vee Q$ is false when:
A. P false, Q true B. P true, Q false C. Both P and Q are false D. Both P and Q are true
3. $P \Rightarrow Q$ is false only when:
A. P false, Q false B. P false, Q true C. P true, Q false D. P true, Q true
4. $\neg \forall x P(x)$ is equivalent to:
A. $\forall x \neg P(x)$ B. $\exists x \neg P(x)$ C. $\exists x P(x)$ D. $\neg \exists x P(x)$
5. Euclid's proof of infinitely many primes uses:
A. Direct proof B. Mathematical induction C. Proof by contradiction D. Constructive proof
6. Which correctly represents the empty set?
A. $\{0\}$ B. $\{\}$ C. $\{\emptyset\}$ D. 0
7. If $x < 4$, then $x < 6$ is:
A. An independent condition B. A superfluous condition C. A contradictory condition D. A necessary and sufficient condition
8. The main difference between mathematical OR and everyday 'or' is:
A. Math OR excludes both being true B. Math OR includes both being true C. No difference D. Math OR only applies to numbers
9. $A \subseteq B$ means:
A. A equals B B. Every element of A is in B C. Every element of B is in A D. A and B share no elements
10. The birth of the theorem marked:
A. Math started using numbers B. Precise assertions can be proved by formal arguments C. Algebra separated from geometry D. The invention of calculus

B. True/False (5 questions)

1. In mathematics, $P \vee Q$ is still true when both P and Q are true. ()

2. $P \Rightarrow Q$ is always false when P is false.()
3. $\neg \exists x P(x)$ is equivalent to $\forall x \neg P(x)$.()
4. The empty set $\emptyset$ and $\{\emptyset\}$ are the same set.()
5. A single example can prove a universal statement.()

C. Short Answer (2 questions)

1. Explain why mathematics requires more precise language than everyday speech. Give an example where everyday ambiguity leads to misunderstanding.
2. Outline Euclid's proof of infinitely many primes using proof by contradiction. Clearly state the assumption, construction, and contradiction.

8. Answer Key

Multiple Choice

1.B 2.C 3.C 4.B 5.C 6.B 7.B 8.B 9.B 10.B

True/False

1. True 2. False 3. True 4. False 5. False

Short Answer Key Points

1. Everyday 'or' often means exclusive (one or the other), but $\vee$ includes both; everyday 'if' implies causation, but $\Rightarrow$ only checks truth values. Precision eliminates ambiguity.
2. Assume primes are finite $p_1, \dots, p_n$; construct $N = p_1 p_2 \cdots p_n + 1$; no p_i divides N , so either N is prime or has a new prime factor—contradiction.