

# ON THE HEIGHT OF TREES

A. RÉNYI and G. SZEKERES<sup>1</sup>

(Received 12 October 1965)

## 1. Introduction

In this note we shall deal with the enumeration of labelled trees of given order and given height over a selected point.

An undirected graph is called a tree if it is connected and contains no cycle. If we select any two vertices  $P$  and  $Q$  of a tree  $T$ , there is evidently a uniquely determined path in  $T$  leading from  $P$  to  $Q$ . We shall call the length of this path (i.e. the number of edges in the path) the distance of  $P$  and  $Q$  in  $T$  and denote it by  $d_T(P, Q)$ . If a vertex  $P$  is distinguished as the root of  $T$ , we define the height of  $T$  over  $P$  as the length of the longest path in  $T$  starting from  $P$ ; thus if  $h_P(T)$  denotes the height of  $T$  over the root  $P$ , we have

$$(1.1) \quad h_P(T) = \max_{Q \in T} d_T(P, Q).$$

Let us consider the set  $\mathcal{T}_n$  of all possible trees with  $n$  given labelled vertices  $P_1, P_2, \dots, P_n$ . According to a classical result of Cayley [1] if  $t_n$  denotes the number of elements of  $\mathcal{T}_n$ , we have

$$(1.2) \quad t_n = n^{n-2}$$

Let  $t_n(k)$  denote the number of those trees  $T \in \mathcal{T}_n$  for which  $h_{P_1}(T) \leq k$ . Clearly

$$(1.3) \quad t_1(0) = 1, \quad t_n(0) = 0 \quad \text{for } n > 1$$

and

$$(1.4) \quad t_n(k) = t_n \quad \text{for } k \geq n-1.$$

J. Riordan [2] has shown that the enumerator

$$(1.5) \quad G_k(x) = \sum_{n=1}^{\infty} \frac{t_n(k)}{(n-1)!} x^n \quad (k = 0, 1, \dots)$$

<sup>1</sup> Part of this work was done at the fifth Summer Research Institute of the Australian Mathematical Society in Canberra. The first author is greatly indebted to the Australian National University for its generous support of his visit to the Institute.

satisfies the recursion formula

$$(1.6) \quad G_{k+1}(x) = x \exp G_k(x) \quad (k = 0, 1, \dots)$$

with

$$(1.7) \quad G_0(x) = x;$$

the latter follows from (1.3) and (1.5).

From the recursion formula (1.6) one can determine  $t_n(k)$  for any  $k$  and  $n$  ( $0 \leq k \leq n-1$ ). For instance

$$(1.8) \quad G_1(x) = xe^x, \quad G_2(x) = xe^{xe^x}, \quad \text{etc.},$$

and thus

$$(1.9) \quad t_n(1) = 1 \quad (n = 1, 2, \dots)$$

$$(1.10) \quad t_n(2) = \sum_{m=0}^{n-1} \binom{n-1}{m} m^{n-m-1} \quad (n = 1, 2, \dots)$$

and generally for  $k \geq 1$

$$(1.11) \quad t_n(k) = \sum_{\substack{m_1 + \dots + m_k = n-1 \\ m_i \geq 0}} \frac{(n-1)!}{m_1! m_2! \dots m_k!} m_1^{m_1} m_2^{m_2} \dots m_k^{m_k} \quad (i = 1, 2, \dots, k).$$

In these formulae  $0^0$  always means 1.

In view of (1.2) and (1.4) one has

$$(1.12) \quad \lim_{k \rightarrow \infty} G_k(x) = \sum_{n=1}^{\infty} \frac{n^{n-2}}{(n-1)!} x^n$$

provided that the series on the right of (1.12) is convergent. But the series

$$(1.13) \quad y = \sum_{n=1}^{\infty} \frac{n^{n-2}}{(n-1)!} x^n$$

converges for  $|x| \leq 1/e$  and represents the inverse function of

$$(1.14) \quad x = ye^{-y}.$$

This equation also follows from (1.6) and (1.12).

Riordan [2] obtained the formula (1.6) as a special case of a more general result on enumerators of trees. In § 2 we shall give direct proofs of (1.6) and (1.11).

In § 3 we shall investigate the asymptotic distribution of

$$(1.15) \quad d_n(k) = t_n(k) - t_n(k-1),$$

i.e. the number of trees  $T \in \mathcal{T}_n$  having exact height  $k$  over  $P_1$ . Let us

mention that if  $D(T)$  denotes the diameter of  $T$  (i.e. the length of the longest path in  $T$ ) one has evidently

$$(1.16) \quad \frac{1}{2}D(T) \leq \min_i h_{P_i}(T) \leq h_{P_i}(T) \leq \max_i h_{P_i}(T) \leq D(T).$$

Thus the study of the distribution of  $h_{P_i}(T)$  for  $T \in \mathcal{T}_n$  gives us also some information on the distribution of  $D(T)$ .

Our thanks are due to F. Harary and J. W. Moon for calling our attention to the paper [2] of Riordan.

## 2. Proof of the recursion formula

To prove (1.6) we start from the formula

$$(2.1) \quad t_n(k) = \sum_{p=1}^{n-1} \binom{n-1}{p} \sum_{m_1+\dots+m_p=n-1} \frac{(n-1-p)!}{(m_1-1)! \cdots (m_p-1)!} t_{m_1}(k-1) \cdots t_{m_p}(k-1).$$

(2.1) can be proved as follows: Let  $E$  denote the set of those points of  $T \in \mathcal{T}_n$  which are directly connected (i.e. connected by an edge) with  $P_1$ . If  $p$  is the number of elements of  $E$  then  $1 \leq p \leq n-1$  and denoting these points by  $Q_1, \dots, Q_p$ , the points  $Q_i$  can be selected in  $\binom{n-1}{p}$  ways. All the remaining  $n-1-p$  points  $P_m$  ( $P_m \neq Q_j$ ,  $1 \leq j \leq p$ ;  $P_m \neq P_1$ ) can be classified into  $p$  classes which are defined as follows:  $P_m$  lies in the  $j$ -th class ( $j = 1, 2, \dots, p$ ) if the unique path from  $P_1$  to  $P_m$  goes through  $Q_j$ . Clearly if the  $j$ -th class contains  $m_j-1$  points, these points together with  $Q_j$  form a tree of order  $m_j$  and height  $\leq k-1$  over the basic point  $Q_j$ . Thus (2.1) follows.

Multiplication of (2.1) by  $x^n/(n-1)!$  and summation for  $n = 1, 2, \dots$  leads immediately to (1.6). (1.11) can be deduced from (1.6) by using several times the power series of the exponential function. It can also be proved directly as follows:

Let  $T \in \mathcal{T}_n$  be a tree the height of which over the basic point  $P_1$  is  $\leq k$ . Then all points of  $T$  different from  $P_1$  can be classified into  $k$  classes, the  $j$ -th class  $\mathcal{C}_j$  consisting of those points whose distance from  $P_1$  is equal to  $j$  ( $1 \leq j \leq k$ ). Let  $m_j$  denote the number of points in the class  $\mathcal{C}_j$  ( $1 \leq j \leq k$ ); then

$$\sum_{j=1}^k m_j = n-1.$$

If the numbers  $m_j$  are fixed, the distribution of the  $n-1$  points in the classes  $\mathcal{C}_j$  can be carried out in  $(n-1)!/m_1! \cdots m_k!$  ways. Now evidently each point in the class  $\mathcal{C}_1$  is directly connected with  $P_1$ , each point in  $\mathcal{C}_2$  is directly connected with some point in  $\mathcal{C}_1$  etc., each point in  $\mathcal{C}_k$  is directly

connected with some point of  $\mathcal{C}_{k-1}$ . As the connections can be established in  $m_1^{m_2} m_2^{m_3} \cdots m_{k-1}^{m_k}$  different ways and by choosing these connections the tree  $T$  is completely determined, (1.11) follows.

For  $d_n(k) = t_n(k) - t_n(k-1)$  the proof of (1.11) gives

$$(2.2) \quad d_n(k) = \sum_{\substack{m_1 + \cdots + m_k = n-1 \\ m_i \geq 1}} \frac{(n-1)!}{m_1! \cdots m_k!} m_1^{m_2} m_2^{m_3} \cdots m_{k-1}^{m_k}, \quad (i = 1, \dots, k).$$

If  $d_n(k, m)$  denotes the number of trees  $T \in \mathcal{T}_n$  for which  $h_{P_1}(T) = k$  and in which there are exactly  $m$  points connected with  $P_1$  by an edge then (2.2) gives

$$(2.3) \quad d_n(k, m) = \frac{1}{m!} \sum_{\substack{m_1 + \cdots + m_k = n-1-m \\ m_i \geq 1}} \frac{(n-1)!}{m_1! \cdots m_k!} m_1^{m_2} \cdots m_{k-1}^{m_k}.$$

From here the following recursion formula can be deduced:

$$(2.4) \quad d_n(k, m) = \binom{n-1}{m} \sum_{p=1}^{n-1-m} m^p d_{n-m}(k-1, p).$$

Similarly if  $t_n(k, m)$  is the number of those trees  $T \in \mathcal{T}_n$  which have height  $\leq k$  over  $P_1$  and in which the number of points having distance  $k$  from  $P_1$  equals  $m$ , then

$$(2.5) \quad t_n(k, m) = \frac{1}{m!} \sum_{m_1 + \cdots + m_{k-1} = n-1-m} \frac{(n-1)!}{m_1! \cdots m_{k-1}!} m_1^{m_2} \cdots m_{k-2}^{m_{k-1}} m_{k-1}^m,$$

and

$$(2.6) \quad t_n(k, m) = \binom{n-1}{m} \sum_{p=1}^{n-1-m} p^m t_{n-m}(k-1, p).$$

Thus putting

$$(2.7) \quad F_k(x, z) = \sum_{n=1}^{\infty} \sum_{m=0}^{n-1} \frac{t_n(k, m)}{(n-1)!} x^{n-m} z^m$$

we have the recursion formula

$$(2.8) \quad F_k(x, z) = F_{k-1}(x, xe^z)$$

with  $F_0(x, z) = z$ . We obtain

$$F_1(x, z) = xe^z, \quad F_2(x, z) = xe^{xe^z}, \text{ etc.,}$$

hence

$$(2.9) \quad F_{k+1}(x, z) = x \exp F_k(x, z)$$

further

$$(2.10) \quad F_k(x, x) = F_{k+1}(x, 0) = G_k(x).$$

### 3. The asymptotic distribution of $d_n(k)$

We consider now the asymptotic distribution of  $d_n(k)$  when  $n$  and  $k$  are large. We shall make use of the generating function

$$(3.1) \quad G_k(x) - G_{k-1}(x) = \sum_{n=1}^{\infty} \frac{d_n(k)}{(n-1)!} x^n$$

where

$$(3.2) \quad G_0(x) = x, \quad G_k(x) = x \exp G_{k-1}(x) \quad (k = 1, 2, \dots)$$

by (1.5) and (1.15). From (3.2) it is seen that  $G_k(z) - G_{k-1}(z)$  is an entire function and hence

$$(3.3) \quad \frac{d_n(k)}{(n-1)!} = \frac{1}{2\pi i} \int_{C+} \frac{G_k(z) - G_{k-1}(z)}{z^{n+1}} dz$$

where  $C$  is any circular path with centre 0. For the radius of  $C$  we shall take  $r = e^{-1}$ ; this is the largest positive value of  $r$  for which the sequence  $G_k(r)$ ,  $k = 1, 2, \dots$  tends to a finite limit, namely  $\lim_{k \rightarrow \infty} G_k(e^{-1}) = 1$ . Moreover if  $k$  is of order  $\sqrt{n}$  which is the case of principal interest then the point  $e^{-1}$  lies very close to a saddle point of the integrand.

As in (2.9), write for a fixed complex number  $\zeta$

$$(3.4) \quad F_0(\zeta, z) = z,$$

$$(3.5) \quad F_k(\zeta, z) = \zeta \exp F_{k-1}(\zeta, z) \quad (k = 1, 2, \dots).$$

Thus  $F_k(\zeta, z)$  is the  $k$ -th iterate of

$$(3.6) \quad F(\zeta, z) = \zeta e^z \quad (\zeta \text{ fixed})$$

and

$$(3.7) \quad G_k(\zeta) = F_k(\zeta, \zeta) = F_{k+1}(\zeta, 0),$$

as in (2.10).

In the particular case of  $\zeta = e^{-1}$ ,  $z = 1$  is a fixed point with multiplier 1 of the function  $F(e^{-1}, z) = e^{z-1}$ ; in fact  $F(e^{-1}, 1) = 1$ ,  $F'(e^{-1}, 1) = 1$ .

The sequence

$$(3.8) \quad \gamma_k = F_k(e^{-1}, e^{-1}) = G_k(e^{-1}), \quad k = 0, 1, 2, \dots$$

satisfies

$$(3.9) \quad \gamma_k = \exp(\gamma_{k-1} - 1), \quad k = 1, 2, \dots$$

and has an asymptotic expansion

\*  $z = a$  is a fixed-point with multiplier  $\mu$  of the function  $F(z)$  if  $F(a) = a$  and  $F'(a) = \mu$ .  
If  $|\mu| < 1$ , the fixed-point is called attractive. (Fatou [4], p. 186).

$$(3.10) \quad \gamma_k \cong 1 - \frac{2}{k} + \frac{2}{3} \frac{\log k}{k^2} + \frac{c}{k^2} + \dots \quad (k \rightarrow \infty)$$

where  $c$  is a certain constant (see e.g. [3], Lemma 3, p. 247). For all other values of  $\zeta$  on the circle

$$\zeta = e^{-1+it}, \quad -\pi \leq t \leq \pi,$$

$F(\zeta, z)$  has an attractive fixed-point  $\omega = u + iv$  with multiplier  $\omega$  ( $|\omega| < 1$ ), given by the equation

$$(3.11) \quad \omega = \zeta e^\omega = e^{-1+it+\omega}.$$

These fixed-points lie on the curve

$$(3.12) \quad u^2 + v^2 = e^{2(u-1)}, \quad u \leq 1$$

with

$$(3.13) \quad \tan(v+t) = v/u = u^{-1} (e^{2(u-1)} - u^2)^{\frac{1}{2}}.$$

Thus to each  $\zeta = e^{-1+it}$  there corresponds a unique  $\omega = \omega(\zeta) = u + iv$  on the curve (3.12). In the neighbourhood of  $t = 0$ , i.e. of  $u = 1$ ,  $v = 0$ , the curve of fixed-points has a double point and satisfies an expansion

$$(3.14) \quad \begin{aligned} u &= 1 - \sqrt{t} + 0 \cdot t + at\sqrt{t} + \dots, \\ v &= \sqrt{t} - \frac{2}{3}t + bt\sqrt{t} + \dots \quad \text{if } t > 0 \\ &= -\sqrt{-t} - \frac{2}{3}t - bt\sqrt{-t} + \dots \quad \text{if } t < 0. \end{aligned}$$

The fixed-points  $\omega$  on (3.12) are attractive for all  $z$  on the circle  $|z| = e^{-1}$ ; in fact

$$(3.15) \quad |F(\zeta, z)| = |\zeta e^z| < 1$$

for all  $|\zeta| = e^{-1}$ ,  $\operatorname{Re} z < 1$  and the functions

$$F_k(\zeta, z), \quad |\zeta| = e^{-1}, \quad k = 0, 1, 2, \dots$$

form a normal family on the half plane  $\operatorname{Re} z < 1$ .

By a more refined argument one can show that if we set

$$(3.16) \quad D_k(\zeta) = G_k(\zeta) - \omega(\zeta)$$

then  $D_k(\zeta)/\omega(\zeta)^k$  is uniformly bounded for all  $\zeta = e^{-1+it}$ ,  $-\pi \leq t \leq \pi$  and  $k = 0, 1, 2, \dots$ , i.e.

$$(3.17) \quad |D_k(\zeta)| < A |\omega(\zeta)|^k$$

for a suitable positive constant  $A$ . For we have

$$(3.18) \quad D_k(\zeta) = \omega(\zeta) [\exp D_{k-1}(\zeta) - 1], \quad k = 1, 2, \dots$$

by (3.2), (3.11) and (3.16), and the sequence behaves very nearly like the sequence  $D_k^*$  given by the recursion

$$(3.19) \quad D_k^* = \omega D_{k-1}^* / (1 - \frac{1}{2} D_{k-1}^*).$$

For this sequence the statement can be verified by direct calculation since

$$(3.20) \quad D_k^* = - \frac{2\omega^k}{1 + \omega + \cdots + \omega^{k-1} - a}, \quad a = 1/D_0^*.$$

We omit details.

For  $|t| < ((\log^2 k)/k)^2$  the sequence  $G_k(\zeta)$ ,  $k = 0, 1, 2, \dots$  has a uniform asymptotic expansion

$$(3.21) \quad G_k(e^{-1+it}) \cong 1 - \frac{2}{k} \tau \cot \tau + \frac{2}{3} \frac{\tau^2}{\sin^2 \tau} \frac{\log k}{k^2} \\ + \frac{1}{k^2} \left[ c \frac{\tau^2}{\sin^2 \tau} + \frac{2}{3} \tau^2 \left( 2 + \log \frac{\sin \tau}{\tau} \right) \right] + \cdots$$

where

$$(3.22) \quad it = 2\tau^2/k^2$$

and  $c$  is the constant in (3.10). The expansion becomes (3.10) for  $\tau = 0$  and can be verified formally by setting

$$(3.23) \quad G_k(e^{-1+it}) \cong 1 - \frac{\theta_1(\tau)}{k} + \theta_2(\tau) \frac{\log k}{k^2} + \frac{\theta_3(\tau)}{k^2} + \cdots$$

and using (3.2).

We obtain (for fixed  $t$ ) by (3.22)

$$G_{k+1}(e^{-1+it}) \cong 1 - \frac{\theta_1(\tau + \tau/k)}{k+1} + \theta_2(\tau + \tau/k) \frac{\log(k+1)}{(k+1)^2} \\ + \theta_3(\tau + \tau/k) \frac{1}{(k+1)^2} + \cdots \\ \cong 1 - \theta_1/k + \theta_1/k^2 - \theta_1/k^3 - \tau\theta_1'/k^2 \\ + \tau\theta_1'/k^3 - \tau^2\theta_1''/2k^3 + \theta_2 \log k/k^2 \\ - 2\theta_2 \log k/k^3 + \theta_2/k^3 + \tau\theta_2' \log k/k^3 \\ + \theta_3/k^2 - 2\theta_3/k^3 + \tau\theta_3'/k^3 + \cdots, \\ \exp(G_k - 1 + 2\tau^2/k^2) \cong \exp \left[ - \frac{\theta_1}{k} + \theta_2 \frac{\log k}{k^2} + \frac{\theta_3}{k^2} + \frac{2\tau^2}{k^2} + \cdots \right] \\ \cong 1 - \theta_1/k + \theta_2 \log k/k^2 + (\theta_3 + 2\tau^2)/k^2 \\ + \frac{1}{2}\theta_1^2/k^2 - \theta_1\theta_2 \log k/k^3 - \theta_1\theta_3/k^3 \\ - 2\theta_1\tau^2/k^3 - \frac{1}{6}\theta_1^3/k^3 + \cdots$$

where values of the function  $\theta_i$  and their derivatives are taken at  $\tau$ . These two expressions are equal by (3.2), hence comparing coefficients

$$(3.24) \quad \theta_1 - \tau\theta'_1 = \frac{1}{2}\theta_1^2 + 2\tau^2,$$

$$(3.25) \quad \tau\theta'_2 - 2\theta_2 = -\theta_1\theta_2,$$

$$(3.26) \quad -\theta_1 + \tau\theta'_1 - \frac{1}{2}\tau^2\theta''_1 + \theta_2 - 2\theta_3 + \tau\theta'_3 = -\theta_1\theta_3 - 2\tau^2\theta_1 - \frac{1}{6}\theta_1^3.$$

With the initial conditions  $\theta_1(0) = 2$ ,  $\theta_2(0) = \frac{2}{3}$ ,  $\theta_3(0) = c$ , obtained from (3.10), the equations (3.24)–(3.26) give

$$\begin{aligned} \theta_1(\tau) &= 2\tau \cot \tau, \quad \theta_2(\tau) = \frac{2}{3} \frac{\tau^2}{\sin^2 \tau}, \\ \theta_3(\tau) &= c \frac{\tau^2}{\sin^2 \tau} + \frac{2}{3}\tau^2 \left( 2 + \log \frac{\sin \tau}{\tau} \right), \quad \text{i.e. (3.21).} \end{aligned}$$

Thus we only have to prove the existence of an expansion of the form (3.23). This can be achieved by a step by step method such as the one used in [3] for the proof of general expansions of the type (3.11); we omit details. Actually we only need the expansion in the weaker form

$$(3.27) \quad G_k(e^{-1+it}) = 1 - \frac{2}{k} \tau \cot \tau + O(k^{-1-2\delta})$$

for some  $\delta > 0$  when  $|t| \leq ((\log^2 k)/k)^2$ .

We then get from (3.2), since

$$2\tau \cot \tau = O(k \sqrt{|t|}) = O(\log^2 k),$$

$$\begin{aligned} G_k(e^{-1+it}) - G_{k-1}(e^{-1+it}) &= \exp(-1+it+G_{k-1}) - G_{k-1} \\ &= it + \frac{1}{2}(G_{k-1}-1)^2 + O(k^{-3+\delta}) \\ &= 2\tau^2(1+\cot^2 \tau)/k^2 + O(k^{-2-\delta}), \end{aligned}$$

$$(3.28) \quad G_k(e^{-1+it}) - G_{k-1}(e^{-1+it}) = \frac{2}{k^2} \frac{\tau^2}{\sin^2 \tau} + O(k^{-2-\delta})$$

for  $|t| \leq ((\log^2 k)/k)^2$ .

Now from (3.17)

$$\begin{aligned} G_k(e^{-1+it}) - G_{k-1}(e^{-1+it}) &= D_k(e^{-1+it}) - D_{k-1}(e^{-1+it}) \\ &= O\left(\left(1 - \frac{\log^2 k}{k}\right)^k\right) = O(e^{-\log^2 k}) \end{aligned}$$

for  $|t| \geq ((\log^2 k)/k)^2$ , by (3.14), hence by (3.3) and (3.28)



$$\begin{aligned}
 (3.29) \quad \frac{d_n(k)}{(n-1)!} &= \frac{1}{2\pi i} \frac{2e^n}{k^2} \left( \int_{|t| \leq ((\log^2 k)/k)^{1/2}} \frac{\tau^2}{\sin^2 \tau} e^{-nit} idt + O(k^{-2-\delta}) \right) \\
 &\cong \frac{1}{2\pi i} \frac{8e^n}{k^4} \int_{\Gamma} \frac{\tau^3}{\sin^2 \tau} e^{-\beta \tau^2} d\tau
 \end{aligned}$$

by (3.22), where

$$(3.30) \quad \beta = 2n/k^2$$

and  $\Gamma$  is the path

$$\begin{aligned}
 \tau &= (i-1)u, \quad -\frac{1}{2} \log^2 k \leq u \leq 0 \\
 \tau &= (i+1)u, \quad 0 \leq u \leq \frac{1}{2} \log^2 k.
 \end{aligned}$$

Hence

$$\frac{d_n(k)}{(n-1)!} \cong \frac{8e^n}{k^4} \sum_{p=1}^{\infty} \operatorname{res}_{\tau=p\pi} \left( \frac{\tau^3}{\sin^2 \tau} e^{-\beta \tau^2} \right).$$

By using Stirling's formula we obtain from here

$$(3.31) \quad p_n(k) = \frac{d_n(k)}{n^{n-2}} \cong 2 \left( \frac{2\pi}{n} \right)^{\frac{1}{2}} \beta^2 \sum_{p=1}^{\infty} (2p^4 \pi^4 \beta - 3p^2 \pi^2) e^{-\beta \pi^2 p^2}$$

for large  $n$  and  $k$  where  $\beta$  is given by (3.30).

This is the required asymptotic probability distribution. Note that

$$\begin{aligned}
 \sum_{k=1}^{n-1} p_n(k) &\sim 2 \left( \frac{2\pi}{n} \right)^{\frac{1}{2}} \int_0^{n-1} \beta^2 \sum_{p=1}^{\infty} (2p^4 \pi^4 \beta - 3p^2 \pi^2) e^{-\beta \pi^2 p^2} dk \\
 &\cong 2\pi^{\frac{1}{2}} \int_0^{\infty} \sum_{p=1}^{\infty} \beta^{\frac{1}{2}} (-3p^2 \pi^2 + 2p^4 \pi^4 \beta) e^{-\beta \pi^2 p^2} d\beta \\
 &= 2\pi^{\frac{1}{2}} \lim_{\epsilon \rightarrow 0} \left[ - \sum_{p=1}^{\infty} 2p^2 \pi^2 \beta^{\frac{3}{2}} e^{-\beta \pi^2 p^2} \right]_{\epsilon}^{\infty} \\
 &= \lim_{\beta \rightarrow 0} 4\pi^{\frac{1}{2}} \beta^{\frac{3}{2}} \sum_{p=1}^{\infty} p^2 e^{-\beta \pi^2 p^2} \\
 &= 4\pi^{-\frac{1}{2}} \int_0^{\infty} u^2 e^{-u^2} du = 1,
 \end{aligned}$$

as required.

The maximum of the distribution curve is reached when in (3.31),  $(d/d\beta)p_n(k) = 0$ , i.e.

$$\sum_{p=1}^{\infty} (9p^4 \pi^4 \beta^2 - 6p^2 \pi^2 \beta - 2p^6 \pi^6 \beta^3) e^{-\beta \pi^2 p^2} = 0.$$

Numerically  $\beta(\max) = 0.373138525$  and

$$(3.32) \quad k(\max) = 2.31515436 \sqrt{n}.$$

#### 4. Conclusion

The result of the previous section can be stated as follows:

Let  $\mathcal{H}_n$  be the height over  $P_1$  of a labelled random tree of order  $n$  i.e. of a tree selected at random from the set of  $n^{n-2}$  elements of  $\mathcal{T}_n$  with uniform probability distribution. Then

$$(4.1) \quad \lim_{n \rightarrow \infty} P\left(\frac{\mathcal{H}_n}{\sqrt{2n}} < x\right) = H(x)$$

where

$$(4.2) \quad H(x) = 4x^{-3} \pi^{\frac{1}{2}} \sum_{p=1}^{\infty} p^2 e^{-\pi^2 p^2 / x^2}.$$

This can be transformed (e.g. by means of Poisson's formula) to the form

$$(4.3) \quad H(x) = \sum_{v=-\infty}^{\infty} e^{-v^2 x^2} (1 - 2v^2 x^2)$$

whence

$$(4.4) \quad h(x) = H'(x) = 4x \sum_{v=1}^{\infty} v^2 (2v^2 x^2 - 3) e^{-v^2 x^2}.$$

From (4.4) we can calculate all moments of the distribution function  $H(x)$ :

$$(4.5) \quad \begin{aligned} M_s &= \int_0^{\infty} x^s h(x) dx \\ &= 2\Gamma(\tfrac{1}{2}s + 1)(s-1)\zeta(s) \end{aligned} \quad (s > 1)$$

where  $\zeta(s) = \sum_{m=1}^{\infty} 1/m^s$ . In particular we obtain for  $M_1$ , since

$$\lim_{s \rightarrow 1} (s-1)\zeta(s) = 1,$$

$$(4.6) \quad M_1 = \sqrt{\pi}.$$

Hence the expectation value of  $\mathcal{H}_n$  is

$$(4.7) \quad E(\mathcal{H}_n) \sim \sqrt{2n\pi} = 2.50663\sqrt{n}$$

and the variance is

$$(4.8) \quad D^2 = M_2 - M_1^2 = \frac{\pi(\pi-3)}{3}.$$

#### References

- [1] Cayley, A., 'A theorem on trees', *Quart. J. of Pure and Applied Math.* **23** (1889), 376–378.
- [2] Riordan, J., 'The enumeration of trees by height and diameter', *IBM Journal of Research and Development* **4** (1960), 473–478.

- [3] Szekeres, G., 'Regular iteration of real and complex functions', *Acta Math.* 100 (1958), 103–258.
- [4] Fatou, M. P., 'Sur les équations fonctionnelles', *Bull. Soc. Math. France* 47 (1919), 161–271.

Mathematical Institute of the Academy, Budapest  
and  
University of New South Wales