

AST4320 Oblig2

Daniel Heinesen, daniehei

28. oktober 2019

1 1

1.1 a)

To find the mean molecular weight if the intergalactic medium (IGM) we use that

$$\frac{1}{\mu} = \sum_i \frac{X_i}{A_i} = 2X + 4/3 \cdot Y, \quad (1)$$

where X_i is the fraction of species i and A_i is the atomic mass of said species divided by the total number of particles it ionizes to. For the IGM we have a fraction of hydrogen $X = 0.76$ and a fraction of helium $Y = 0.24$, giving us

$$\mu = \frac{1}{2 \cdot 0.76 + 4/3 \cdot 0.24} = 0.59. \quad (2)$$

1.2 b)

The Jeans length is given as

$$\lambda_j = c_s \left(\frac{\pi}{G\rho} \right)^{1/2}, \quad (3)$$

where the speed of sound c_s for a gas is given as

$$c_s = \sqrt{\frac{k_b T}{\mu m_p}} = 11.83 \text{ m/s}. \quad (4)$$

We are using that $T = 10^4$ K. We have to find the pressure of the IGM. For this we assume that the gas only have baryonic matter, so

$$\rho = \rho_b(1+z)^3 = \Omega_b \rho_c(1+z)^3, \quad (5)$$

where $\Omega_b = 0.049$ and $\rho_c = 9.47 \cdot 10^{-27} \text{ g/cm}^3$. We then see that get

$$\lambda_j(z) = 3.9 \text{ Mpc} \cdot (1+z)^{-3/2}. \quad (6)$$

This gives k

$$k = \frac{2\pi}{\lambda_j(z)} = 1.61 \text{ Mpc}^{-1}(1+z)^{3/2}. \quad (7)$$

1.3 c)

We can use Hubble's law to find the different in velocity across the Jeans length of the IGM, thereby giving us the velocity width. Through Hubble's law we get

$$\Delta v = H(z)\lambda_j, \quad (8)$$

where

$$H(z) = H_0(\Omega_m(1+z)^3 + \Omega_r(1+z)^4 + \Omega_\Lambda)^{1/2}. \quad (9)$$

This gives us that

$$\Delta v = H_0(\Omega_m(1+z)^3 + \Omega_r(1+z)^4 + \Omega_\Lambda)^{1/2} 3.9 \text{ Mpc} \cdot (1+z)^{-3/2} \quad (10)$$

$$= 264.5 \text{ km/s} \cdot (\Omega_m + \Omega_r(1+z) + \Omega_\Lambda(1+z)^{-3})^{1/2} = 264.5 \text{ km/s} \cdot (0.308 + 0.692(1+z)^{-3})^{1/2} \quad (11)$$

1.4 d)

See over
this!

1.5 e)

The thermal broadening scale is given as

$$v_{th} = \sqrt{\frac{2k_b T}{m_p}}, \quad (12)$$

where $T = 10^4$ K. This gives us

$$v_{th} = 12.85 \text{ km/s}, \quad (13)$$

which is an order small that all values of z .

2 Exercise 2

We would like to find the optical depth τ_e of the IGM as a function of redshift z . The optical depth is given as


$$\tau_e(z) = c \int_0^z \frac{n_e(z) \sigma_T dz}{H(z)(1+z)}, \quad (14)$$

where σ_t is the Thompson cross section, n_e the electron density and $H(z)$ the Hubble parameter. Since the Universe is taken to be completely ionized, he assume that $n_e \approx n_H = 1.9 \cdot 10^{-7} (1+z)^3 \text{ cm}^{-3}$. We get the Hubble parameter from the Friedmann equation

$$H(z) = H_0 \sqrt{\Omega_m (1+z)^3 + \Omega_r (1+z)^4 + \Omega_\Lambda}, \quad (15)$$

where $\Omega_\Lambda = 0.692$, $\Omega_m = 0.308$ and $\Omega_r = 0$.

We will integrate (14) from $z = 0$ to 10. We assume that $\tau_e(0) \approx 0$.



Figur 1: The optical depth of the intergalactic medium. We see that for larger redshift the optical depth increases.

3 Exercise 3

3.1 a)

We have the differential equation for an isothermal halo

$$-\frac{k_b T}{m_{DM} r^2} \frac{d}{dr} r^2 \frac{d}{dr} \ln \rho = 4\pi G \rho. \quad (16)$$

We have an ansatz that

$$\rho(r) = \frac{A}{r^2}, \quad A = \frac{k_b T}{2\pi G m_{DM}}. \quad (17)$$

To see that this is a solution, we put (17) into (16). Looking at the RHS of (16) we get

$$-\frac{k_b T}{m_{DM} r^2} \frac{d}{dr} r^2 \frac{d}{dr} \ln \rho = -\frac{k_b T}{m_{DM} r^2} \frac{d}{dr} r^2 \frac{r^2}{A} \cdot \left(-\frac{2A}{r^3} \right) \quad (18)$$

$$2 \frac{k_b T}{m_{DM} r^2} \frac{d}{dr} r = 2 \frac{k_b T}{m_{DM} r^2}. \quad (19)$$

Thus we get

$$2 \frac{k_b T}{m_{DM} r^2} = 4\pi G \frac{A}{r^2} \Rightarrow A = \frac{k_b T}{2\pi G m_{DM}}. \quad (20)$$

Thus (17) solves (16).

3.2 b)

We have that a gas in hydrostatic equilibrium

$$\frac{dp}{dr} = -\frac{GM(< r)\rho}{r^2}, \quad (21)$$

where $M(< r)$ is the mass within a radius r and p is the pressure. We can see that our isothermal gas, with the density defined in (17), behaves in the similar way. We start by finding the mass, which for a spherical symmetric mass is given as

$$M(< r) = 4\pi \int_0^r \rho(r')r'^2 dr' = 4\pi \int_0^r \frac{A}{r'^2} r'^2 dr' = 4\pi \int_0^r A dr' = 4\pi Ar. \quad (22)$$

In our expression of A we have the dark matter mass m_{DM} . Since we now have a gas, we let $m_{DM} \rightarrow m_p$, which is the proton mass. Thus we have

$$\rho = \frac{A}{r^2} = \frac{k_b T}{2\pi G m_p r^2}. \quad (23)$$

We then use that for an isothermal gas the pressure is given as

$$p = \frac{k_b T}{m_p} \rho = \frac{k_b T A}{m_p r^2}. \quad (24)$$

We can now take the differentiation of this with respect to r and use (22) and (17) to find

$$\frac{dp}{dr} = -\frac{2k_b T A}{m_p r^3} = -\frac{2k_b T}{m_p r^3} \cdot \frac{M(< r)}{4\pi r} = -\frac{GM(< r)}{r^2} \frac{k_b T}{2\pi G m_p r^2} = -\frac{GM(< r)\rho}{r^2}. \quad (25)$$

This is the same as for the gas in hydrostatic equilibrium from (21).

4 Exercise 3

4.1 a)

We have that for cusp density profile

$$\rho^{cusp}(r) = \begin{cases} \rho_0 \left(\frac{r}{r_s}\right)^{-1} & \text{if } r < r_s \\ \rho_0 \left(\frac{r}{r_s}\right)^{-3} & \text{if } r \geq r_s \end{cases} \quad (26)$$

And for the cored profile, we have

$$\rho^{cusp}(r) = \begin{cases} \rho_0 & \text{if } r < r_s \\ \rho_0 \left(\frac{r}{r_s}\right)^{-3} & \text{if } r \geq r_s \end{cases} \quad (27)$$

The mass of a spherical symmetric system can be written as

$$M(< r) = 4\pi \int_0^r \rho(r')r'^2 dr'. \quad (28)$$

So for $r < r_s$ we get

$$M^{cusp}(< r) = 4\pi \int_0^r \rho_0 \left(\frac{r'}{r_s}\right)^{-1} r'^2 dr' = 4\pi r_s \rho_0 \int_0^r r' dr' = 2\pi r_s \rho_0 r^2, \quad (29)$$

and

$$M^{core}(< r) = 4\pi \int_0^r \rho_0 r'^2 dr' = \frac{4}{3}\pi \rho_0 r^3. \quad (30)$$

For $r \geq r_s$ we integrate from r_s to r and include all the mass that was within r_s , so

$$M^{cusp}(< r) = M(< r_s) + 4\pi \int_{r_s}^r \rho_0 \left(\frac{r'}{r_s}\right)^{-3} r'^2 dr' = M(< r_s) + 4\pi r_s^3 \rho_0 \int_0^r r'^{-1} dr' \quad (31)$$

$$= M(< r_s) + 4\pi r_s^3 \rho_0 \ln r \Big|_{r_s}^r = 2\pi r_s \rho_0 r_s^2 + 4\pi r_s^3 \rho_0 (\ln r - \ln r_s), \quad (32)$$

and similarly for the core mass

$$M^{core}(< r) = M^{core}(< r_s) + 4\pi \int_{r_s}^r \rho_0 \left(\frac{r'}{r_s}\right)^{-3} r'^2 dr' = \frac{4}{3}\pi \rho_0 r_s^3 + 4\pi r_s^3 \rho_0 (\ln r - \ln r_s). \quad (33)$$

So we get that

$$M^{cusp}(< r) = \begin{cases} 2\pi r_s \rho_0 r^2 & \text{if } r < r_s \\ 2\pi r_s \rho_0 r_s^2 + 4\pi r_s^3 \rho_0 (\ln r - \ln r_s) & \text{if } r \geq r_s \end{cases} \quad (34)$$

and

$$M^{core}(< r) = \begin{cases} \frac{4}{3}\pi \rho_0 r^3 & \text{if } r < r_s \\ \frac{4}{3}\pi \rho_0 r_s^3 + 4\pi r_s^3 \rho_0 (\ln r - \ln r_s) & \text{if } r \geq r_s \end{cases} \quad (35)$$

4.2 b)

We get the gravitation potential energy by integrating over the energies felt by a small shell of mass from the mass inside it. First we get the energy of one such shell

$$dW(r) = -\frac{GM(< r)m_{shell}}{r_{shell}} = -4\pi GM(< r)\frac{\rho r_{shell}^2 dr}{r_{shell}} = -4\pi GM(< r)\rho r_{shell} dr, \quad (36)$$

where we have used that

$$m_{shell} = \rho V_{shell} = 4\pi \rho r_{shell}^2 dr. \quad (37)$$

Thus the gravitational potential energy is

$$W = \int_0^{r_{vir}} dW = -4\pi G \int_0^{r_{vir}} M(< r)\rho(r)r dr. \quad (38)$$

4.3 c)

We can now find the minimal energy needed to form the cored profile, given as

$$\Delta E = \frac{W^{core} - W^{cusp}}{2} = -2\pi G \int_0^{r_{vir}} (\rho^{core} M^{core}(< r) - \rho^{cusp} M^{cusp}(< r)) r dr \quad (39)$$

We split this into two parts. First we look at $r = 0$ to r_s

$$-2\pi G \int_0^{r_s} (\rho^{core} M^{core}(< r) - \rho^{cusp} M^{cusp}(< r)) r dr = -2\pi G \int_0^{r_s} \left(\rho_0 \frac{4}{3}\pi \rho_0 r^3 - \rho_0 \left(\frac{r}{r_s}\right)^{-1} 2\pi r_s \rho_0 r^2 \right) r dr \quad (40)$$

$$= -2\pi^2 G \rho_0^2 \int_0^{r_s} \left(\frac{4}{3}r^4 - 2r^2 r_s^2 \right) dr = -2\pi^2 G \rho_0^2 \left(\frac{4}{15}r_s^5 - \frac{2}{3}r_s^5 \right) = \frac{4}{5}\pi^2 G \rho_0^2 r_s^5. \quad (41)$$

We then look from r_s to r_{vir}

$$-2\pi G \int_{r_s}^{r_{vir}} (\rho^{core} M^{core}(< r) - \rho^{cusp} M^{cusp}(< r)) r dr \quad (42)$$

$$= -2\pi G \int_{r_s}^{r_{vir}} \left(\rho_0 \left(\frac{r}{r_s} \right)^{-3} \left(\frac{4}{3} \pi \rho_0 r_s^3 + 4\pi r_s^3 \rho_0 (\ln r - \ln r_s) \right) - \rho_0 \left(\frac{r}{r_s} \right)^{-3} (2\pi \rho_0 r_s^3 + 4\pi r_s^3 \rho_0 (\ln r - \ln r_s)) \right) r dr \quad (43)$$

$$= -2\pi G \int_{r_s}^{r_{vir}} \left(\rho_0 \left(\frac{r}{r_s} \right)^{-3} \left(\frac{4}{3} \pi \rho_0 r_s^3 \right) - \rho_0 \left(\frac{r}{r_s} \right)^{-3} (2\pi \rho_0 r_s^3) \right) r dr \quad (44)$$

$$= 2\pi^2 G \rho_0^2 r_s^6 \int_{r_s}^{r_{vir}} \frac{1}{r^2} \frac{2}{3} dr = \frac{4}{3} \pi^2 G \rho_0^2 r_s^6 \frac{-1}{1} (r_{vir}^{-1} - r_s^{-1}) \approx \frac{4}{3} \pi^2 G \rho_0^2 r_s^5, \quad (45)$$

where the last step is because $r_s \ll r_{vir}$. We then get

$$\Delta E = \frac{4}{5} \pi^2 \rho_0^2 G r_s^5 + \frac{4}{3} \pi^2 G \rho_0^2 r_s^5 = \frac{32}{15} \pi^2 G \rho_0^2 r_s^5. \quad (46)$$

4.4 d)

We are now looking at a dwarf galaxy at $z = 0$ with $M_{vir} = 3 \cdot 10^{10} M_\odot$ and $R_{vir} = 45$ kpc, with $r_s = 1$ kpc. With this mass and radius, we can find

$$\rho_0 = \frac{M_{vir}}{4/3 r_s^3 \pi + 4\pi r_s^3 (\ln R_{vir} - \ln r_s)} = 6.13 \cdot 10^{-23} \text{ g cm}^{-3}. \quad (47)$$

This gives us that

$$\Delta E = \frac{32}{15} \pi^2 G \rho_0^2 r_s^5 = 1.477 \cdot 10^{57} \text{ ergs}. \quad (48)$$

Check
this!!!

4.5 e)