

FYS3150 - Project 2

Daniel Heinesen, Halvard Sutterud, Gunnar Lange

October 6, 2016

Abstract

We investigate a numerical solution to Schroedinger's equation for two electrons confined in a three-dimensional harmonic oscillator well. We present a solution to the case where we disregard the interaction between the electrons, and a solution where we include a repulsive (Coloumb) interaction between the electrons.

Contents

1	Introduction	2
2	Theoretical model	2
2.1	Introducing the relevant Schroedinger equation	2
2.2	Reformulating the equation as an Eigenvalue problem	3
2.3	Jacobi's algorithm for solving an eigenvalue problem	3
2.4	Investigating the properties of orthogonal transformations	3
3	Results	4
3.1	Non-interacting	4
3.2	Interacting	4

1 Introduction

The study of electrons confined in a potential trap is a highly active research field today **INSERT SOME REFERENCES HERE**, due to its potential applicability in solid-state physics. Some proposed applications for these systems include semi-conductors and quantum gates for future quantum computers, as well as nano-medical applications.

We investigate Schroedinger's equation for two non-interacting electrons confined in a three-dimensional potential well, and reformulate it as an Eigenvalue problem. We then solve this problem by use of Jacobi's method for finding eigenvalues and eigenvectors. Once we have developed this general method, it is straightforward to implement the potential for the interacting electrons.

2 Theoretical model

2.1 Introducing the relevant Schroedinger equation

A three-dimensional potential well is described by the harmonic oscillator potential:

$$V(r) = \frac{1}{2}kr^2 \quad (1)$$

Here r is the distance from the origin, and k is a constant describing the steepness of the well. Note how this potential is radially symmetric. This means that we can ignore the angular parts of the Schroedinger's equation, leaving us with the radial equation:

$$-\frac{\hbar^2}{2m} \left(\frac{1}{r^2} \frac{d}{dr} r^2 \frac{d}{dr} - \frac{l(l+1)}{r^2} \right) R(r) + V(r)R(r) = ER(r) \quad (2)$$

Here E is the energy of the harmonic oscillator, and l are orbital momentum numbers¹. Substituting $R(r) = \frac{u(r)}{r}$ as done **INSERT REFERENCE HERE**, we obtain:

$$-\frac{\hbar^2}{2m} \frac{d^2}{dr^2} u(r) + \left(V(r) + \frac{l(l+1)}{r^2} \frac{\hbar^2}{2m} \right) u(r) = E u(r) \quad (3)$$

From this transformation, it also follows that the boundary conditions are necessarily $u(0) = u(\infty) = 0$.

2.2 Reformulating the equation as an Eigenvalue problem

After some rescaling of equation 3 described **HERE**, we can reformulate equation 3 as:

$$-\frac{d^2}{d\rho^2} u(\rho) + V(\rho) u(\rho) = \lambda u(\rho) \quad (4)$$

Where ρ is a dimensionless quantity describing a characteristic length and λ are the (sought-after) eigenvalues of the system. V is the chosen potential. To reformulate this as an Eigenvalue problem, we discretize the second derivative. This implies that we write the continuous function $u(\rho)$ as a function at discrete points, $u(\rho_i) = u_i$, with $i = 0, 1, \dots, N$. We can then expand the second derivative in terms of Taylor polynomials to get an iterative scheme for solving the differential equation. The detailed procedure can be found **HERE**, with the final result:

$$d_i u_i + e_{i-1} u_{i-1} + e_{i+1} u_{i+1} = \lambda u_i \quad (5)$$

Where $d_i = 2/h^2 + V_i$ and $e_i = -1/h^2$, with h being the step length (distance between ρ_i and ρ_{i+1}). Note how every e is the same, i.e. we can define $e_i = e$. This can now be reformulated as an eigenvalue problem with a tridiagonal matrix, \mathbf{A} that is:

$$\mathbf{A}\mathbf{u} = \lambda\mathbf{u} \quad (6)$$

The matrix \mathbf{A} is found by inserting the u and e from above, giving:

$$\mathbf{A} = \begin{bmatrix} d_0 & e & 0 & 0 & \dots & 0 & 0 \\ e & d_1 & e & 0 & \dots & 0 & 0 \\ 0 & e & d_2 & e & 0 \dots & 0 & \\ \dots & \dots & \dots & \dots & \dots & \dots & \dots \\ 0 & \dots & \dots & \dots & e & d_{N-1} & e \\ 0 & \dots & \dots & \dots & \dots & e & d_N \end{bmatrix}$$

Note, however, that u_0 and u_N are known, and thus we can remove those rows and columns from the matrix.

2.3 Jacobi's algorithm for solving an eigenvalue problem

Equation 6 can be solved via a general algorithm known as Jacobi's method. Details of this method can be found **HERE**, but the general idea is rather straightforward.

2.4 Investigating the properties of orthogonal transformations

We will here briefly investigate some properties of orthogonal transforms, to arrive at possible unit tests to implement in our programs. We will show that this transformation preserves the dot product and orthogonality. Thus let \mathbf{v}_i and \mathbf{v}_j be two arbitrary vectors. Consider an orthogonal transformation matrix, \mathbf{U} , that is $\mathbf{U}^T = \mathbf{U}^{-1}$. Define $\mathbf{w}_i = \mathbf{U}\mathbf{v}_i$ and $\mathbf{w}_j = \mathbf{U}\mathbf{v}_j$. Consider:

$$\mathbf{w}_i \cdot \mathbf{w}_j = (\mathbf{U}\mathbf{v}_i)^T (\mathbf{U}\mathbf{v}_j)$$

¹Our background in quantum mechanics being nonexistent, we assume this sentence to be meaningful.




Figure 1: Plots showing the probability distribution for the relative position of the two electrons. Omega corresponds to the frequency of the harmonic oscillator potential, which can be seen as the width of the potential well.

Where the last equality follows from the definition of the inner product. Using the properties of the transpose, the first parenthesis can be expanded as:

$$(\mathbf{U}\mathbf{v}_i)^T = \mathbf{v}_i^T \mathbf{U}^T$$

Multiplying this out gives:

$$\mathbf{w}_i \cdot \mathbf{w}_j = \mathbf{v}_i^T \mathbf{U}^T \mathbf{U} \mathbf{v}_j = \mathbf{v}_i^T \mathbf{I} \mathbf{v}_j = \mathbf{v}_i \cdot \mathbf{v}_j$$

This shows immediately that the orthogonal transformation preserves the dot product. It also shows, however, that it preserves orthogonality. To see this, assume therefore that we have a basis of orthonormal vectors, $\mathcal{B} = \{\mathbf{v}_i\}$, i.e. that $\mathbf{v}_i \cdot \mathbf{v}_j = \delta_{ij}$. Inserting this into the last expression gives:

$$\mathbf{w}_i \cdot \mathbf{w}_j = \delta_{ij}$$

Which shows that orthogonal transformations also preserve orthogonality between vectors.

3 Results

After solving the the Schrödingers equation for both the 2 interacting electrons and 2 non-interacting electrons, for four different frequencies for the harmonic oscillator ω_r , we get these graphs(FIG1).

As can be seen, the larger ω_r is, the closer $|u(\rho)|^2$ for the interacting and non-interacting electrons are; $|u(\rho)|^2$ being the probability distribution for the relative position of the two electrons. We can interpret these results by looking at the different potential energy in the two cases:

3.1 Non-interacting

In the non-interaction case the only potential energy is the harmonic oscillator potential, given dimensionless as: (IKKE SIKKERT VI TRENGER Å GJENGI FORMELENE HER, OM DE ER DEFINERT TIDLIGERE I ARTIKKELEN)

$$V(\rho) = \omega \rho^2$$

ω will determine the width of this potential well, and therefor determine the probability of the relative distance for the electron: the greater ω is, the smaller the narrower the potential well is and the closer the electrons are probable to be eachother. This is also what we see in the results.

3.2 Interacting

In the interacting case, we have two potentials competing, the harmonic oscillator potential (HO) and the coulomb potential, given by

$$V(\rho) = \underbrace{\omega\rho^2}_{HO} + \underbrace{1/\rho}_{Coulomb}$$

As mentioned above, for larger ω the closer the interacting solution is to the non-interacting solution. If we look at the two parts of the potential, we can see that for small ρ , the Coulomb potential is dominant, and for larger ρ , the HO is dominant. We can find that the ρ where domain where the coulomb potential is dominant is

$$\rho < \omega_r^{-\frac{1}{3}}$$

This means that the larger ω_r is, the larger the domain where the coulomb potential is dominant is, and thus we expect a greater probability for a greater relative distance between the two electrons. This is exactly what can be seen from the results. For the larger ω 's the HO well is quite small, and is therefore wholly dominant, causing the solutions for both potentials to be similar. For smaller ω 's the HO wells are large and the electrons have the possibility to have a large relative distance from each other.