

FYS3120 Oblig 4

Daniel Heinesen, daniehei

16. februar 2017

1)

a)

We are going to start by finding the coordinates, we only need to find y , since we need it to find the potential energy

$$y = -\frac{b}{2} \cos \theta$$

The potential energy is then given by

$$V = mgy = -mg\frac{b}{2} \cos \theta$$

There is only rotational energy for the rod, it has a inertia of $I = \frac{1}{3}mb^2$. So its kinetic energy is given as

$$T = \frac{1}{2}I\omega^2 = \frac{1}{6}mb^2\dot{\theta}^2$$

Giving us the Lagrangian

$$L = T - V = \frac{1}{6}mb^2\dot{\theta}^2 + mg\frac{b}{2} \cos \theta$$

We can now find the e.o.m

$$\frac{\partial L}{\partial \theta} = -\frac{mgb}{2} \sin \theta$$
$$\frac{\partial L}{\partial \dot{\theta}} = \frac{1}{3}mb^2\dot{\theta}, \quad \frac{d}{dt} \left(\frac{\partial L}{\partial \dot{\theta}} \right) = \frac{1}{3}mb^2\ddot{\theta}$$

Giving us that

$$\frac{1}{3}mb^2\ddot{\theta} + \frac{mgb}{2} \sin \theta = 0$$

or

$$\ddot{\theta} + \frac{3g}{2b} \sin \theta = 0$$

b)

We can see that for $\theta = 0$

$$\ddot{\theta} = 0$$

So this is a equilibrium. This is a minimum or maximum of the potential. We need to differentiate the potential twice to see if this is a minimum

$$\frac{d^2}{d\theta^2} - \frac{mgb}{2} \cos \theta = \frac{mgb}{2} \cos \theta$$

at $\theta = 0$

$$\Rightarrow \frac{mgb}{2} \cos(0) = \frac{mgb}{2} > 0$$

This shows that we have a minimum, and the equilibrium is thus stable.

We can now look at the e.o.m for small angles. Then $\sin \theta \approx \theta$, so

$$\ddot{\theta} = -\frac{3g}{2b}\theta$$

This is harmonic oscillator with

$$\omega^2 = \frac{3g}{2b}$$

Using that $T = \frac{2\pi}{\omega}$ we find that

$$T_0 = 2\pi\sqrt{\frac{2b}{3g}}$$

c)

Since

$$\frac{\partial L}{\partial t} = 0$$

and

$$\frac{dH}{dt} = -\frac{\partial L}{\partial t} = 0$$

We can see that the Hamiltonian is a constant of motion. Since the constraints are holonomic and time-independent, we know that the Hamiltonian is the same as the total energy of the system. Thus the total energy is conserved.

The Hamiltonian is related to the e.o.m by Hamilton's equations

$$\dot{p} = -\frac{\partial H}{\partial \theta}, \quad \dot{\theta} = \frac{\partial H}{\partial p}$$

d)

This derivation is heavily inspired by a similar derivation for a normal pendulum, found on wikipedia.

We are going to start by looking at the energy of the pendulum. The pendulum has no kinetic energy at θ_0 and potential $mg \cos \theta_0$. At some θ the kinetic energy has to be the difference in potential energy between these angles. So

$$\frac{1}{6}mb^2\dot{\theta}^2 = \frac{mgb}{2}(\cos \theta - \cos \theta_0)$$

giving us that

$$\dot{\theta} = \frac{d\theta}{dt} = \sqrt{\frac{3g}{b}}\sqrt{(\cos \theta - \cos \theta_0)}$$

Doing some physicist math we get that

$$dt = \sqrt{\frac{b}{3g}} \frac{d\theta}{\sqrt{\cos \theta - \cos \theta_0}}$$

The pendulum needs to fall from 0 to θ_0 , then to $-\theta_0$, and then all the way back to θ_0 , meaning that the period is 4 times the time it take to fall from 0 to θ_0 . Meaning that

$$T = 4 \int dt = 4 \sqrt{\frac{b}{3g}} \int_0^{\theta_0} \frac{d\theta}{\sqrt{\cos \theta - \cos \theta_0}}$$

And we know that $T_0 = 2\pi\sqrt{\frac{2b}{3g}}$, giving us that

$$T = T_0 \frac{\sqrt{2}}{\pi} \int_0^{\theta_0} \frac{d\theta}{\sqrt{\cos \theta - \cos \theta_0}}$$

We can now solve this for $\theta_0 = \pi/2$

$$T = T_0 \frac{\sqrt{2}}{\pi} \int_0^{\pi/2} \frac{d\theta}{\sqrt{\cos \theta}}$$

From good old Rottmann page 161 eq. 116, we get that this is

$$T = T_0 \frac{\sqrt{2}}{\pi} \frac{\pi \Gamma(\frac{1}{2})}{\sqrt{2} \Gamma(\frac{3}{4})^2} = T_0 \frac{\sqrt{\pi}}{\Gamma(\frac{3}{4})^2} \approx 1.18 \cdot T_0$$

2)

a)

First we write the coordinates of the ball

$$x = r \sin \theta \cos(\omega t), \quad y = r \sin \theta \sin(\omega t), \quad z = -r \cos \theta$$

$$\dot{x} = r\dot{\theta} \cos \theta \cos(\omega t) - r\omega \sin \theta \sin(\omega t)$$

$$\dot{y} = r\dot{\theta} \cos \theta \sin(\omega t) - r\omega \sin \theta \cos(\omega t)$$

$$\dot{z} = r\dot{\theta} \sin \theta$$

with out writing down the quite long expressions for \dot{x} , \dot{y} and \dot{z} , we get that

$$\dot{x}^2 + \dot{y}^2 + \dot{z}^2 = r^2 \dot{\theta}^2 + r^2 \omega^2 \sin^2 \theta$$

We have the energies

$$V = mgy = -mgr \cos \theta$$

$$T = \frac{1}{2}m(r^2 \dot{\theta}^2 + r^2 \omega^2 \sin^2 \theta)$$

Giving us the Lagrangian

$$L = \frac{1}{2}m(r^2 \dot{\theta}^2 + r^2 \omega^2 \sin^2 \theta) + mgr \cos \theta$$

We can rewrite this as

$$L = \frac{1}{2}mr^2 \dot{\theta}^2 + \left(\frac{1}{2}mr^2 \omega^2 \sin^2 \theta + mgr \cos \theta\right)$$

We can therefor say that we have a potential

$$W(\theta) = -\left(\frac{1}{2}mr^2\omega^2 \sin^2 \theta + mgr \cos \theta\right)$$

The term

$$-\frac{1}{2}mr^2\omega^2 \sin^2 \theta$$

Can be said to be the potential energy due to the centripetal force.

b)

We can find the e.o.m

$$\frac{\partial L}{\partial \theta} = mr^2\omega^2 \sin \theta \cos \theta - mgr \sin \theta = \frac{1}{2}mr^2\omega^2 \sin 2\theta - mgr \sin \theta$$

$$\frac{\partial L}{\partial \dot{\theta}} = mr^2\dot{\theta}, \quad \frac{d}{dt} \left(\frac{\partial L}{\partial \dot{\theta}} \right) = mr^2\ddot{\theta}$$

Giving us that

$$mr^2\ddot{\theta} - \frac{1}{2}mr^2\omega^2 \sin 2\theta + mgr \sin \theta = 0$$

We can look at this for small angles. For small angles $\sin \theta \approx \theta$, so

$$mr^2\ddot{\theta} - mr^2\omega^2 \theta + mg\theta = 0$$

$$\Rightarrow \ddot{\theta} = -\left(\frac{g}{r} - \omega^2\right) \theta$$

This is a harmonic oscillation, giving us

$$\Omega = \sqrt{\frac{g - r\omega^2}{r}}$$

c)

We have the potential

$$W(\theta) = -\left(\frac{1}{2}mr^2\omega^2 \sin^2 \theta + mgr \cos \theta\right)$$

To find the equilibrium positions we have to differentiate

$$W'(\theta) = -mr^2\omega^2 \sin \theta \cos \theta + mgr \sin \theta = -\frac{1}{2}mr^2\omega^2 \sin 2\theta + mgr \sin \theta \quad (1)$$

We can see that this is 0 for $\theta = 0$, which indeed is an equilibrium. To find if it is a stable equilibrium, we have to check for which ω $W''(\theta = 0) > 0$

$$W''(\theta) = -mr^2\omega^2 \cos 2\theta + mgr \cos \theta \quad (2)$$

$$W''(0) = -mr^2\omega^2 + mgr > 0$$

This gives us the condition that

$$\omega < \sqrt{\frac{g}{r}} = \omega_{cr}$$

We are now going to look at $\omega > \omega_{cr}$. We are going to presuppose that $\sin \theta \neq 0$, and can therefor rewrite $W'(\theta)$ as

$$-mr^2\omega_{cr}\cos\theta = mgr$$

$$\Rightarrow \cos\theta = \frac{g}{r\omega^2}$$

Giving us the new equilibrium points

$$\theta_{\pm} = \arccos\left(\frac{g}{r\omega^2}\right) + n\pi, \quad n = 0, 1$$

d)

When we now are going to check if these points are stable I am going to use a trick which makes it alot easier. I am going to say that

$$\omega = a\omega_{cr}$$

Where $a > 1$. This gives us that the equilibrium points can be written as

$$\theta_{\pm} = \arccos\left(\frac{1}{a^2}\right) + n\pi, \quad n = 0, 1$$

We can now use this to find if $W''(\theta_{\pm}) > 0$. Using 2

$$W''(\arccos(\frac{1}{a^2})) = -mr^2\omega^2 \cos 2(\arccos(\frac{1}{a^2})) + mgr \cos(\arccos(\frac{1}{a^2}))$$

We can obviously see that $\cos(2(\arccos(x))) = 2x^2 - 1$, so

$$-mr^2\omega^2 \left(\frac{2}{a^2} - 1\right) + \frac{mgr}{a^2} > 0$$

$$\Rightarrow \frac{g}{a^2} > ra^2\omega_{cr} \left(\frac{2}{a^2} - 1\right)$$

$$\Rightarrow \frac{1}{a^2} > 2 - a^2$$

$$\Rightarrow 1 > 2a^2 - a^4 = a^2(2 - a^2)$$

Since $\cos(2(\arccos(x)) + n\pi) = 2x^2 - 1$ we get the same inequality for $W''(\arccos(\frac{1}{a^2}) + n\pi)$.

Looking at the inequality $1 > a^2(2 - a^2)$ for $a = 1 \Rightarrow a^2(2 - a^2) = 1$, we defined $a > 1$ and for $a > 1$ the function $a^2(2 - a^2)$ is monotonically decreasing function, thus the inequality holds, and θ_{\pm} are stable equilibriums.

e)

The Hamiltonian is defined here as

$$H = p\dot{\theta} - L \tag{3}$$

where

$$p = \frac{\partial L}{\partial \dot{\theta}} = mr^2\dot{\theta}$$

We then get a Hamiltonian

$$\begin{aligned}
H &= mr^2\dot{\theta}^2 - L = \frac{p}{mr^2} - L \\
&= \frac{p}{mr^2} - \frac{p}{2mr^2} - \frac{1}{2}mr^2\omega^2\sin^2\theta - mgr\cos\theta \\
&= \frac{p}{2mr^2} - \frac{1}{2}mr^2\omega^2\sin^2\theta - mgr\cos\theta
\end{aligned}$$