

FYS3121 Oblig 3

Daniel Heinesen, daniehei

8. februar 2017

1)

a)

To find the Lagrangian, we need to find the Cartesian coordinates expressed with the general coordinate s

$$y = h - s \sin 30^\circ = h - \frac{s}{2}, \quad x = s \cos 30^\circ = s \frac{\sqrt{3}}{2} \quad (1)$$

We then find that

$$\dot{y} = -\frac{\dot{s}}{2}, \quad \dot{x} = \dot{s} \frac{\sqrt{3}}{2}$$

We can then write the energies

$$V = mgy = mg(h - \frac{s}{2})$$
$$T = \frac{1}{2}mv^2 = \frac{1}{2}m(\frac{\dot{s}^2}{4} + \frac{3}{4}\dot{s}^2) = \frac{1}{2}m\dot{s}^2$$

We then find that the Lagrangian is

$$\mathcal{L} = T - V = \frac{1}{2}m\dot{s}^2 - mg(h - \frac{s}{2})$$

b)

With an accelerated incline, we have to rewrite the coordinates

$$y = h - \frac{s}{2}, \quad x = \frac{1}{2}at^2 + s \frac{\sqrt{3}}{2}$$
$$\dot{y} = -\frac{\dot{s}}{2}, \quad \dot{x} = at + \dot{s} \frac{\sqrt{3}}{2}$$

For the energies we see that

$$V = mgy = mg(h - \frac{s}{2})$$
$$T = \frac{1}{2}mv^2 = \frac{1}{2}m(\frac{\dot{s}^2}{4} + a^2t^2 + \dot{s}at\sqrt{3} + \dot{s}^2\frac{3}{4})$$
$$= \frac{1}{2}m(\dot{s}^2 + at(at + \dot{s}\sqrt{3}))$$

So we get the Lagrangian

$$\mathcal{L} = T - V = \frac{1}{2}m(\dot{s}^2 + at(at + \dot{s}\sqrt{3})) - mg(h - \frac{s}{2})$$

c)

We can now find the e.o.m. for this system by solving Lagrange's equation. We first find that

$$\begin{aligned}\frac{\partial \mathcal{L}}{\partial s} &= \frac{mg}{2} \\ \frac{\partial \mathcal{L}}{\partial \dot{s}} &= m\dot{s} + ma\frac{\sqrt{3}}{2} \\ \frac{d}{dt} \left(\frac{\partial \mathcal{L}}{\partial \dot{s}} \right) &= m\ddot{s} + ma\frac{\sqrt{3}}{2}\end{aligned}$$

We can then write the e.o.m

$$\begin{aligned}m\ddot{s} + ma\frac{\sqrt{3}}{2} - \frac{mg}{2} &= 0 \\ \Rightarrow \ddot{s} &= \frac{1}{2}(g - \sqrt{3}a)\end{aligned}$$

We can now integrate this twice with respect to s .

$$s(t) = s_0 + v_0 t + \frac{1}{4}(g - \sqrt{3}a)t^2$$

Knowing that at $t = 0$: $r_0 = 0$ and $v_0 = 0$ we finally get that

$$s(t) = \frac{1}{4}(g - \sqrt{3}a)t^2$$

2)

First we write the coordinates of the ball

$$x = r \sin \theta \cos(\omega t), \quad y = r \sin \theta \sin(\omega t), \quad z = r \cos \theta$$

$$\dot{x} = r\dot{\theta} \cos \theta \cos(\omega t) - r\omega \sin \theta \sin(\omega t)$$

$$\dot{y} = r\dot{\theta} \cos \theta \sin(\omega t) + r\omega \sin \theta \cos(\omega t)$$

$$\dot{z} = -r\dot{\theta} \sin \theta$$

with out writing down the quite long expressions for \dot{x} , \dot{y} and \dot{z} , we get that

$$\dot{x}^2 + \dot{y}^2 + \dot{z}^2 = r^2\dot{\theta}^2 + r^2\omega^2 \sin^2 \theta$$

Since there are no gravitation, we get that $V = 0$, and therefor

$$\mathcal{L} = T = \frac{1}{2}m(r^2\dot{\theta}^2 + r^2\omega^2 \sin^2 \theta)$$

We can now find Lagrange's equation

$$\begin{aligned}\frac{\partial \mathcal{L}}{\partial \theta} &= mr^2\omega^2 \sin \theta \cos \theta \\ \frac{\partial \mathcal{L}}{\partial \dot{\theta}} &= mr^2\dot{\theta}\end{aligned}$$

$$\frac{d}{dt} \left(\frac{\partial \mathcal{L}}{\partial \dot{\theta}} \right) = mr^2 \ddot{\theta}$$

An we find the e.o.m

$$mr^2 \ddot{\theta} - mr^2 \omega^2 \sin \theta \cos \theta = 0$$

b)

c)

3)

a)

Here we start by finding the energies. If we define the plate as being where $V = 0$, we can see that only the lower mass has a potential energy. If the rope has length L , we see that

$$V = mg(-(L - r)) = mg(r - L)$$

The lower mass have a kinetic energy gives as

$$\frac{1}{2} m \dot{r}^2$$

While the upper mass has kinetic energy from being dragged towards the hole, and from the rotation around the hole. Its kinetic energy is then

$$\frac{1}{2} m \dot{r}^2 + \frac{1}{2} m r^2 \dot{\theta}^2$$

Giving us a total kinetic energy of

$$T = 2 \cdot \frac{1}{2} m \dot{r}^2 + \frac{1}{2} m r^2 \dot{\theta}^2$$

And a Lagrangian

$$\mathcal{L} = \frac{1}{2} m (2\dot{r}^2 + r^2 \dot{\theta}^2) - mg(r - L)$$

We can now find the e.o.m

$$\frac{\partial \mathcal{L}}{\partial \theta} = 0 \tag{2}$$

$$\frac{\partial \mathcal{L}}{\partial \dot{\theta}} = mr^2 \dot{\theta} \tag{3}$$

$$\frac{d}{dt} \left(\frac{\partial \mathcal{L}}{\partial \dot{\theta}} \right) = mr^2 \ddot{\theta} + 2mr\dot{r}\dot{\theta}$$

So the first e.o.m is:

$$mr^2 \ddot{\theta} + 2mr\dot{r}\dot{\theta} = 0$$

For the second

$$\frac{\partial \mathcal{L}}{\partial r} = mr\dot{\theta}^2 - mg$$

$$\frac{\partial \mathcal{L}}{\partial \dot{r}} = 2m\dot{r}$$

$$\frac{d}{dt} \left(\frac{\partial \mathcal{L}}{\partial \dot{r}} \right) = 2m\ddot{r}$$

So the second e.o.m becomes

$$2m\ddot{r} - mr\dot{\theta}^2 + mg = 0$$

b)

As we saw above from 2

$$\frac{\partial \mathcal{L}}{\partial \theta} = 0$$

This means that θ is a cyclic coordinate, and therefor

$$\frac{\partial \mathcal{L}}{\partial \dot{\theta}} = mr^2\dot{\theta} = \text{constant} = \ell$$

We can there for rewrite $\dot{\theta}$ as

$$\dot{\theta} = \frac{\ell}{mr^2}$$

If we set this into the second e.o.m we get a description of the system, only dependent on r

$$2m\ddot{r} - \frac{\ell^2}{mr^3} + mg = 0$$

4)

a)

For convenience I am going to label the upper mass m_1 and the pendulum bob m_2 . The coordinates then becomes

$$x_1 = s, \quad y_1 = 0$$

$$\dot{x}_1 = \dot{s}, \quad \dot{y}_1 = 0$$

and

$$x_2 = s + d \sin \theta, \quad y_2 = -d \cos \theta$$

$$\dot{x}_2 = \dot{s} + d \cos \theta, \quad \dot{y}_2 = d \sin \theta$$

And the energies

$$V = mg(y_1 + y_2) = -mgd \cos \theta$$

$$T = \frac{1}{2}m(\dot{x}_1^2 + \dot{y}_1^2 + \dot{x}_2^2 + \dot{y}_2^2)$$

$$= \frac{1}{2}m(2\dot{s}^2 + d^2\dot{\theta}^2 + 2\dot{\theta}\dot{s}d \cos \theta)$$

This gives us

$$\mathcal{L} = T - V = \frac{1}{2}m(2\dot{s}^2 + d^2\dot{\theta}^2 + 2\dot{\theta}\dot{s}d \cos \theta) + mgd \cos \theta$$

b)

If we differentiate \mathcal{L} with respect to s

$$\frac{\partial \mathcal{L}}{\partial s} = 0$$

So s is a cyclic coordinate, meaning that the Lagrangian does not depend on s . It can also be shown that since s is cyclic, there is a conserved quantity in this system, namely

$$\frac{\partial \mathcal{L}}{\partial \dot{s}} = 2m\dot{s} + md\dot{\theta} \cos \theta$$

Since this quantity is conserved, it is constant

$$2m\dot{s} + md\dot{\theta} \cos \theta = \ell$$

We can therefore rewrite this as

$$\dot{s} = \frac{1}{2} \left(\frac{\ell}{m} - d\dot{\theta} \cos \theta \right)$$

If we insert this into the Lagrangian, we get

$$\begin{aligned} \mathcal{L} &= \frac{1}{2}m \left(\frac{1}{2} \left(\frac{\ell}{m} - d\dot{\theta} \cos \theta \right)^2 + d^2\dot{\theta}^2 + \dot{\theta} \left(\frac{\ell}{m} - d\dot{\theta} \cos \theta \right) d \cos \theta \right) + mgd \cos \theta \\ &= \frac{1}{2}m \left(\frac{\ell^2}{2m^2} + \frac{d^2\dot{\theta}^2}{2} \cos^2 \theta + d^2\dot{\theta} \right) + mgd \sin \theta \end{aligned}$$

c)

We can now find the e.o.m

$$\begin{aligned} \frac{\partial \mathcal{L}}{\partial \theta} &= -mgd \sin \theta + \frac{1}{2}md^2\dot{\theta}^2 \cos \theta \sin \theta \\ \frac{\partial \mathcal{L}}{\partial \dot{\theta}} &= -\frac{1}{2}md^2\dot{\theta} \cos^2 \theta + md^2\dot{\theta} = md^2 \left(\dot{\theta} - \frac{1}{2}\dot{\theta} \cos^2 \theta \right) \\ \frac{d}{dt} \left(\frac{\partial \mathcal{L}}{\partial \dot{\theta}} \right) &= md^2 \left(\ddot{\theta} - \frac{1}{2}\ddot{\theta} \cos^2 \theta + \dot{\theta}^2 \sin \theta \cos \theta \right) \end{aligned}$$

This gives us that

$$\begin{aligned} md^2 \left(\ddot{\theta} - \frac{1}{2}\ddot{\theta} \cos^2 \theta + \dot{\theta}^2 \sin \theta \cos \theta \right) + mgd \sin \theta - \frac{1}{2}md^2\dot{\theta}^2 \cos \theta \sin \theta &= 0 \\ \Rightarrow \ddot{\theta} \left(1 - \frac{1}{2} \cos^2 \theta \right) + \frac{\dot{\theta}^2}{2} \sin \theta \cos \theta + \frac{g}{d} \sin \theta &= 0 \end{aligned}$$

d)

We can find a small-angle solution to this problem. If we Taylor expand the above expression around $\theta = 0$. We get that the first order Taylor series is

$$T_2(f(\theta)) = \frac{1}{2}\ddot{\theta} + \frac{g}{d}\theta + \frac{1}{2}\theta\dot{\theta}^2 = 0$$

We are only interested in the first order, and for small θ see get that $\dot{\theta}^2 \rightarrow 0$, so this expression becomes

$$\frac{1}{2}\ddot{\theta} + \frac{g}{d}\theta = 0$$

This is a harmonic oscillator equation

$$\ddot{\theta} = -\frac{2g}{d}\theta$$

The angular frequency of this system will be

$$\omega = \sqrt{\frac{2g}{d}}$$