

# Total positivity of the Eulerian triangle: A big generalisation of Brenti's conjecture

Bishal Deb (he/him)

University College London

June 8, 2021

Young Researchers in Mathematics (YRM) 2021

University of Bristol

*Based on Ongoing Joint Work With*

Xi Chen, Alexander Dyachenko, Tomack Gilmore and Alan D. Sokal

## Definition (Total Positivity (TP))

A matrix of real numbers said to be **totally positive (TP)** if all its minors are non-negative

## Definition (Total Positivity (TP))

A matrix of real numbers said to be **totally positive (TP)** if all its minors are non-negative i.e., determinants of all finite square submatrices are non-negative.

## Definition (Total Positivity (TP))

A matrix of real numbers said to be **totally positive (TP)** if all its minors are non-negative i.e., determinants of all finite square submatrices are non-negative.

Array of numbers and not linear operator.

## Definition (Total Positivity (TP))

A matrix of real numbers said to be **totally positive (TP)** if all its minors are non-negative i.e., determinants of all finite square submatrices are non-negative.

Array of numbers and not linear operator.

Need not be a square matrix,

## Definition (Total Positivity (TP))

A matrix of real numbers said to be **totally positive (TP)** if all its minors are non-negative i.e., determinants of all finite square submatrices are non-negative.

Array of numbers and not linear operator.

Need not be a square matrix, or finite!

## Definition (Total Positivity (TP))

A matrix of real numbers said to be **totally positive (TP)** if all its minors are non-negative i.e., determinants of all finite square submatrices are non-negative.

Array of numbers and not linear operator.

Need not be a square matrix, or finite!


Will consider matrix of polynomials soon!

# Historical Note

First defined independently by two different groups in the 30s



(a) M.G. Krein (1907-1989)



(b) I.G. Schoenberg  
(1903-1990)

Source: MacTutor History of Mathematics Archive

We use Schoenberg's terminology.



# Example and basic properties

# Example and basic properties

- Example: Bidiagonal matrices with entries  $\geq 0$

$$\begin{bmatrix} 0 & a_1 & b_1 & 0 & 0 & 0 & 0 \\ 0 & 0 & a_2 & b_2 & 0 & 0 & 0 \\ 0 & 0 & 0 & a_3 & b_3 & 0 & 0 \\ 0 & 0 & 0 & 0 & a_4 & b_4 & 0 \\ 0 & 0 & 0 & 0 & 0 & a_5 & b_5 \\ 0 & 0 & 0 & 0 & 0 & 0 & a_6 \end{bmatrix}$$

# Example and basic properties

- Example: Bidiagonal matrices with entries  $\geq 0$

$$\begin{bmatrix} 0 & a_1 & b_1 & 0 & 0 & 0 & 0 \\ 0 & 0 & a_2 & b_2 & 0 & 0 & 0 \\ 0 & 0 & 0 & a_3 & b_3 & 0 & 0 \\ 0 & 0 & 0 & 0 & a_4 & b_4 & 0 \\ 0 & 0 & 0 & 0 & 0 & a_5 & b_5 \\ 0 & 0 & 0 & 0 & 0 & 0 & a_6 \end{bmatrix}$$

- Matrix addition doesn't preserve TP.

# Example and basic properties

- Example: Bidiagonal matrices with entries  $\geq 0$

$$\begin{bmatrix} 0 & a_1 & b_1 & 0 & 0 & 0 & 0 \\ 0 & 0 & a_2 & b_2 & 0 & 0 & 0 \\ 0 & 0 & 0 & a_3 & b_3 & 0 & 0 \\ 0 & 0 & 0 & 0 & a_4 & b_4 & 0 \\ 0 & 0 & 0 & 0 & 0 & a_5 & b_5 \\ 0 & 0 & 0 & 0 & 0 & 0 & a_6 \end{bmatrix}$$

- Matrix addition doesn't preserve TP. Example:

$$\begin{bmatrix} 0 & 0 \\ 1 & 1 \end{bmatrix} + \begin{bmatrix} 1 & 2 \\ 0 & 0 \end{bmatrix} = \begin{bmatrix} 1 & 2 \\ 1 & 1 \end{bmatrix}$$

# Example and basic properties

- Example: Bidiagonal matrices with entries  $\geq 0$

$$\begin{bmatrix} 0 & a_1 & b_1 & 0 & 0 & 0 & 0 \\ 0 & 0 & a_2 & b_2 & 0 & 0 & 0 \\ 0 & 0 & 0 & a_3 & b_3 & 0 & 0 \\ 0 & 0 & 0 & 0 & a_4 & b_4 & 0 \\ 0 & 0 & 0 & 0 & 0 & a_5 & b_5 \\ 0 & 0 & 0 & 0 & 0 & 0 & a_6 \end{bmatrix}$$

- Matrix addition doesn't preserve TP. Example:

$$\begin{bmatrix} 0 & 0 \\ 1 & 1 \end{bmatrix} + \begin{bmatrix} 1 & 2 \\ 0 & 0 \end{bmatrix} = \begin{bmatrix} 1 & 2 \\ 1 & 1 \end{bmatrix}$$

- Matrix multiplication preserves TP

# Example and basic properties

- Example: Bidiagonal matrices with entries  $\geq 0$

$$\begin{bmatrix} 0 & a_1 & b_1 & 0 & 0 & 0 & 0 \\ 0 & 0 & a_2 & b_2 & 0 & 0 & 0 \\ 0 & 0 & 0 & a_3 & b_3 & 0 & 0 \\ 0 & 0 & 0 & 0 & a_4 & b_4 & 0 \\ 0 & 0 & 0 & 0 & 0 & a_5 & b_5 \\ 0 & 0 & 0 & 0 & 0 & 0 & a_6 \end{bmatrix}$$

- Matrix addition doesn't preserve TP. Example:

$$\begin{bmatrix} 0 & 0 \\ 1 & 1 \end{bmatrix} + \begin{bmatrix} 1 & 2 \\ 0 & 0 \end{bmatrix} = \begin{bmatrix} 1 & 2 \\ 1 & 1 \end{bmatrix}$$

- Matrix multiplication preserves TP (Cauchy-Binet formula)

# Applications to Combinatorics

- Brenti in 80s started studying TP in combinatorics.



- Brenti in 80s started studying TP in combinatorics.
- Many important combinatorial triangles are TP.

# Example 1: Binomial triangle

Entries  $\binom{n}{k}$

$$\begin{bmatrix} 1 & & & & & \\ 1 & 1 & & & & \\ 1 & 2 & 1 & & & \\ 1 & 3 & 3 & 1 & & \\ 1 & 4 & 6 & 4 & 1 & \\ \vdots & \vdots & \vdots & \vdots & \vdots & \ddots \end{bmatrix}$$

## Example 2: Stirling subset triangle

Entries  $\left\{ \begin{smallmatrix} n+1 \\ k \end{smallmatrix} \right\}$ : set partitions of  $\{1, \dots, n+1\}$  with  $k$  blocks

$$\begin{bmatrix} 1 & & & & & \\ 1 & 1 & & & & \\ 1 & 3 & 1 & & & \\ 1 & 7 & 6 & 1 & & \\ 1 & 15 & 25 & 10 & 1 & \\ \vdots & \vdots & \vdots & \vdots & \vdots & \ddots \end{bmatrix}$$

## Example 3: Stirling cycle triangle

Entries  $\left[ \begin{smallmatrix} n+1 \\ k \end{smallmatrix} \right]$ : permutations on  $n + 1$  letters with  $k$  cycles

$$\begin{bmatrix} 1 & & & & & \\ 1 & 1 & & & & \\ 2 & 3 & 1 & & & \\ 6 & 11 & 6 & 1 & & \\ 24 & 50 & 35 & 10 & 1 & \\ \vdots & \vdots & \vdots & \vdots & \vdots & \ddots \end{bmatrix}$$

## Example 4: Reversed Stirling cycle triangle

Original Matrix

$$\begin{bmatrix} 1 & & & & & \\ 1 & 1 & & & & \\ 2 & 3 & 1 & & & \\ 6 & 11 & 6 & 1 & & \\ 24 & 50 & 35 & 10 & 1 & \\ \vdots & \vdots & \vdots & \vdots & \vdots & \ddots \end{bmatrix}$$

Reversal

$$\begin{bmatrix} 1 & & & & & \\ 1 & 1 & & & & \\ 1 & 3 & 2 & & & \\ 1 & 6 & 11 & 6 & & \\ 1 & 10 & 35 & 50 & 24 & \\ \vdots & \vdots & \vdots & \vdots & \vdots & \ddots \end{bmatrix}$$

# Reversal of a lower triangular matrix

Matrix	Total positivity of Reversal
--------	------------------------------

# Reversal of a lower triangular matrix

Matrix	Total positivity of Reversal
Binomial triangle	True

# Reversal of a lower triangular matrix

Matrix	Total positivity of Reversal
Binomial triangle	True
Stirling cycle triangle	True (Brenti, 1995)



# Reversal of a lower triangular matrix

Matrix	Total positivity of Reversal
Binomial triangle	True
Stirling cycle triangle	True (Brenti, 1995)
Stirling subset triangle	Empirically True

# Reversal of a lower triangular matrix

Matrix	Total positivity of Reversal
Binomial triangle	True
Stirling cycle triangle	True (Brenti, 1995)
Stirling subset triangle	Empirically True
Arbitrary lower triangular TP matrix	Not true

# Reversed Stirling Subset Triangle

Reversed Stirling subset triangle:

$$\left( \left\{ \begin{matrix} n+1 \\ k \end{matrix} \right\}^{\text{rev}} \right)_{n,k \geq 0} = \left( \left\{ \begin{matrix} n+1 \\ n-k+1 \end{matrix} \right\} \right)_{n,k \geq 0} = \begin{bmatrix} 1 & & & & & \\ 1 & 1 & & & & \\ 1 & 3 & 1 & & & \\ 1 & 6 & 7 & 1 & & \\ 1 & 10 & 25 & 15 & 1 & \\ \vdots & \vdots & \vdots & \vdots & \vdots & \ddots \end{bmatrix}$$

# Reversed Stirling Subset Triangle

Reversed Stirling subset triangle:

$$\left( \left\{ \begin{matrix} n+1 \\ k \end{matrix} \right\}^{\text{rev}} \right)_{n,k \geq 0} = \left( \left\{ \begin{matrix} n+1 \\ n-k+1 \end{matrix} \right\} \right)_{n,k \geq 0} = \begin{bmatrix} 1 & & & & & \\ 1 & 1 & & & & \\ 1 & 3 & 1 & & & \\ 1 & 6 & 7 & 1 & & \\ 1 & 10 & 25 & 15 & 1 & \\ \vdots & \vdots & \vdots & \vdots & \vdots & \ddots \end{bmatrix}$$

Conjecture (Us 2019)

*The infinite lower triangular matrix  $\left( \left\{ \begin{matrix} n+1 \\ k \end{matrix} \right\}^{\text{rev}} \right)_{n,k \geq 0} = \left( \left\{ \begin{matrix} n+1 \\ n-k+1 \end{matrix} \right\} \right)_{n,k \geq 0}$  is TP.*

# Eulerian Numbers

Descent of a permutation  $\sigma$ :

Descent of a permutation  $\sigma$ : Position  $i$  such that  $\sigma(i) > \sigma(i+1)$ .

# Eulerian Numbers

Descent of a permutation  $\sigma$ : Position  $i$  such that  $\sigma(i) > \sigma(i+1)$ .

$\text{des}(\sigma)$ : number of descents of  $\sigma$ .



# Eulerian Numbers

Descent of a permutation  $\sigma$ : Position  $i$  such that  $\sigma(i) > \sigma(i+1)$ .

$\text{des}(\sigma)$ : number of descents of  $\sigma$ .

Example: For  $\sigma = 943127685$ ,  $\text{des}(\sigma) = 5$

Descent of a permutation  $\sigma$ : Position  $i$  such that  $\sigma(i) > \sigma(i+1)$ .

$\text{des}(\sigma)$ : number of descents of  $\sigma$ .

Example: For  $\sigma = 943127685$ ,  $\text{des}(\sigma) = 5$

## Definition

The Eulerian number  $\left\langle n \atop k \right\rangle$  is defined to be the cardinality of the set  $\{\sigma \in \mathfrak{S}_n \mid \text{des}(\sigma) = k\}$ .

## Definition

The Eulerian number  $\left\langle n \atop k \right\rangle$  is defined to be the cardinality of the set  $\{\sigma \in \mathfrak{S}_n \mid \text{des}(\sigma) = k\}$ .

## Definition

The Eulerian number  $\langle n \rangle_k$  is defined to be the cardinality of the set  $\{\sigma \in \mathfrak{S}_n \mid \text{des}(\sigma) = k\}$ .

$$\left( \left\langle \begin{matrix} n+1 \\ k \end{matrix} \right\rangle \right)_{n,k \geq 0} = \begin{bmatrix} 1 & & & & & & \\ 1 & 1 & & & & & \\ 1 & 4 & 1 & & & & \\ 1 & 11 & 11 & 1 & & & \\ 1 & 26 & 66 & 26 & 1 & & \\ 1 & 57 & 302 & 302 & 57 & 1 & \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \ddots \end{bmatrix}$$

# Eulerian Triangle Conjecture

$$\left( \left\langle \begin{matrix} n+1 \\ k \end{matrix} \right\rangle \right)_{n,k \geq 0} = \begin{bmatrix} 1 & & & & & & \\ 1 & 1 & & & & & \\ 1 & 4 & 1 & & & & \\ 1 & 11 & 11 & 1 & & & \\ 1 & 26 & 66 & 26 & 1 & & \\ 1 & 57 & 302 & 302 & 57 & 1 & \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \ddots \end{bmatrix}$$

Conjecture (Brenti 1996)

*The infinite lower triangular matrix  $\left( \left\langle \begin{matrix} n+1 \\ k \end{matrix} \right\rangle \right)_{n,k \geq 0}$  is TP.*

# Eulerian Triangle Conjecture

$$\left( \left\langle \begin{matrix} n+1 \\ k \end{matrix} \right\rangle \right)_{n,k \geq 0} = \begin{bmatrix} 1 & & & & & & \\ 1 & 1 & & & & & \\ 1 & 4 & 1 & & & & \\ 1 & 11 & 11 & 1 & & & \\ 1 & 26 & 66 & 26 & 1 & & \\ 1 & 57 & 302 & 302 & 57 & 1 & \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \ddots \end{bmatrix}$$

Conjecture (Brenti 1996)

*The infinite lower triangular matrix  $\left( \left\langle \begin{matrix} n+1 \\ k \end{matrix} \right\rangle \right)_{n,k \geq 0}$  is TP.*

Verified for  $512 \times 512$  using an efficient bidiagonal factorisation algorithm.

# Eulerian Triangle Conjecture

$$\left( \left\langle \begin{matrix} n+1 \\ k \end{matrix} \right\rangle \right)_{n,k \geq 0} = \begin{bmatrix} 1 & & & & & & \\ 1 & 1 & & & & & \\ 1 & 4 & 1 & & & & \\ 1 & 11 & 11 & 1 & & & \\ 1 & 26 & 66 & 26 & 1 & & \\ 1 & 57 & 302 & 302 & 57 & 1 & \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \ddots \end{bmatrix}$$

Conjecture (Brenti 1996)

*The infinite lower triangular matrix  $\left( \left\langle \begin{matrix} n+1 \\ k \end{matrix} \right\rangle \right)_{n,k \geq 0}$  is TP.*

Verified for  $512 \times 512$  using an efficient bidiagonal factorisation algorithm.

Rational numbers with large denominators appear.

# Eulerian Triangle Conjecture

$$\left( \left\langle \begin{matrix} n+1 \\ k \end{matrix} \right\rangle \right)_{n,k \geq 0} = \begin{bmatrix} 1 & & & & & & \\ 1 & 1 & & & & & \\ 1 & 4 & 1 & & & & \\ 1 & 11 & 11 & 1 & & & \\ 1 & 26 & 66 & 26 & 1 & & \\ 1 & 57 & 302 & 302 & 57 & 1 & \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \ddots \end{bmatrix}$$

Conjecture (Brenti 1996)

*The infinite lower triangular matrix  $\left( \left\langle \begin{matrix} n+1 \\ k \end{matrix} \right\rangle \right)_{n,k \geq 0}$  is TP.*

Verified for  $512 \times 512$  using an efficient bidiagonal factorisation algorithm.

Rational numbers with large denominators appear.

General pattern not clear.



# A comparison of the two triangles

Reversed Stirling subset triangle:

$$\left( \left\{ \begin{matrix} n+1 \\ k \end{matrix} \right\}^{\text{rev}} \right)_{n,k \geq 0} = \begin{bmatrix} 1 & & & & & \\ 1 & 1 & & & & \\ 1 & 3 & 1 & & & \\ 1 & 6 & 7 & 1 & & \\ 1 & 10 & 25 & 15 & 1 & \\ \vdots & \vdots & \vdots & \vdots & \vdots & \ddots \end{bmatrix}$$

Eulerian triangle:

$$\left( \left\langle \begin{matrix} n+1 \\ k \end{matrix} \right\rangle \right)_{n,k \geq 0} = \begin{bmatrix} 1 & & & & & \\ 1 & 1 & & & & \\ 1 & 4 & 1 & & & \\ 1 & 11 & 11 & 1 & & \\ 1 & 26 & 66 & 26 & 1 & \\ \vdots & \vdots & \vdots & \vdots & \vdots & \ddots \end{bmatrix}$$

# A comparison of the two triangles

Reversed Stirling subset triangle:

$$\left( \left\{ \begin{matrix} n+1 \\ k \end{matrix} \right\}^{\text{rev}} \right)_{n,k \geq 0} = \begin{bmatrix} 1 & & & & & \\ 1 & 1 & & & & \\ 1 & 3 & 1 & & & \\ 1 & 6 & 7 & 1 & & \\ 1 & 10 & 25 & 15 & 1 & \\ \vdots & \vdots & \vdots & \vdots & \vdots & \ddots \end{bmatrix}$$

$$A_{n,k} = (n - k + 1)A_{n-1,k-1} + 1 \cdot A_{n-1,k}$$

Eulerian triangle:

$$\left( \left\langle \begin{matrix} n+1 \\ k \end{matrix} \right\rangle \right)_{n,k \geq 0} = \begin{bmatrix} 1 & & & & & \\ 1 & 1 & & & & \\ 1 & 4 & 1 & & & \\ 1 & 11 & 11 & 1 & & \\ 1 & 26 & 66 & 26 & 1 & \\ \vdots & \vdots & \vdots & \vdots & \vdots & \ddots \end{bmatrix}$$

$$A_{n,k} = (n - k + 1)A_{n-1,k-1} + (k + 1)A_{n-1,k}$$

# A comparison of the two triangles

Reversed Stirling subset triangle:

$$\left( \left\{ \begin{matrix} n+1 \\ k \end{matrix} \right\}^{\text{rev}} \right)_{n,k \geq 0} = \begin{bmatrix} 1 & & & & & \\ 1 & 1 & & & & \\ 1 & 3 & 1 & & & \\ 1 & 6 & 7 & 1 & & \\ 1 & 10 & 25 & 15 & 1 & \\ \vdots & \vdots & \vdots & \vdots & \vdots & \ddots \end{bmatrix}$$

$$A_{n,k} = (n - k + 1) A_{n-1,k-1} + 1 \cdot A_{n-1,k}$$

Introduce  
variables

Eulerian triangle:

$$\left( \left\langle \begin{matrix} n+1 \\ k \end{matrix} \right\rangle \right)_{n,k \geq 0} = \begin{bmatrix} 1 & & & & & \\ 1 & 1 & & & & \\ 1 & 4 & 1 & & & \\ 1 & 11 & 11 & 1 & & \\ 1 & 26 & 66 & 26 & 1 & \\ \vdots & \vdots & \vdots & \vdots & \vdots & \ddots \end{bmatrix}$$

$$A_{n,k} = (n - k + 1) A_{n-1,k-1} + (k + 1) A_{n-1,k}$$

## Definition (Coefficientwise TP)

A matrix of polynomials with real coefficients is said to be **coefficientwise totally positive (coefficientwise TP)** if all its minors have non-negative coefficients.

## Definition (Coefficientwise TP)

A matrix of polynomials with real coefficients is said to be **coefficientwise totally positive (coefficientwise TP)** if all its minors have non-negative coefficients.

One or several variables.

## Definition (Coefficientwise TP)

A matrix of polynomials with real coefficients is said to be **coefficientwise totally positive (coefficientwise TP)** if all its minors have non-negative coefficients.

One or several variables.

Coefficientwise TP  $\implies$  Pointwise TP.

## Definition (Coefficientwise TP)

A matrix of polynomials with real coefficients is said to be **coefficientwise totally positive (coefficientwise TP)** if all its minors have non-negative coefficients.

One or several variables.

Coefficientwise TP  $\implies$  Pointwise TP.

But much stronger.

# Putting variables in the recurrence

Replace

$$A_{n,k} = (n - k + 1)A_{n-1,k-1} + (k + 1)A_{n-1,k}$$

$$A_{n,k} = (n - k + 1)A_{n-1,k-1} + 1 \cdot A_{n-1,k}$$

Introduce variables  $a, b, c, d, e$ .

$$A_{n,k} = [a(n - k) + b(k - 1) + c]A_{n-1,k-1} + (dk + e)A_{n-1,k}$$



# Putting variables in the recurrence

Replace

$$A_{n,k} = (n - k + 1)A_{n-1,k-1} + (k + 1)A_{n-1,k}$$

$$A_{n,k} = (n - k + 1)A_{n-1,k-1} + 1 \cdot A_{n-1,k}$$

Introduce variables  $a, b, c, d, e$ .

$$A_{n,k} = [a(n - k) + b(k - 1) + c]A_{n-1,k-1} + (dk + e)A_{n-1,k}$$

Is it coefficientwise TP?

(Minors are polynomials with non-negative coefficients?)

# Putting variables in the recurrence

Replace

$$A_{n,k} = (n - k + 1)A_{n-1,k-1} + (k + 1)A_{n-1,k}$$

$$A_{n,k} = (n - k + 1)A_{n-1,k-1} + 1 \cdot A_{n-1,k}$$

Introduce variables  $a, b, c, d, e$ .

$$A_{n,k} = [a(n - k) + b(k - 1) + c]A_{n-1,k-1} + (dk + e)A_{n-1,k}$$

Is it coefficientwise TP?

(Minors are polynomials with non-negative coefficients?)

More general conjecture!

# Putting variables in the recurrence

Replace

$$A_{n,k} = (n - k + 1)A_{n-1,k-1} + (k + 1)A_{n-1,k}$$

$$A_{n,k} = (n - k + 1)A_{n-1,k-1} + 1 \cdot A_{n-1,k}$$

Introduce variables  $a, b, c, d, e$ .

$$A_{n,k} = [a(n - k) + b(k - 1) + c]A_{n-1,k-1} + (dk + e)A_{n-1,k}$$

Is it coefficientwise TP?

(Minors are polynomials with non-negative coefficients?)

More general conjecture!

True till  $11 \times 11$  - a week.

# Putting variables in the recurrence

Replace

$$A_{n,k} = (n - k + 1)A_{n-1,k-1} + (k + 1)A_{n-1,k}$$

$$A_{n,k} = (n - k + 1)A_{n-1,k-1} + 1 \cdot A_{n-1,k}$$

Introduce variables  $a, b, c, d, e$ .

$$A_{n,k} = [a(n - k) + b(k - 1) + c]A_{n-1,k-1} + (dk + e)A_{n-1,k}$$

Is it coefficientwise TP?

(Minors are polynomials with non-negative coefficients?)

More general conjecture!

True till  $11 \times 11$  - a week.

False for  $12 \times 12$  - 99 days CPU time.

# Putting variables in the recurrence

Replace

$$A_{n,k} = (n - k + 1)A_{n-1,k-1} + (k + 1)A_{n-1,k}$$

$$A_{n,k} = (n - k + 1)A_{n-1,k-1} + 1 \cdot A_{n-1,k}$$

Introduce variables  $a, b, c, d, e$ .

$$A_{n,k} = [a(n - k) + b(k - 1) + c]A_{n-1,k-1} + (dk + e)A_{n-1,k}$$

Is it coefficientwise TP?

(Minors are polynomials with non-negative coefficients?)

More general conjecture!

True till  $11 \times 11$  - a week.

False for  $12 \times 12$  - 99 days CPU time. ☹

# Is all hope lost?

Put  $b = 0$  to obtain

$$A_{n,k} = [a(n-k) + c]A_{n-1,k-1} + (dk + e)A_{n-1,k}$$

# Is all hope lost?

Put  $b = 0$  to obtain

$$A_{n,k} = [a(n-k) + c]A_{n-1,k-1} + (dk + e)A_{n-1,k}$$

Empirically true upto  $13 \times 13$ .

# Is all hope lost?

Put  $b = 0$  to obtain

$$A_{n,k} = [a(n-k) + c]A_{n-1,k-1} + (dk + e)A_{n-1,k}$$

Empirically true upto  $13 \times 13$ .

Special cases:

$(a, c, d, e)$	Matrix obtained
----------------	-----------------



# Is all hope lost?

Put  $b = 0$  to obtain

$$A_{n,k} = [a(n-k) + c]A_{n-1,k-1} + (dk + e)A_{n-1,k}$$

Empirically true upto  $13 \times 13$ .

Special cases:

$(a, c, d, e)$	Matrix obtained
$(1, 1, 1, 1)$	clean Eulerian triangle, conjecture
$(1, 0, 1, 1)$	shifted Eulerian triangle, conjecture
$(1, 1, 1, 0)$	shifted Eulerian triangle, conjecture

# Is all hope lost?

Put  $b = 0$  to obtain

$$A_{n,k} = [a(n-k) + c]A_{n-1,k-1} + (dk + e)A_{n-1,k}$$

Empirically true upto  $13 \times 13$ .

Special cases:

$(a, c, d, e)$	Matrix obtained
$(1, 1, 1, 1)$	clean Eulerian triangle, conjecture
$(1, 0, 1, 1)$	shifted Eulerian triangle, conjecture
$(1, 1, 1, 0)$	shifted Eulerian triangle, conjecture
$(0, 1, 0, 1)$	Binomial triangle, TP

# Is all hope lost?

Put  $b = 0$  to obtain

$$A_{n,k} = [a(n-k) + c]A_{n-1,k-1} + (dk + e)A_{n-1,k}$$

Empirically true upto  $13 \times 13$ .

Special cases:

$(a, c, d, e)$	Matrix obtained
$(1, 1, 1, 1)$	clean Eulerian triangle, conjecture
$(1, 0, 1, 1)$	shifted Eulerian triangle, conjecture
$(1, 1, 1, 0)$	shifted Eulerian triangle, conjecture
$(0, 1, 0, 1)$	Binomial triangle, TP
$(0, 1, 1, 1)$	Stirling subset numbers, TP proved by Brenti 1995

# Is all hope lost?


Put  $b = 0$  to obtain

$$A_{n,k} = [a(n-k) + c]A_{n-1,k-1} + (dk + e)A_{n-1,k}$$

Empirically true upto  $13 \times 13$ .

Special cases:

$(a, c, d, e)$	Matrix obtained
$(1, 1, 1, 1)$	clean Eulerian triangle, conjecture
$(1, 0, 1, 1)$	shifted Eulerian triangle, conjecture
$(1, 1, 1, 0)$	shifted Eulerian triangle, conjecture
$(0, 1, 0, 1)$	Binomial triangle, TP
$(0, 1, 1, 1)$	Stirling subset numbers, TP proved by Brenti 1995
$(1, 1, 0, 1)$	Row reversed matrix of Stirling subset numbers, conjecture



$$A_{n,k} = (n - k + 1)A_{n-1,k-1}$$

$$+ (k + 1)A_{n-1,k}$$

$$A_{n,k} = (n - k + 1)A_{n-1,k-1} + 1 \cdot A_{n-1,k}$$



$$A_{n,k} = [a(n - k) + c]A_{n-1,k-1}$$

$$+ (dk + e)A_{n-1,k}$$

# Is all hope lost?

Put  $b = 0$  to obtain

$$A_{n,k} = [a(n-k) + c]A_{n-1,k-1} + (dk + e)A_{n-1,k}$$

This seems to be coefficientwise TP so far.

In fact, we have the following special cases for different substitutions:

$(a, c, d, e)$	Matrix obtained
$(1, 1, 1, 1)$	clean Eulerian triangle, conjecture
$(1, 0, 1, 1)$	Down-shifted Eulerian triangle, conjecture
$(1, 1, 1, 0)$	Diagonally down-shifted Eulerian triangle, conjecture
$(0, 1, 0, 1)$	Binomial triangle, TP
$(0, 1, 1, 1)$	Stirling subset numbers, TP proved by Brenti 1995
$(1, 1, 0, 1)$	<b>Row reversed matrix of Stirling subset numbers, conjecture</b>

# Is all hope lost?

Put  $b = 0$  to obtain

$$A_{n,k} = [a(n-k) + c]A_{n-1,k-1} + (dk + e)A_{n-1,k}$$

This seems to be coefficientwise TP so far.

In fact, we have the following special cases for different substitutions:

$(a, c, d, e)$	Matrix obtained
$(1, 1, 1, 1)$	clean Eulerian triangle, conjecture
$(1, 0, 1, 1)$	Down-shifted Eulerian triangle, conjecture
$(1, 1, 1, 0)$	Diagonally down-shifted Eulerian triangle, conjecture
$(0, 1, 0, 1)$	Binomial triangle, TP
$(0, 1, 1, 1)$	Stirling subset numbers, TP proved by Brenti 1995
$(1, 1, 0, 1)$	<b>Row reversed matrix of Stirling subset numbers, conjecture</b>

## Reversed Stirling subset triangle

$$\left( \left\{ \begin{matrix} n+1 \\ k \end{matrix} \right\}^{\text{rev}} \right)_{n,k \geq 0} = \begin{bmatrix} 1 & & & & & \\ 1 & 1 & & & & \\ 1 & 3 & 1 & & & \\ 1 & 6 & 7 & 1 & & \\ 1 & 10 & 25 & 15 & 1 & \\ \vdots & \vdots & \vdots & \vdots & \vdots & \ddots \end{bmatrix}.$$



ace triangle

$$\begin{bmatrix} 1 & & & & & \\ e & c & & & & \\ e^2 & ae + 2ce & c^2 & & & \\ e^3 & 3ae^2 + 3ce^2 & a^2e + 3ace + 3c^2e & c^3 & & \\ e^4 & 6ae^3 + 4ce^3 & 7a^2e^2 + 12ace^2 + 6c^2e^2 & a^3e + 4a^2ce + 6ac^2e + 4c^3e & c^4 & \\ \vdots & \vdots & \vdots & \vdots & \vdots & \ddots \end{bmatrix}.$$

recurrence

$$A_{n,k} = [\textcolor{green}{a}(n-k) + \textcolor{red}{c}]A_{n-1,k-1} + \textcolor{blue}{e}A_{n-1,k}$$

ace triangle

$$\begin{bmatrix} 1 & & & & & \\ e & c & & & & \\ e^2 & ae + 2ce & c^2 & & & \\ e^3 & 3ae^2 + 3ce^2 & a^2e + 3ace + 3c^2e & c^3 & & \\ e^4 & 6ae^3 + 4ce^3 & 7a^2e^2 + 12ace^2 + 6c^2e^2 & a^3e + 4a^2ce + 6ac^2e + 4c^3e & c^4 & \\ \vdots & \vdots & \vdots & \vdots & \vdots & \ddots \end{bmatrix}.$$


recurrence

$$A_{n,k} = [\textcolor{green}{a}(n-k) + \textcolor{red}{c}]A_{n-1,k-1} + \textcolor{blue}{e}A_{n-1,k}$$

We have three proofs:

- Two Digraph proofs using the Lindström-Gessel-Viennot lemma
  - Bijection between paths in the digraph to set partitions.
  - Exploiting the structure of the digraph.
- Algebraic proof

# An *ace* digraph




1

*a*

*c*

*e*

# An *ace* digraph



We have a weight preserving bijection between paths from  $u_n \rightarrow v_k$  and set partitions of  $\{1, \dots, n+1\}$  into  $n-k+1$  parts.

$$A_{n,k} = (n - k + 1)A_{n-1,k-1} \\ + (k + 1)A_{n-1,k}$$

Eulerian triangle conjecture

$$A_{n,k} = [a(n - k) + c]A_{n-1,k-1} \\ + (dk + e)A_{n-1,k}$$



*acde* conjecture

$$A_{n,k} = [a(n - k) + c]A_{n-1,k-1} \\ + (dk + e)A_{n-1,k} \\ + [f(n - 2) + g]A_{n-2,k-1}$$

*acdefg* conjecture

$$A_{n,k} = [a(n - k) + c]A_{n-1,k-1} \\ + \left( \sum_{i=0}^k d_i \right) A_{n-1,k} \\ + [f(n - 2) + g]A_{n-2,k-1}$$

*acd<sup>d</sup>fg* conjecture



10/27/20

J-fraction for new generating polynomials of TTSa with  $g=f$

$$\delta_n = [c + \binom{n}{2}d] + [c + \binom{n}{2}a]x$$

$$\beta_n = n[cd + ae + f + (n-1)ad]x$$

see conf from only action "Eulerian polynomials ..."

Stokel-Zeng have first J-fraction for permutations

$$\delta_0 = w_0$$

$$\delta_n = [x_2 + (n-1)u_2] + [y_2 + (n-1)v_2] + w_n \text{ for } n \geq 1$$

$$\beta_n = [x_1 + (n-1)u_1][y_1 + (n-1)v_1]$$

so we could take ...

# Thank you

Meme images from internet and digraph drawn by Tomack Gilmore.

*The essence of mathematics is proving theorems - and so, that is what mathematicians do: They prove theorems. But to tell the truth, what they really want to prove, once in their lifetime, is a Lemma, like the one by Fatou in analysis, the Lemma of Gauss in number theory, or the Burnside-Frobenius Lemma in combinatorics.*

...

- Chapter 25, Lattice Paths and Determinants  
Proofs from the Book

# LGV lemma on planar digraphs

- Let  $G$  be a directed acyclic graph embedded in the plane such that



# LGV lemma on planar digraphs

- Let  $G$  be a directed acyclic graph embedded in the plane such that
  - Edges directed left to right
  - Edges have non-negative weights

# LGV lemma on planar digraphs

- Let  $G$  be a directed acyclic graph embedded in the plane such that
  - Edges directed left to right
  - Edges have non-negative weights
  - Sources ( $u_n$ ) and sinks ( $v_n$ ) on boundary

# LGV lemma on planar digraphs


- Let  $G$  be a directed acyclic graph embedded in the plane such that
  - Edges directed left to right
  - Edges have non-negative weights
  - Sources ( $u_n$ ) and sinks ( $v_n$ ) on boundary
  - Sources – clockwise direction

# LGV lemma on planar digraphs

- Let  $G$  be a directed acyclic graph embedded in the plane such that
  - Edges directed left to right
  - Edges have non-negative weights
  - Sources ( $u_n$ ) and sinks ( $v_n$ ) on boundary
  - Sources – clockwise direction
  - Sinks – anticlockwise direction.

# LGV lemma on planar digraphs

- Let  $G$  be a directed acyclic graph embedded in the plane such that
  - Edges directed left to right
  - Edges have non-negative weights
  - Sources ( $u_n$ ) and sinks ( $v_n$ ) on boundary
  - Sources – clockwise direction
  - Sinks – anticlockwise direction.



Example. Edges weighted 1.

# LGV lemma on planar digraphs

- Let  $G$  be a directed acyclic graph embedded in the plane such that
  - Edges directed left to right
  - Edges have non-negative weights
  - Sources ( $u_n$ ) and sinks ( $v_n$ ) on boundary
  - Sources – clockwise direction
  - Sinks – anticlockwise direction.
- Weight of a path – product of edges


# LGV lemma on planar digraphs


- Let  $G$  be a directed acyclic graph embedded in the plane such that
  - Edges directed left to right
  - Edges have non-negative weights
  - Sources ( $u_n$ ) and sinks ( $v_n$ ) on boundary
  - Sources – clockwise direction
  - Sinks – anticlockwise direction.
- Weight of a path – product of edges
- Path matrix

# LGV lemma on planar digraphs

- Let  $G$  be a directed acyclic graph embedded in the plane such that
  - Edges directed left to right
  - Edges have non-negative weights
  - Sources ( $u_n$ ) and sinks ( $v_n$ ) on boundary
  - Sources – clockwise direction
  - Sinks – anticlockwise direction.
- Weight of a path – product of edges
- Path matrix
  - Entry  $(n, k)$  – sum of paths from  $u_n$  to  $v_k$








Path matrix

$$\begin{bmatrix} 1 & & & & & \\ 1 & 1 & & & & \\ 1 & 2 & 1 & & & \\ 1 & 3 & 3 & 1 & & \\ 1 & 4 & 6 & 4 & 1 & \\ 1 & 5 & 10 & 10 & 5 & 1 \end{bmatrix}$$



Path matrix

$$\begin{bmatrix} 1 & & & & & \\ 1 & 1 & & & & \\ 1 & 2 & 1 & & & \\ 1 & 3 & 3 & 1 & & \\ 1 & 4 & 6 & 4 & 1 & \\ 1 & 5 & 10 & 10 & 5 & 1 \end{bmatrix}$$

Binomial triangle!

# LGV lemma on planar digraphs

- Let  $G$  be a directed acyclic graph embedded in the plane such that
  - Edges directed left to right
  - Edges have non-negative weights
  - Sources ( $u_n$ ) and sinks ( $v_n$ ) on boundary
  - Sources – clockwise direction
  - Sinks – anticlockwise direction.
- Weight of a path – product of edges
- Path matrix
  - Entry  $(n, k)$  – sum of paths from  $u_n$  to  $v_k$

# LGV lemma on planar digraphs

- Let  $G$  be a directed acyclic graph embedded in the plane such that
  - Edges directed left to right
  - Edges have non-negative weights
  - Sources ( $u_n$ ) and sinks ( $v_n$ ) on boundary
  - Sources – clockwise direction
  - Sinks – anticlockwise direction.
- Weight of a path – product of edges
- Path matrix
  - Entry  $(n, k)$  – sum of paths from  $u_n$  to  $v_k$
- LGV lemma says that the path matrix is totally positive

10/27/20

J-fraction for new generating polynomials of TTSa with  $g=f$

$$\delta_n = [c + \binom{n}{2}d] + [c + \binom{n}{2}a]x$$

$$\beta_n = n[cd + ae + f + (n-1)ad]x$$

see conf from only action "Eulerian polynomials ..."

Stokel-Zeng have first J-fraction for permutations

$$\delta_0 = w_0$$

$$\delta_n = [x_2 + (n-1)u_2] + [y_2 + (n-1)v_2] + w_n \text{ for } n \geq 1$$

$$\beta_n = [x_1 + (n-1)u_1][y_1 + (n-1)v_1]$$

so we could take ...

# Thank you

Meme images from internet and *ace*-digraph drawn by Tomack Gilmore.