

Continued fractions using a Laguerre digraph interpretation of the Foata–Zeilberger bijection and its variants

Bishal Deb (he/him)

Sorbonne Université and Université Paris-Cité, CNRS, LPSM

April 16, 2024
Institut Camille Jordan

arxiv: 2304.14487

- ① Continued fractions and enumerative combinatorics
 - ① Classical continued fractions
 - ② Sokal–Zeng’s results for factorials
 - ③ D.–Sokal’s results for Genocchi and median Genocchi numbers
 - ④ Conjectures
- ② Proof overview of existing results
 - ① Flajolet’s combinatorial interpretation
 - ② Foata–Zeilberger bijection
- ③ What’s new
 - ① Laguerre digraphs
 - ② New interpretation of the FZ bijection
- ④ The story continues ...

- ➊ Continued fractions and enumerative combinatorics
 - ➊ Classical continued fractions
 - ➋ Sokal–Zeng's results for factorials
 - ➌ D.–Sokal's results for Genocchi and median Genocchi numbers
 - ➍ Conjectures
- ➋ Proof overview of existing results
 - ➊ Flajolet's combinatorial interpretation
 - ➋ Foata–Zeilberger bijection
- ➌ What's new
 - ➊ Laguerre digraphs
 - ➋ New interpretation of the FZ bijection
- ➍ The story continues ...

Continued fractions for ogf

Given sequence $(a_n)_{n \geq 0}$

Continued fractions for ogf

Given sequence $(a_n)_{n \geq 0}$

want to write

$$\sum_{n=0}^{\infty} a_n t^n = \frac{1}{1 - \frac{\alpha_1 t}{1 - \frac{\alpha_2 t}{1 - \ddots}}}$$

Continued fractions for ogf

Given sequence $(a_n)_{n \geq 0}$

want to write

$$\sum_{n=0}^{\infty} a_n t^n = \frac{1}{1 - \frac{\alpha_1 t}{1 - \frac{\alpha_2 t}{1 - \ddots}}}$$

Stieltjes-type continued fractions (S-fractions)

An example

$$\sum_{n=0} n! t^n = \frac{1}{1 - \frac{1 \cdot t}{1 - \frac{1 \cdot t}{1 - \frac{2 \cdot t}{1 - \frac{2 \cdot t}{1 - \frac{3 \cdot t}{1 - \frac{3 \cdot t}{1 - \dots}}}}}}$$

This line of thought goes back to Euler

$$A = \frac{x}{1+x} + \frac{\frac{x}{1+x}}{1+\frac{x}{1+x}} + \frac{\frac{\frac{x}{1+x}}{1+\frac{x}{1+x}}}{1+\frac{\frac{x}{1+x}}{1+\frac{x}{1+x}}} + \dots$$

§. 22. Quomodo autem huiusmodi fractionum, continuorum valor sit investigandus, alibi ostendi: Scilicet cum singulorum denominatorum partes integrae sint unitates, soli numeratores in computum veniunt; sit ergo $x=1$, atque investigatio summae A sequenti modo instituetur:

$$A = \begin{matrix} 0 & 1 & 2 & 3 & 4 & 5 & 6 & 7 & 8 & 9 & 10 & 11 & 12 & 13 & 14 & 15 & 16 & 17 & 18 & 19 & 20 & 21 & 22 & 23 & 24 & 25 & 26 & 27 & 28 & 29 & 30 & 31 & 32 & 33 & 34 & 35 & 36 & 37 & 38 & 39 & 40 & 41 & 42 & 43 & 44 & 45 & 46 & 47 & 48 & 49 & 50 & 51 & 52 & 53 & 54 & 55 & 56 & 57 & 58 & 59 & 60 & 61 & 62 & 63 & 64 & 65 & 66 & 67 & 68 & 69 & 70 & 71 & 72 & 73 & 74 & 75 & 76 & 77 & 78 & 79 & 80 & 81 & 82 & 83 & 84 & 85 & 86 & 87 & 88 & 89 & 90 & 91 & 92 & 93 & 94 & 95 & 96 & 97 & 98 & 99 \end{matrix}$$

num. 1, 1, 2, 2, 3, 3, 4, 4, 5, 5, etc:

Fractiones nimirum hic exhibitae continuo propius ad verum valorem ipsius A accedunt, et quidem alternim eo sunt maiores et minores; ita ut fit:

Tom. V. Nou. Com.

Ff

As

Euler (1760)

Jacobi-type continued fractions

$$\frac{1}{1 - \gamma_0 t - \frac{\beta_1 t^2}{1 - \gamma_1 t - \frac{\beta_2 t^2}{1 - \ddots}}}$$

J-fractions in short

J-fraction for $n!$

$$\sum_{n=0}^{\infty} n! t^n = \frac{1}{1 - \gamma_0 t - \frac{\beta_1 t^2}{1 - \gamma_1 t - \frac{\beta_2 t^2}{1 - \ddots}}}$$

where

$$\begin{aligned}\gamma_n &= 2n + 1 \\ \beta_n &= n^2\end{aligned}$$

Sokal–Zeng's (2022) reverse program

Start with

$$\sum_{n=0}^{\infty} n! t^n = \frac{1}{1 - \gamma_0 t - \frac{\beta_1 t^2}{1 - \gamma_1 t - \frac{\beta_2 t^2}{1 - \ddots}}}$$

where

$$\gamma_n = 2n + 1$$

$$\beta_n = n^2$$

Sokal–Zeng's (2022) reverse program

Start with

$$\sum_{n=0}^{\infty} n! t^n = \frac{1}{1 - \gamma_0 t - \frac{\beta_1 t^2}{1 - \gamma_1 t - \frac{\beta_2 t^2}{1 - \ddots}}}$$

where

$$\begin{aligned}\gamma_n &= 2n + 1 \\ \beta_n &= n^2\end{aligned}$$

Instead consider

$$\begin{aligned}\gamma_0 &= w_0 \\ \gamma_n &= [x_2 + (n-1)u_2] + [y_2 + (n-1)v_2] + w_n \\ \beta_n &= [x_1 + (n-1)u_1][y_1 + (n-1)v_1]\end{aligned}$$

Sokal–Zeng's (2022) reverse program

Start with

$$\sum_{n=0}^{\infty} n! t^n = \frac{1}{1 - \gamma_0 t - \frac{\beta_1 t^2}{1 - \gamma_1 t - \frac{\beta_2 t^2}{1 - \ddots}}}$$

where

$$\begin{aligned}\gamma_n &= 2n + 1 \\ \beta_n &= n^2\end{aligned}$$

Instead consider

$$\begin{aligned}\gamma_0 &= w_0 \\ \gamma_n &= [x_2 + (n-1)u_2] + [y_2 + (n-1)v_2] + w_n \\ \beta_n &= [x_1 + (n-1)u_1][y_1 + (n-1)v_1]\end{aligned}$$

Question

Find the 10 permutation statistics $x_1, x_2, y_1, y_2, u_1, u_2, v_1, v_2, w_0, (w_n)_{n \geq 1}$

Continued fractions by Sokal–Zeng (2022) for permutations

“First” (Cycles not counted)	“Second” (Cycles counted)
J-fraction with 10-statistics	Conjecture: J-fraction with 10-statistics

Continued fractions by Sokal–Zeng (2022) for permutations

“First” (Cycles not counted)	“Second” (Cycles counted)
J-fraction with 10-statistics	Conjecture: J-fraction with 10-statistics ↓ J-fraction with 9-statistics

Continued fractions by Sokal–Zeng (2022) for permutations

“First” (Cycles not counted)	“Second” (Cycles counted)
<p data-bbox="83 370 529 405">J-fraction with 10-statistics</p> <p data-bbox="157 418 179 453">↑↑</p> <p data-bbox="83 458 513 534">p, q-generalisation: J-fraction with 18 statistics</p>	<p data-bbox="665 280 1308 314">Conjecture: J-fraction with 10-statistics</p> <p data-bbox="739 328 761 363">⇓</p> <p data-bbox="665 370 1090 405">J-fraction with 9-statistics</p> <p data-bbox="739 418 761 453">↑↑</p> <p data-bbox="665 458 1094 534">p, q-generalisation: J-fraction with 15 statistics</p>

Continued fractions by Sokal–Zeng (2022) for permutations

“First” (Cycles not counted)	“Second” (Cycles counted)
<p data-bbox="83 370 529 405">J-fraction with 10-statistics</p> <p data-bbox="159 418 179 453">↑↑</p> <p data-bbox="83 458 406 493"><i>p, q</i>-generalisation:</p> <p data-bbox="83 503 513 537">J-fraction with 18 statistics</p> <p data-bbox="159 540 179 576">↑↑</p>	<p data-bbox="665 280 1308 314">Conjecture: J-fraction with 10-statistics</p> <p data-bbox="740 328 761 363">⇓</p> <p data-bbox="665 368 1092 403">J-fraction with 9-statistics</p> <p data-bbox="740 416 761 451">↑↑</p> <p data-bbox="665 456 985 491"><i>p, q</i>-generalisation:</p> <p data-bbox="665 501 1094 535">J-fraction with 15 statistics</p> <p data-bbox="740 538 761 573">↑↑</p>

Continued fractions by Sokal–Zeng (2022) for permutations

“First” (Cycles not counted)	“Second” (Cycles counted)
<p>J-fraction with 10-statistics</p> <p>↑↑</p> <p>p, q-generalisation:</p> <p>J-fraction with 18 statistics</p> <p>↑↑</p> <p>Master J-fraction:</p> <p>four infinite 2-parameter families</p> <p>one infinite 1-parameter family</p>	<p>Conjecture: J-fraction with 10-statistics</p> <p>⇓</p> <p>J-fraction with 9-statistics</p> <p>↑↑</p> <p>p, q-generalisation:</p> <p>J-fraction with 15 statistics</p> <p>↑↑</p> <p>Master J-fraction:</p> <p>three infinite 2-parameter families</p> <p>two infinite 1-parameter family</p> <p>and one statistic for counting cycles</p>

Continued fractions by Sokal–Zeng (2022) for permutations

“First” (Cycles not counted)	“Second” (Cycles counted)
<p data-bbox="83 370 529 405">J-fraction with 10-statistics</p> <p data-bbox="159 418 179 452">↑↑</p> <p data-bbox="83 459 404 494">p, q-generalisation:</p> <p data-bbox="83 505 513 539">J-fraction with 18 statistics</p> <p data-bbox="159 536 179 570">↑↑</p> <p data-bbox="83 598 392 632">Master J-fraction:</p> <p data-bbox="83 644 598 678">four infinite 2-parameter families</p> <p data-bbox="83 689 568 724">one infinite 1-parameter family</p> <p data-bbox="83 837 618 919">Proof: Foata–Zeilberger bijection (1990)</p>	<p data-bbox="665 280 1308 314">Conjecture: J-fraction with 10-statistics</p> <p data-bbox="740 328 761 362">⇓</p> <p data-bbox="665 369 1092 404">J-fraction with 9-statistics</p> <p data-bbox="740 417 761 451">↑↑</p> <p data-bbox="665 458 986 493">p, q-generalisation:</p> <p data-bbox="665 504 1094 538">J-fraction with 15 statistics</p> <p data-bbox="740 535 761 569">↑↑</p> <p data-bbox="665 577 972 611">Master J-fraction:</p> <p data-bbox="665 622 1195 656">three infinite 2-parameter families</p> <p data-bbox="665 668 1149 702">two infinite 1-parameter family</p> <p data-bbox="665 713 1230 747">and one statistic for counting cycles</p> <p data-bbox="665 837 1012 919">Proof: Biane bijection (1993)</p>

At around the same time, Blitvić–Steingrímsson (2021) independently came up with a 14-variable continued fraction

At around the same time, Blitvić–Steingrímsson (2021) independently came up with a 14-variable continued fraction

They later discovered that Randrianarivony (1998) had a 17-variable continued fraction.

Results and conjecture for factorials

Cycle classification

For a permutation σ , compare each i with $\sigma(i)$ and $\sigma^{-1}(i)$:

Cycle classification

For a permutation σ , compare each i with $\sigma(i)$ and $\sigma^{-1}(i)$:

- cycle valley $\sigma^{-1}(i) > i < \sigma(i)$
- cycle peaks $\sigma^{-1}(i) < i > \sigma(i)$
- cycle double rise $\sigma^{-1}(i) < i < \sigma(i)$
- cycle double fall $\sigma^{-1}(i) > i > \sigma(i)$
- fixed point $i = \sigma(i) = \sigma^{-1}(i)$

Consider σ as $\sigma(1)\sigma(2)\dots\sigma(n)$:

Record classification

Consider σ as $\sigma(1)\sigma(2)\dots\sigma(n)$:

- i is record if for every $j < i$ we have $\sigma(j) < \sigma(i)$
left-to-right-maxima
- i is antirecord if for every $j > i$ we have $\sigma(j) > \sigma(i)$
right-to-left-minima

Record classification

Consider σ as $\sigma(1)\sigma(2)\dots\sigma(n)$:

- i is record if for every $j < i$ we have $\sigma(j) < \sigma(i)$
left-to-right-maxima
- i is antirecord if for every $j > i$ we have $\sigma(j) > \sigma(i)$
right-to-left-minima

Each i is one of the following four types:

Record classification

Consider σ as $\sigma(1)\sigma(2)\dots\sigma(n)$:

- i is record if for every $j < i$ we have $\sigma(j) < \sigma(i)$
left-to-right-maxima
- i is antirecord if for every $j > i$ we have $\sigma(j) > \sigma(i)$
right-to-left-minima

Each i is one of the following four types:





- rar - record-antirecord
- erec - exclusive record
- earec - exclusive antirecord
- nrar - neither record-antirecord

Record classification

Consider σ as $\sigma(1)\sigma(2)\dots\sigma(n)$:

- i is record if for every $j < i$ we have $\sigma(j) < \sigma(i)$
left-to-right-maxima
- i is antirecord if for every $j > i$ we have $\sigma(j) > \sigma(i)$
right-to-left-minima

Each i is one of the following four types:

- rar - record-antirecord 
- errec - exclusive record 
- earec - exclusive antirecord 
- nrar - neither record-antirecord 

Each i is one of the following ten (not 20) types:

Record-and-cycle classification

Each i is one of the following ten (not 20) types:

	cpeak	cval	cdrise	cdfall	fix
erec	eareccpeak	ereccval	ereccdrise	eareccdfall	rar nrfix
earec					
rar	nrcpeak	nrcval	nrhdrise	nrhdrfall	
nrar					

Continued fractions counting permutation statistics

Consider 10-variable polynomials

$$P_n(x_1, x_2, y_1, y_2, u_1, u_2, v_1, v_2, w, z) = \sum_{\sigma \in \mathfrak{S}_n} x_1^{\text{eareccpeak}(\sigma)} x_2^{\text{eareccdfall}(\sigma)} y_1^{\text{ereccval}(\sigma)} y_2^{\text{ereccdrise}(\sigma)} z^{\text{rar}(\sigma)} \times \\ u_1^{\text{nrcpeak}(\sigma)} u_2^{\text{nrcdfall}(\sigma)} v_1^{\text{nrcval}(\sigma)} v_2^{\text{nrcdrise}(\sigma)} w^{\text{nrfix}(\sigma)}$$

Continued fractions counting permutation statistics

Consider 10-variable polynomials

$$P_n(x_1, x_2, y_1, y_2, u_1, u_2, v_1, v_2, w, z) = \sum_{\sigma \in \mathfrak{S}_n} x_1^{\text{eareccpeak}(\sigma)} x_2^{\text{eareccdfall}(\sigma)} y_1^{\text{ereccval}(\sigma)} y_2^{\text{ereccdrise}(\sigma)} z^{\text{rar}(\sigma)} \times \\ u_1^{\text{nrcpeak}(\sigma)} u_2^{\text{nrcdfall}(\sigma)} v_1^{\text{nrcval}(\sigma)} v_2^{\text{nrcdrise}(\sigma)} w^{\text{nrfix}(\sigma)}$$

Theorem (Sokal–Zeng (2022) First J-fraction for permutations)

$$\sum_{n=0}^{\infty} P_n(x_1, x_2, y_1, y_2, u_1, u_2, v_1, v_2, w, z) t^n \\ = \frac{1}{1 - z \cdot t - \frac{1}{x_1 y_1 \cdot t^2} \frac{(x_1 + u_1)(y_1 + v_1) \cdot t^2}{1 - ((x_2 + u_2) + (y_2 + v_2) + w) \cdot t - \frac{(x_1 + 2u_1)(y_1 + 2v_1) \cdot t^2}{1 - \ddots}}}$$

Continued fractions counting permutation statistics

Consider 10-variable polynomials

$$P_n(x_1, x_2, y_1, y_2, u_1, u_2, v_1, v_2, w, z) = \sum_{\sigma \in \mathfrak{S}_n} x_1^{\text{eareccpeak}(\sigma)} x_2^{\text{eareccdfall}(\sigma)} y_1^{\text{ereccval}(\sigma)} y_2^{\text{ereccdrise}(\sigma)} z^{\text{rar}(\sigma)} \times \\ u_1^{\text{nrcpeak}(\sigma)} u_2^{\text{nrcdfall}(\sigma)} v_1^{\text{nrcval}(\sigma)} v_2^{\text{nrcdrise}(\sigma)} w^{\text{nrfix}(\sigma)}$$

Theorem (Sokal–Zeng (2022) First J-fraction for permutations)

$$\sum_{n=0}^{\infty} P_n(x_1, x_2, y_1, y_2, u_1, u_2, v_1, v_2, w, z) t^n \\ = \frac{1}{1 - z \cdot t - \frac{1}{x_1 y_1 \cdot t^2} \frac{(x_1 + u_1)(y_1 + v_1) \cdot t^2}{1 - ((x_2 + u_2) + (y_2 + v_2) + w) \cdot t - \frac{(x_1 + 2u_1)(y_1 + 2v_1) \cdot t^2}{1 - \ddots}}}$$

Proof uses the Foata–Zeilberger bijection (1990)

Can we count cycles as well?

Consider 11-variable polynomials

$$P_n(x_1, x_2, y_1, y_2, u_1, u_2, v_1, v_2, w, z) = \sum_{\sigma \in \mathfrak{S}_n} x_1^{\text{eareccpeak}(\sigma)} x_2^{\text{eareccdfall}(\sigma)} y_1^{\text{ereccval}(\sigma)} y_2^{\text{ereccdrise}(\sigma)} z^{\text{rar}(\sigma)} \times \\ u_1^{\text{nrcpeak}(\sigma)} u_2^{\text{nrcdfall}(\sigma)} v_1^{\text{nrcval}(\sigma)} v_2^{\text{nrcdrise}(\sigma)} w^{\text{nrfix}(\sigma)} \lambda^{\text{cyc}(\sigma)}$$

Can we count cycles as well?

Consider 11-variable polynomials

$$P_n(x_1, x_2, y_1, y_2, u_1, u_2, v_1, v_2, w, z) = \sum_{\sigma \in \mathfrak{S}_n} x_1^{\text{eareccpeak}(\sigma)} x_2^{\text{eareccdfall}(\sigma)} y_1^{\text{ereccval}(\sigma)} y_2^{\text{ereccdrise}(\sigma)} z^{\text{rar}(\sigma)} \times \\ u_1^{\text{nrcpeak}(\sigma)} u_2^{\text{nrcdfall}(\sigma)} v_1^{\text{nrcval}(\sigma)} v_2^{\text{nrcdrise}(\sigma)} w^{\text{nrfix}(\sigma)} \lambda^{\text{cyc}(\sigma)}$$

No nice J-fraction!

Can we count cycles as well?

Consider 11-variable polynomials

$$P_n(x_1, x_2, y_1, y_2, u_1, u_2, v_1, v_2, w, z) = \sum_{\sigma \in \mathfrak{S}_n} x_1^{\text{eareccpeak}(\sigma)} x_2^{\text{eareccdfall}(\sigma)} y_1^{\text{ereccval}(\sigma)} y_2^{\text{ereccdrise}(\sigma)} z^{\text{rar}(\sigma)} \times \\ u_1^{\text{nrcpeak}(\sigma)} u_2^{\text{nrcdfall}(\sigma)} v_1^{\text{nrcval}(\sigma)} v_2^{\text{nrcdrise}(\sigma)} w^{\text{nrfix}(\sigma)} \lambda^{\text{cyc}(\sigma)}$$

No nice J-fraction!

But can obtain J-fraction by specialising $y_1 = v_1$:

Can we count cycles as well?

Consider 11-variable polynomials

$$P_n(x_1, x_2, y_1, y_2, u_1, u_2, v_1, v_2, w, z) = \sum_{\sigma \in \mathfrak{S}_n} x_1^{\text{eareccpeak}(\sigma)} x_2^{\text{eareccdfall}(\sigma)} y_1^{\text{ereccval}(\sigma)} y_2^{\text{ereccdrise}(\sigma)} z^{\text{rar}(\sigma)} \times \\ u_1^{\text{nrcpeak}(\sigma)} u_2^{\text{nrcdfall}(\sigma)} v_1^{\text{nrcval}(\sigma)} v_2^{\text{nrcdrise}(\sigma)} w^{\text{nrfix}(\sigma)} \lambda^{\text{cyc}(\sigma)}$$

No nice J-fraction!

But can obtain J-fraction by specialising $y_1 = v_1$:

Conjecture (Sokal–Zeng (2022))

$$\sum_{n=0}^{\infty} P_n(x_1, x_2, y_1, y_2, u_1, u_2, y_1, v_2, w, z, \lambda) t^n \\ = \frac{1}{1 - \lambda z \cdot t - \frac{\lambda x_1 y_1 \cdot t^2}{1 - (x_2 + y_2 + \lambda w) \cdot t - \frac{(\lambda + 1)(x_1 + u_1) y_1 \cdot t^2}{1 - ((x_2 + v_2) + (y_2 + v_2) + \lambda w) \cdot t - \frac{(\lambda + 2)(x_1 + 2u_1) y_1 \cdot t^2}{1 - \dots}}}}$$

Sokal–Zeng conjectured a continued fraction for 11 variable polynomials involving one specialisation $y_1 = v_1$.

Sokal–Zeng conjectured a continued fraction for 11 variable polynomials involving one specialisation $y_1 = v_1$.

They could only prove with two specialisations $y_1 = v_1$ and $y_2 = v_2$

Sokal–Zeng conjectured a continued fraction for 11 variable polynomials involving one specialisation $y_1 = v_1$.

They could only prove with two specialisations $y_1 = v_1$ and $y_2 = v_2$
Second J-fraction for permutations

Sokal–Zeng conjectured a continued fraction for 11 variable polynomials involving one specialisation $y_1 = v_1$.

They could only prove with two specialisations $y_1 = v_1$ and $y_2 = v_2$
Second J-fraction for permutations

Used Biane bijection (1993).

Sokal–Zeng conjectured a continued fraction for 11 variable polynomials involving one specialisation $y_1 = v_1$.

They could only prove with two specialisations $y_1 = v_1$ and $y_2 = v_2$
Second J-fraction for permutations

Used Biane bijection (1993).

Genocchi and median Genocchi numbers

The Genocchi numbers are given by

$$t \tan\left(\frac{t}{2}\right) = \sum_{n=0}^{\infty} g_n \frac{t^{2n+2}}{(2n+2)!}$$

Genocchi numbers

The Genocchi numbers are given by

$$t \tan\left(\frac{t}{2}\right) = \sum_{n=0}^{\infty} g_n \frac{t^{2n+2}}{(2n+2)!}$$

The first few numbers are 1, 1, 3, 17, 155, 2073, ...

Genocchi numbers:

D-e-semiderangements

$$g_n = \#\{\sigma \in \mathfrak{S}_{2n} \mid 2i > \sigma(2i) \text{ and } 2i - 1 \leq \sigma(2i - 1)\}$$

D-o-semiderangements

$$= \#\{\sigma \in \mathfrak{S}_{2n} \mid 2i \geq \sigma(2i) \text{ and } 2i - 1 < \sigma(2i - 1)\}$$

Genocchi numbers:

D-e-semiderangements

$$g_n = \#\{\sigma \in \mathfrak{S}_{2n} \mid 2i > \sigma(2i) \text{ and } 2i - 1 \leq \sigma(2i - 1)\}$$

D-o-semiderangements

$$= \#\{\sigma \in \mathfrak{S}_{2n} \mid 2i \geq \sigma(2i) \text{ and } 2i - 1 < \sigma(2i - 1)\}$$

Median Genocchi numbers:

D-permutations (introduced by Lazar and Wachs (2019))

$$h_{n+1} = |\mathfrak{D}_{2n}| = \#\{\sigma \in \mathfrak{S}_{2n} \mid 2i \geq \sigma(2i) \text{ and } 2i - 1 \leq \sigma(2i - 1)\}$$

D-derangements

$$h_n = \#\{\sigma \in \mathfrak{S}_{2n} \mid 2i > \sigma(2i) \text{ and } 2i - 1 < \sigma(2i - 1)\}$$

Genocchi numbers:

D-e-semiderangements

$$g_n = \#\{\sigma \in \mathfrak{S}_{2n} \mid 2i > \sigma(2i) \text{ and } 2i - 1 \leq \sigma(2i - 1)\}$$

D-o-semiderangements

$$= \#\{\sigma \in \mathfrak{S}_{2n} \mid 2i \geq \sigma(2i) \text{ and } 2i - 1 < \sigma(2i - 1)\}$$

Median Genocchi numbers:

D-permutations (introduced by Lazar and Wachs (2019))

$$h_{n+1} = |\mathfrak{D}_{2n}| = \#\{\sigma \in \mathfrak{S}_{2n} \mid 2i \geq \sigma(2i) \text{ and } 2i - 1 \leq \sigma(2i - 1)\}$$

D-derangements

$$h_n = \#\{\sigma \in \mathfrak{S}_{2n} \mid 2i > \sigma(2i) \text{ and } 2i - 1 < \sigma(2i - 1)\}$$

D — Dumont-like

Continued fractions for median Genocchi numbers

The once-shifted median Genocchi numbers h_{n+1} have the following Thron-type continued fraction

$$\sum_{n=0}^{\infty} h_{n+1} t^n = \frac{1}{1 - t - \frac{1t}{1 - \frac{4t}{1 - \frac{4t}{1 - \frac{9t}{1 - \dots}}}}}$$

Story for Genocchi and median Genocchi

D.-Sokal (2022):

Story for Genocchi and median Genocchi

D.-Sokal (2022):

“First” (Cycles not counted)	“Second” (Cycles counted)
<p>0-T-fraction with 12-statistics</p> <p>↑↑</p> <p>p, q-generalisation:</p> <p>0-T-fraction with 22 statistics</p> <p>↑↑</p> <p>Master T-fraction:</p> <p>four infinite 2-parameter families two infinite 1-parameter family</p> <p>Proof:</p> <p>FZ-like bijection</p>	<p>Conjecture: 0-T-fraction with 12-statistics</p> <p>0-T-fraction with 12-statistics</p> <p>↑↑</p> <p>p, q-generalisation:</p> <p>0-T-fraction with 21 statistics</p> <p>↑↑</p> <p>Master T-fraction:</p> <p>three infinite 2-parameter families, three infinite 1-parameter families, and one statistic for counting cycles</p> <p>Proof:</p> <p>Biane-like bijection</p>
Variant forms with slightly different statistics	

Randrianarivony–Zeng conjecture (1996)

Define the polynomials

where
$$G_n(x, y, \bar{x}, \bar{y}) = \sum_{\sigma \in \mathfrak{D}_{2n}^o} x^{\text{comi}(\sigma)} y^{\text{lema}(\sigma)} \bar{x}^{\text{cemi}(\sigma)} \bar{y}^{\text{remi}(\sigma)}$$

Randrianarivony–Zeng conjecture (1996)

Define the polynomials

$$G_n(x, y, \bar{x}, \bar{y}) = \sum_{\sigma \in \mathfrak{D}_{2n}^o} x^{\text{comi}(\sigma)} y^{\text{lema}(\sigma)} \bar{x}^{\text{cemi}(\sigma)} \bar{y}^{\text{remi}(\sigma)}$$

where

- comi – odd cycle minima,
- lema – left-to-right maxima whose value is even
- cemi – even cycle minima
- remi – right-to-left minima whose value is even

Randrianarivony–Zeng conjecture (1996)

Define the polynomials

$$G_n(x, y, \bar{x}, \bar{y}) = \sum_{\sigma \in \mathfrak{D}_{2n}^o} x^{\text{comi}(\sigma)} y^{\text{lema}(\sigma)} \bar{x}^{\text{cemi}(\sigma)} \bar{y}^{\text{remi}(\sigma)}$$

where

- comi – odd cycle minima,
- lema – left-to-right maxima whose value is even
- cemi – even cycle minima
- remi – right-to-left minima whose value is even

Conjecture

The ogf of the polynomials $G_n(x, y, \bar{x}, \bar{y})$ has the following S-fraction

$$\sum_{n=0}^{\infty} G_n(x, y, \bar{x}, \bar{y}) t^n = \cfrac{1}{1 - \cfrac{xyt}{1 - \cfrac{1(\bar{x} + \bar{y})t}{1 - \cfrac{(x+1)(y+1)t}{1 - \cfrac{2(\bar{x} + \bar{y} + 1)t}{1 - \cfrac{(x+2)(y+2)t}{1 - \cfrac{3(\bar{x} + \bar{y} + 2)t}{\dots}}}}}} \quad (1)$$

Plot twist for Sokal–Zeng conjecture

“First” (Cycles not counted)

J-fraction with 10 statistics



p, q -generalisation:

J-fraction with 18 statistics



Master J-fraction:

four infinite 2-parameter families
one infinite 1-parameter family



Proof:

Foata–Zeilberger bijection (1990)

“Second” (Cycles counted)

~~Conjecture:~~ **J-fraction with 10 statistics**



J-fraction with 9 statistics



p, q -generalisation:

J-fraction with 15 statistics



Master J-fraction:



three infinite 2-parameter families
two infinite 1-parameter family
and one statistic for counting cycles



Proof:

Biane bijection (1993)

Conjectures of D.-Sokal, Randrianarivony–Zeng

“First” (Cycles not counted)	“Second” (Cycles counted)
<p data-bbox="72 267 563 301">0-T-fraction with 12 statistics</p> <p data-bbox="148 313 168 350">↑↑</p> <p data-bbox="72 355 395 389"><i>p, q</i>-generalisation:</p> <p data-bbox="72 399 543 434">0-T-fraction with 22 statistics</p> <p data-bbox="148 436 168 472">↑↑</p> <p data-bbox="72 474 609 601">Master T-fraction: four infinite 2-parameter families two infinite 1-parameter families</p> <div data-bbox="72 636 540 778"> Proof: FZ-like bijection</div>	<p data-bbox="653 179 1142 256">Conjecture: 0-T-fraction with 12 statistics</p> <p data-bbox="653 267 1142 301">0-T-fraction with 12 statistics</p> <p data-bbox="728 313 749 350">↑↑</p> <p data-bbox="653 355 964 389"><i>p, q</i>-generalisation</p> <p data-bbox="653 399 1123 434">0-T-fraction with 21 statistics</p> <p data-bbox="728 436 749 472">↑↑</p> <p data-bbox="653 474 1219 643">Master T-fraction: three infinite 2-parameter families, three infinite 1-parameter families, and one statistic for counting cycles</p> <div data-bbox="653 636 1120 767"> Proof: Biane-like bijection</div>
Variant forms	Variant forms \implies Randrianarivony–Zeng (1996) S-fraction with 4 statistics

Pan–Zeng ('23) came up with multivariate continued fractions for other objects enumerated by Genocchi numbers also introduced in work of Lazar's PhD thesis


- ① Continued fractions and enumerative combinatorics
 - ① Classical continued fractions
 - ② Sokal–Zeng’s results for factorials
 - ③ D.–Sokal’s results for Genocchi and median Genocchi numbers
 - ④ Conjectures
- ② Proof overview of existing results
 - ① Flajolet’s combinatorial interpretation
 - ② Foata–Zeilberger bijection
- ③ What’s new
 - ① Laguerre digraphs
 - ② New interpretation of the FZ bijection
- ④ The story continues ...

Motzkin paths

Consider a Motzkin path, let's say


Motzkin paths

Consider a Motzkin path, let's say



Motzkin paths

Consider a Motzkin path, let's say




Assign weights:

- $\nearrow : 1$
- \rightarrow from height $i \rightarrow i : \gamma_i$
- \searrow from height $i \rightarrow (i-1) : \beta_i$

Motzkin paths

Consider a Motzkin path, let's say




Assign weights:

- $\nearrow : 1$
- \rightarrow from height $i \rightarrow i : \gamma_i$
- \searrow from height $i \rightarrow (i-1) : \beta_i$

Motzkin paths

Consider a Motzkin path, let's say



$$\text{Weight} = \beta_1 \beta_2 \beta_3 \beta_4^2 \gamma_2 \gamma_3$$

Assign weights:

- $\nearrow : 1$
- \rightarrow from height $i \rightarrow i : \gamma_i$
- \searrow from height $i \rightarrow (i-1) : \beta_i$

Combinatorial Interpretation of J-fraction

J-fraction

$$\cfrac{1}{1 - \gamma_0 t - \cfrac{\beta_1 t^2}{1 - \gamma_1 t - \cfrac{\beta_2 t^2}{1 - \gamma_2 t - \cfrac{\beta_3 t^2}{\ddots}}}}$$

Combinatorial Interpretation of J-fraction

J-fraction

$$\frac{1}{1 - \gamma_0 t - \frac{\beta_1 t^2}{1 - \gamma_1 t - \frac{\beta_2 t^2}{1 - \gamma_2 t - \frac{\beta_3 t^2}{\ddots}}}} = \sum_{n=0}^{\infty} a_n t^n$$

Combinatorial Interpretation of J-fraction

J-fraction

$$\frac{1}{1 - \gamma_0 t - \frac{\beta_1 t^2}{1 - \gamma_1 t - \frac{\beta_2 t^2}{1 - \gamma_2 t - \frac{\beta_3 t^2}{\ddots}}}} = \sum_{n=0}^{\infty} a_n t^n$$

Theorem (Flajolet '80)

The a_n are weighted sum of Motzkin paths with n steps.

Combinatorial Interpretation of J-fraction

J-fraction

$$\frac{1}{1 - \gamma_0 t - \frac{\beta_1 t^2}{1 - \gamma_1 t - \frac{\beta_2 t^2}{1 - \gamma_2 t - \frac{\beta_3 t^2}{\ddots}}}} = \sum_{n=0}^{\infty} a_n t^n$$

Theorem (Flajolet '80)

The a_n are weighted sum of Motzkin paths with n steps.

- Stieltjes-type continued fractions — weighted Dyck paths (Flajolet '80)
- Thron-type continued fractions — weighted Schröder paths (Oste–Van der Jeugt (2015), Fusy–Guitter(2017), Josuat-Vergès (2018), Sokal (unpublished))

Combinatorial Interpretation of J-fraction

J-fraction

$$\frac{1}{1 - \gamma_0 t - \frac{\beta_1 t^2}{1 - \gamma_1 t - \frac{\beta_2 t^2}{1 - \gamma_2 t - \frac{\beta_3 t^2}{\ddots}}}} = \sum_{n=0}^{\infty} a_n t^n$$

Theorem (Flajolet '80)

The a_n are weighted sum of Motzkin paths with n steps.

- Stieltjes-type continued fractions — weighted Dyck paths (Flajolet '80)
- Thron-type continued fractions — weighted Schröder paths (Oste–Van der Jeugt (2015), Fusy–Guitter (2017), Josuat-Vergès (2018), Sokal (unpublished))

Gateway for proving continued fractions using bijective combinatorics :-D

Foata–Zeilberger bijection

excedance indices $F = \{i \in \sigma : \sigma(i) > i\} = \text{Cdrise} \cup \text{Cval}$

$$\begin{aligned} \text{excedance indices } F &= \{i \in \sigma : \sigma(i) > i\} = \text{Cdrise} \cup \text{Cval} \\ \text{excedance values } F' &= \{i \in \sigma : i > \sigma^{-1}(i)\} = \text{Cdrise} \cup \text{Cpeak} \end{aligned}$$

$$\begin{aligned}
\text{excedance indices } F &= \{i \in \sigma : \sigma(i) > i\} = \text{Cdrise} \cup \text{Cval} \\
\text{excedance values } F' &= \{i \in \sigma : i > \sigma^{-1}(i)\} = \text{Cdrise} \cup \text{Cpeak} \\
\text{antiexcedance indices } G &= \{i \in \sigma : \sigma(i) < i\} = \text{Cdfall} \cup \text{Cpeak}
\end{aligned}$$

$$\begin{aligned}
\text{excedance indices } F &= \{i \in \sigma : \sigma(i) > i\} = \text{Cdrise} \cup \text{Cval} \\
\text{excedance values } F' &= \{i \in \sigma : i > \sigma^{-1}(i)\} = \text{Cdrise} \cup \text{Cpeak} \\
\text{antiexcedance indices } G &= \{i \in \sigma : \sigma(i) < i\} = \text{Cdfall} \cup \text{Cpeak} \\
\text{antiexcedance values } G' &= \{i \in \sigma : i < \sigma^{-1}(i)\} = \text{Cdfall} \cup \text{Cval}
\end{aligned}$$

$$\begin{aligned}
\text{excedance indices } F &= \{i \in \sigma : \sigma(i) > i\} = \text{Cdrise} \cup \text{Cval} \\
\text{excedance values } F' &= \{i \in \sigma : i > \sigma^{-1}(i)\} = \text{Cdrise} \cup \text{Cpeak} \\
\text{antiexcedance indices } G &= \{i \in \sigma : \sigma(i) < i\} = \text{Cdfall} \cup \text{Cpeak} \\
\text{antiexcedance values } G' &= \{i \in \sigma : i < \sigma^{-1}(i)\} = \text{Cdfall} \cup \text{Cval} \\
\text{fixed points } H &= \{i \in \sigma : i = \sigma(i)\} = \text{Fix}
\end{aligned}$$

$$\begin{aligned}
\text{excedance indices } F &= \{i \in \sigma : \sigma(i) > i\} = \text{Cdrise} \cup \text{Cval} \\
\text{excedance values } F' &= \{i \in \sigma : i > \sigma^{-1}(i)\} = \text{Cdrise} \cup \text{Cpeak} \\
\text{antiexcedance indices } G &= \{i \in \sigma : \sigma(i) < i\} = \text{Cdfall} \cup \text{Cpeak} \\
\text{antiexcedance values } G' &= \{i \in \sigma : i < \sigma^{-1}(i)\} = \text{Cdfall} \cup \text{Cval} \\
\text{fixed points } H &= \{i \in \sigma : i = \sigma(i)\} = \text{Fix}
\end{aligned}$$

A permutation can be fully described the following data:

- Sets F, F', G, G', H

$$\begin{aligned}
\text{excedance indices } F &= \{i \in \sigma : \sigma(i) > i\} = \text{Cdrise} \cup \text{Cval} \\
\text{excedance values } F' &= \{i \in \sigma : i > \sigma^{-1}(i)\} = \text{Cdrise} \cup \text{Cpeak} \\
\text{antiexcedance indices } G &= \{i \in \sigma : \sigma(i) < i\} = \text{Cdfall} \cup \text{Cpeak} \\
\text{antiexcedance values } G' &= \{i \in \sigma : i < \sigma^{-1}(i)\} = \text{Cdfall} \cup \text{Cval} \\
\text{fixed points } H &= \{i \in \sigma : i = \sigma(i)\} = \text{Fix}
\end{aligned}$$

A permutation can be fully described the following data:

- Sets F, F', G, G', H
- $\sigma|_F : F \rightarrow F'$
- $\sigma|_G : G \rightarrow G'$

Foata–Zeilberger bijection

Foata–Zeilberger bijection:

$$\sigma \mapsto (\omega, \xi)$$

Foata–Zeilberger bijection

Foata–Zeilberger bijection:

$$\sigma \mapsto (\omega, \xi)$$

where

- ω is a Motzkin path, where the level steps come in three different colours (red, blue, green)

Foata–Zeilberger bijection

Foata–Zeilberger bijection:

$$\sigma \mapsto (\omega, \xi)$$

where

- ω is a Motzkin path, where the level steps come in three different colours (red, blue, green)
Correspond to F, F', G, G', H

Foata–Zeilberger bijection:

$$\sigma \mapsto (\omega, \xi)$$

where

- ω is a Motzkin path, where the level steps come in three different colours (red, blue, green)
Correspond to F, F', G, G', H
- $\xi = (\xi_1, \dots, \xi_n)$ are labels on the steps of the Motzkin paths

Foata–Zeilberger bijection

Foata–Zeilberger bijection:

$$\sigma \mapsto (\omega, \xi)$$

where

- ω is a Motzkin path, where the level steps come in three different colours (red, blue, green)
Correspond to F, F', G, G', H
- $\xi = (\xi_1, \dots, \xi_n)$ are labels on the steps of the Motzkin paths
Correspond to $\sigma|_F : F \rightarrow F'$ and $\sigma|_G : G \rightarrow G'$

Description of $\sigma \rightarrow \omega$

- If i is a cycle valley, step i is \nearrow
- If i is a cycle peak, step i is \searrow
- If i is a cycle double rise, cycle double fall or fixed, step i is \rightarrow , \rightarrow or \rightarrow respectively.

Description of labels $\sigma \rightarrow \xi$

For $i \in [n]$

$$\xi_i = \begin{cases} \#\{j: j < i \text{ and } \sigma(j) > \sigma(i)\} & \text{if } \sigma(i) > i & \text{if } i \in \text{Cval} \cup \text{Cdrise} \\ \#\{j: j > i \text{ and } \sigma(j) < \sigma(i)\} & \text{if } \sigma(i) < i & \text{if } i \in \text{Cpeak} \cup \text{Cdfall} \\ 0 & \text{if } \sigma(i) = i & \text{if } i \in \text{Fix} \end{cases}$$

An example

Let $\sigma = 715492638 = (1762)(3598)(4) \in \mathfrak{S}_9$.


- Cval = $\{1, 3\}$ - Cpeak = $\{7, 9\}$ - Cdrise = $\{5\}$ -
 Cdfall = $\{2, 6, 8\}$
- Fix = $\{4\}$

An example

Let $\sigma = 715492638 = (1762)(3598)(4) \in \mathfrak{S}_9$.

- $C_{\text{val}} = \{1, 3\}$
- $C_{\text{peak}} = \{7, 9\}$
- $C_{\text{drise}} = \{5\}$
- $C_{\text{dfall}} = \{2, 6, 8\}$
- $\text{Fix} = \{4\}$

The Motzkin path ω is




An example

Let $\sigma = 715492638 = (1762)(3598)(4) \in \mathfrak{S}_9$.

- Cval = {1, 3} - Cpeak = {7, 9} - Cdrise = {5} -
- Cdfall = {2, 6, 8}
- Fix = {4}

The Motzkin path ω is



The labels ξ and the sets F, F', G, G' are:

$i \in F$	1	3	5
$\sigma(i) \in F'$	7	5	9
ξ_i	0	1	0

$i \in G$	2	6	7	8	9
$\sigma(i) \in G'$	1	2	6	3	8
ξ_i	0	0	1	0	0

- ① Continued fractions and enumerative combinatorics
 - ① Classical continued fractions
 - ② Sokal–Zeng’s results for factorials
 - ③ D.–Sokal’s results for Genocchi and median Genocchi numbers
 - ④ Conjectures
- ② Proof overview of existing results
 - ① Flajolet’s combinatorial interpretation
 - ② Foata–Zeilberger bijection
- ③ What’s new
 - ① Laguerre digraphs
 - ② New interpretation of the FZ bijection
- ④ The story continues ...

Definition


A **Laguerre digraph** of size n is a directed graph on vertex set $\{1, \dots, n\}$ where each vertex has indegree 0 or 1 and outdegree 0 or 1.

Laguerre digraph

Definition

A **Laguerre digraph** of size n is a directed graph on vertex set $\{1, \dots, n\}$ where each vertex has indegree 0 or 1 and outdegree 0 or 1.

Example:




Laguerre digraph

Definition

A **Laguerre digraph** of size n is a directed graph on vertex set $\{1, \dots, n\}$ where each vertex has indegree 0 or 1 and outdegree 0 or 1.

Example:



Connected components


- Directed cycle
- Directed paths

Laguerre digraph

Definition

A **Laguerre digraph** of size n is a directed graph on vertex set $\{1, \dots, n\}$ where each vertex has indegree 0 or 1 and outdegree 0 or 1.

Example:



Connected components

- Directed cycle
- Directed paths

Generalise permutations

An equivalent object with a different definition was introduced by Foata–Strehl (1984) — Laguerre configurations

An equivalent object with a different definition was introduced by Foata–Strehl (1984) — Laguerre configurations

Other authors often use partial permutations

An equivalent object with a different definition was introduced by Foata–Strehl (1984) — Laguerre configurations

Other authors often use partial permutations

Laguerre digraphs after Sokal (2022)

“History” of Foata–Zeilberger bijection

Start with all n vertices and no edges

“History” of Foata–Zeilberger bijection

Start with all n vertices and no edges

At each stage insert edges $i \rightarrow \sigma(i)$

“History” of Foata–Zeilberger bijection

Start with all n vertices and no edges

At each stage insert edges $i \rightarrow \sigma(i)$ in the following order:

Stage (a): $i \in H$ (fixed points) in increasing order

Stage (b): $i \in G$ (antiexcedances) in increasing order

Stage (c): $i \in F$ (excedances) in decreasing order

“History” of Foata–Zeilberger bijection

Start with all n vertices and no edges

At each stage insert edges $i \rightarrow \sigma(i)$ in the following order:

Stage (a): $i \in H$ (fixed points) in increasing order

Stage (b): $i \in G$ (antiexcedances) in increasing order

Stage (c): $i \in F$ (excedances) in decreasing order

This order is suggested by the inverse bijection and the inversion tables

History with an example

Let $\sigma = 715492638 = (1762)(3598)(4) \in \mathfrak{S}_9$.

$$H = \{4\}$$

$i \in F$	1	3	5
$\sigma(i) \in F'$	7	5	9
ξ_i	0	1	0

$i \in G$	2	6	7	8	9
$\sigma(i) \in G'$	1	2	6	3	8
ξ_i	0	0	1	0	0


History with an example

Let $\sigma = 715492638 = (1762)(3598)(4) \in \mathfrak{S}_9$.

$$H = \{4\}$$

$i \in F$	1	3	5
$\sigma(i) \in F'$	7	5	9
ξ_i	0	1	0

$i \in G$	2	6	7	8	9
$\sigma(i) \in G'$	1	2	6	3	8
ξ_i	0	0	1	0	0



History with an example


Let $\sigma = 715492638 = (1762)(3598)(4) \in \mathfrak{S}_9$.

$$H = \{4\}$$

$i \in F$	1	3	5
$\sigma(i) \in F'$	7	5	9
ξ_i	0	1	0

$i \in G$	2	6	7	8	9
$\sigma(i) \in G'$	1	2	6	3	8
ξ_i	0	0	1	0	0

Stage (a): H in increasing order



History with an example


Let $\sigma = 715492638 = (1762)(3598)(4) \in \mathfrak{S}_9$.

$$H = \{4\}$$

$i \in F$	1	3	5
$\sigma(i) \in F'$	7	5	9
ξ_i	0	1	0

$i \in G$	2	6	7	8	9
$\sigma(i) \in G'$	1	2	6	3	8
ξ_i	0	0	1	0	0

Stage (a): H in increasing order



History with an example


Let $\sigma = 715492638 = (1762)(3598)(4) \in \mathfrak{S}_9$.

$$H = \{4\}$$

$i \in F$	1	3	5
$\sigma(i) \in F'$	7	5	9
ξ_i	0	1	0

$i \in G$	2	6	7	8	9
$\sigma(i) \in G'$	1	2	6	3	8
ξ_i	0	0	1	0	0

Stage (b): G in increasing order



History with an example


Let $\sigma = 715492638 = (1762)(3598)(4) \in \mathfrak{S}_9$.

$$H = \{4\}$$

$i \in F$	1	3	5
$\sigma(i) \in F'$	7	5	9
ξ_i	0	1	0

$i \in G$	2	6	7	8	9
$\sigma(i) \in G'$	1	2	6	3	8
ξ_i	0	0	1	0	0

Stage (b): G in increasing order



History with an example


Let $\sigma = 715492638 = (1762)(3598)(4) \in \mathfrak{S}_9$.

$$H = \{4\}$$

$i \in F$	1	3	5
$\sigma(i) \in F'$	7	5	9
ξ_i	0	1	0

$i \in G$	2	6	7	8	9
$\sigma(i) \in G'$	1	2	6	3	8
ξ_i	0	0	1	0	0

Stage (b): G in increasing order



History with an example


Let $\sigma = 715492638 = (1762)(3598)(4) \in \mathfrak{S}_9$.

$$H = \{4\}$$

$i \in F$	1	3	5
$\sigma(i) \in F'$	7	5	9
ξ_i	0	1	0

$i \in G$	2	6	7	8	9
$\sigma(i) \in G'$	1	2	6	3	8
ξ_i	0	0	1	0	0

Stage (b): G in increasing order



History with an example


Let $\sigma = 715492638 = (1762)(3598)(4) \in \mathfrak{S}_9$.

$$H = \{4\}$$

$i \in F$	1	3	5
$\sigma(i) \in F'$	7	5	9
ξ_i	0	1	0

$i \in G$	2	6	7	8	9
$\sigma(i) \in G'$	1	2	6	3	8
ξ_i	0	0	1	0	0

Stage (b): G in increasing order



History with an example


Let $\sigma = 715492638 = (1762)(3598)(4) \in \mathfrak{S}_9$.

$$H = \{4\}$$

$i \in F$	1	3	5
$\sigma(i) \in F'$	7	5	9
ξ_i	0	1	0

$i \in G$	2	6	7	8	9
$\sigma(i) \in G'$	1	2	6	3	8
ξ_i	0	0	1	0	0

Stage (b): G in increasing order



History with an example


Let $\sigma = 715492638 = (1762)(3598)(4) \in \mathfrak{S}_9$.

$$H = \{4\}$$

$i \in F$	1	3	5
$\sigma(i) \in F'$	7	5	9
ξ_i	0	1	0

$i \in G$	2	6	7	8	9
$\sigma(i) \in G'$	1	2	6	3	8
ξ_i	0	0	1	0	0

Stage (c): F in decreasing order



History with an example


Let $\sigma = 715492638 = (1762)(3598)(4) \in \mathfrak{S}_9$.

$$H = \{4\}$$

$i \in F$	1	3	5
$\sigma(i) \in F'$	7	5	9
ξ_i	0	1	0

$i \in G$	2	6	7	8	9
$\sigma(i) \in G'$	1	2	6	3	8
ξ_i	0	0	1	0	0

Stage (c): F in decreasing order



History with an example


Let $\sigma = 715492638 = (1762)(3598)(4) \in \mathfrak{S}_9$.

$$H = \{4\}$$

$i \in F$	1	3	5
$\sigma(i) \in F'$	7	5	9
ξ_i	0	1	0

$i \in G$	2	6	7	8	9
$\sigma(i) \in G'$	1	2	6	3	8
ξ_i	0	0	1	0	0

Stage (c): F in decreasing order



History with an example


Let $\sigma = 715492638 = (1762)(3598)(4) \in \mathfrak{S}_9$.

$$H = \{4\}$$

$i \in F$	1	3	5
$\sigma(i) \in F'$	7	5	9
ξ_i	0	1	0

$i \in G$	2	6	7	8	9
$\sigma(i) \in G'$	1	2	6	3	8
ξ_i	0	0	1	0	0

Stage (c): F in decreasing order



- ① Continued fractions and enumerative combinatorics
 - ① Classical continued fractions
 - ② Sokal–Zeng’s results for factorials
 - ③ D.–Sokal’s results for Genocchi and median Genocchi numbers
 - ④ Conjectures
- ② Proof overview of existing results
 - ① Flajolet’s combinatorial interpretation
 - ② Foata–Zeilberger bijection
- ③ What’s new
 - ① Laguerre digraphs
 - ② New interpretation of the FZ bijection
- ④ The story continues ...

Flajolet's 1980 paper allowed continued fractions with non-commutative variables

Flajolet's 1980 paper allowed continued fractions with non-commutative variables

However, for this work, it is very important that the variables commute

Flajolet's 1980 paper allowed continued fractions with non-commutative variables

However, for this work, it is very important that the variables commute

So there must be non-commutative analogues of continued fractions coming from Biane bijection or Françon–Viennot bijection

A conjecture of Baril and Kirgizov

Baril and Kirgizov (2021) conjectured the following equidistribution of statistics on \mathfrak{S}_n :

Conjecture

The bistatistics (des_2, cyc) and (pex, cyc) are equidistributed.

A conjecture of Baril and Kirgizov

Baril and Kirgizov (2021) conjectured the following equidistribution of statistics on \mathfrak{S}_n :

Conjecture

The bistatistics (des_2, cyc) and (pex, cyc) are equidistributed.

Han–Mao–Zeng (2021) showed that this conjecture is equivalent to the following:

Conjecture

$$\sum_{n=0}^{\infty} y^{\text{des}_2 \sigma} \lambda^{\text{cyc} \sigma} = \frac{1}{1 - \lambda z - \frac{\lambda y z^2}{1 - (\lambda + 2)z - \frac{(\lambda + 1)(y + 1)z^2}{\dots}}}$$

A conjecture of Baril and Kirgizov

Baril and Kirgizov (2021) conjectured the following equidistribution of statistics on \mathfrak{S}_n :

Conjecture

The bistatistics (des_2, cyc) and (pex, cyc) are equidistributed.

Han–Mao–Zeng (2021) showed that this conjecture is equivalent to the following:

Conjecture

$$\sum_{n=0}^{\infty} y^{\text{des}_2 \sigma} \lambda^{\text{cyc} \sigma} = \frac{1}{1 - \lambda z - \frac{\lambda y z^2}{1 - (\lambda + 2)z - \frac{(\lambda + 1)(y + 1)z^2}{\dots}}}$$

How do we resolve this???

A conjecture of Baril and Kirgizov

Baril and Kirgizov (2021) conjectured the following equidistribution of statistics on \mathfrak{S}_n :

Conjecture

The bistatistics (des_2, cyc) and (pex, cyc) are equidistributed.

Han–Mao–Zeng (2021) showed that this conjecture is equivalent to the following:

Conjecture

$$\sum_{n=0}^{\infty} y^{\text{des}_2 \sigma} \lambda^{\text{cyc} \sigma} = \frac{1}{1 - \lambda z - \frac{\lambda y z^2}{1 - (\lambda + 2)z - \frac{(\lambda + 1)(y + 1)z^2}{\dots}}}$$

How do we resolve this???


An index i is a descent of type 2 if i is a descent and i is a record (left-to-right maxima)

Merci pour votre attention




Sokal–Zeng's master J-fraction

Pictorial representation







Pictorial representation



$$\sigma = 7192548610311121413 = \\ (1, 7, 8, 6, 4, 2) (3, 9, 10) (5) (11) (12) (13, 14) \in \mathfrak{S}_{14}.$$

Due to Corteel (2007)

Crossings, nestings and pseudo-nestings





$\text{ucross}(j, \sigma)$

$\text{lcross}(k, \sigma)$

$\text{unest}(j, \sigma)$

$\text{lnest}(k, \sigma)$



Sokal–Zeng first master J-fraction for permutations

$$Q_n(\mathbf{a}, \mathbf{b}, \mathbf{c}, \mathbf{d}, \mathbf{e}) =$$

$$\sum_{\sigma \in \mathfrak{S}_n} \prod_{i \in C_{\text{val}}(\sigma)} a_{\text{ucross}(i, \sigma), \text{unest}(i, \sigma)} \prod_{i \in C_{\text{peak}}(\sigma)} b_{\text{lcross}(i, \sigma), \text{lnest}(i, \sigma)} \times$$

$$\prod_{i \in C_{\text{dfall}}(\sigma)} c_{\text{lcross}(i, \sigma), \text{lnest}(i, \sigma)} \prod_{i \in C_{\text{drise}}(\sigma)} d_{\text{ucross}(i, \sigma), \text{unest}(i, \sigma)} \times$$

$$\prod_{i \in \text{Fix}(\sigma)} e_{\text{psnest}(i, \sigma)}$$

Sokal–Zeng first master J-fraction for permutations

$$Q_n(\mathbf{a}, \mathbf{b}, \mathbf{c}, \mathbf{d}, \mathbf{e}) =$$

$$\sum_{\sigma \in \mathfrak{S}_n} \prod_{i \in \text{Cval}(\sigma)} a_{\text{ucross}(i, \sigma), \text{unest}(i, \sigma)} \prod_{i \in \text{Cpeak}(\sigma)} b_{\text{lcross}(i, \sigma), \text{lneest}(i, \sigma)} \times \\ \prod_{i \in \text{Cdfall}(\sigma)} c_{\text{lcross}(i, \sigma), \text{lneest}(i, \sigma)} \prod_{i \in \text{Cdrise}(\sigma)} d_{\text{ucross}(i, \sigma), \text{unest}(i, \sigma)} \times \\ \prod_{i \in \text{Fix}(\sigma)} e_{\text{psnest}(i, \sigma)}$$

Theorem

$$\sum_{n=0}^{\infty} Q_n(\mathbf{a}, \mathbf{b}, \mathbf{c}, \mathbf{d}, \mathbf{e}) t^n = \frac{1}{1 - \gamma_0 t - \frac{\beta_1 t^2}{1 - \gamma_1 t - \frac{\beta_2 t^2}{1 - \gamma_2 t - \frac{\beta_3 t^2}{\ddots}}}}$$

Sokal–Zeng first master J-fraction for permutations

$$Q_n(\mathbf{a}, \mathbf{b}, \mathbf{c}, \mathbf{d}, \mathbf{e}) =$$

$$\sum_{\sigma \in \mathfrak{S}_n} \prod_{i \in C_{\text{val}}(\sigma)} a_{\text{ucross}(i, \sigma), \text{unest}(i, \sigma)} \prod_{i \in C_{\text{peak}}(\sigma)} b_{\text{lcross}(i, \sigma), \text{lneest}(i, \sigma)} \times \\ \prod_{i \in C_{\text{dfall}}(\sigma)} c_{\text{lcross}(i, \sigma), \text{lneest}(i, \sigma)} \prod_{i \in C_{\text{drise}}(\sigma)} d_{\text{ucross}(i, \sigma), \text{unest}(i, \sigma)} \times \\ \prod_{i \in \text{Fix}(\sigma)} e_{\text{psnest}(i, \sigma)}$$

Theorem

$$\sum_{n=0}^{\infty} Q_n(\mathbf{a}, \mathbf{b}, \mathbf{c}, \mathbf{d}, \mathbf{e}) t^n = \frac{1}{1 - \gamma_0 t - \frac{\beta_1 t^2}{1 - \gamma_1 t - \frac{\beta_2 t^2}{1 - \gamma_2 t - \frac{\beta_3 t^2}{\ddots}}}}$$

$$\gamma_k = \left(\sum_{\xi=0}^{k-1} c_{k-1-\xi, \xi} \right) + \left(\sum_{\xi=0}^{k-1} d_{k-1-\xi, \xi} \right) + e_k$$

$$\beta_k = \left(\sum_{\xi=0}^{k-1} a_{k-1-\xi, \xi} \right) \left(\sum_{\xi=0}^{k-1} b_{k-1-\xi, \xi} \right)$$

(D. arxiv '23) master J-fraction for permutations

$$Q_n(\mathbf{a}, \mathbf{b}, \mathbf{c}, \mathbf{d}, \mathbf{e}, \lambda) =$$

$$\sum_{\sigma \in \mathfrak{S}_n} \prod_{i \in \text{Cval}(\sigma)} a_{\text{ucross}(i, \sigma) + \text{unest}(i, \sigma)} \prod_{i \in \text{Cpeak}(\sigma)} b_{\text{lcross}(i, \sigma), \text{lnest}(i, \sigma)} \times$$

$$\prod_{i \in \text{Cdfall}(\sigma)} c_{\text{lcross}(i, \sigma), \text{lnest}(i, \sigma)} \prod_{i \in \text{Cdriase}(\sigma)} d_{\text{ucross}(i, \sigma), \text{unest}(i, \sigma)} \times$$

$$\prod_{i \in \text{Fix}(\sigma)} e_{\text{psnest}(i, \sigma)} \lambda^{\text{cyc}(\sigma)}$$

(D. arxiv '23) master J-fraction for permutations

$$Q_n(\mathbf{a}, \mathbf{b}, \mathbf{c}, \mathbf{d}, \mathbf{e}, \lambda) =$$

$$\sum_{\sigma \in \mathfrak{S}_n} \prod_{i \in C_{\text{val}}(\sigma)} a_{\text{ucross}(i, \sigma) + \text{unest}(i, \sigma)} \prod_{i \in C_{\text{peak}}(\sigma)} b_{\text{lcross}(i, \sigma), \text{lnest}(i, \sigma)} \times$$

$$\prod_{i \in C_{\text{dfall}}(\sigma)} c_{\text{lcross}(i, \sigma), \text{lnest}(i, \sigma)} \prod_{i \in C_{\text{drise}}(\sigma)} d_{\text{ucross}(i, \sigma), \text{unest}(i, \sigma)} \times$$

$$\prod_{i \in \text{Fix}(\sigma)} e_{\text{psnest}(i, \sigma)} \lambda^{\text{cyc}(\sigma)}$$

Theorem

$$\sum_{n=0}^{\infty} Q_n(\mathbf{a}, \mathbf{b}, \mathbf{c}, \mathbf{d}, \mathbf{e}, \lambda) t^n = \frac{1}{1 - \gamma_0 t - \frac{\beta_1 t^2}{1 - \gamma_1 t - \frac{\beta_2 t^2}{1 - \gamma_2 t - \frac{\beta_3 t^2}{\ddots}}}}$$

(D. arxiv '23) master J-fraction for permutations

$$Q_n(\mathbf{a}, \mathbf{b}, \mathbf{c}, \mathbf{d}, \mathbf{e}, \lambda) =$$

$$\sum_{\sigma \in \mathfrak{S}_n} \prod_{i \in \text{Cval}(\sigma)} a_{\text{ucross}(i, \sigma) + \text{unest}(i, \sigma)} \prod_{i \in \text{Cpeak}(\sigma)} b_{\text{lcross}(i, \sigma), \text{lnest}(i, \sigma)} \times$$

$$\prod_{i \in \text{Cdfall}(\sigma)} c_{\text{lcross}(i, \sigma), \text{lnest}(i, \sigma)} \prod_{i \in \text{Cdrise}(\sigma)} d_{\text{ucross}(i, \sigma), \text{unest}(i, \sigma)} \times$$

$$\prod_{i \in \text{Fix}(\sigma)} e_{\text{psnest}(i, \sigma)} \lambda^{\text{cyc}(\sigma)}$$

Theorem

$$\sum_{n=0}^{\infty} Q_n(\mathbf{a}, \mathbf{b}, \mathbf{c}, \mathbf{d}, \mathbf{e}, \lambda) t^n = \frac{1}{1 - \gamma_0 t - \frac{\beta_1 t^2}{1 - \gamma_1 t - \frac{\beta_2 t^2}{1 - \gamma_2 t - \frac{\beta_3 t^2}{\ddots}}}}$$

$$\gamma_k = \left(\sum_{\xi=0}^{k-1} c_{k-1-\xi, \xi} \right) + \left(\sum_{\xi=0}^{k-1} d_{k-1-\xi, \xi} \right) + \lambda e_k$$

$$\beta_k = (\lambda + k - 1) a_{k-1} \left(\sum_{\xi=0}^{k-1} b_{k-1-\xi, \xi} \right)$$