

# Total positivity of the Eulerian triangle: A big generalisation of Brenti's conjecture

Bishal Deb (he/him)

University College London

April 24, 2021

Graduate Student Combinatorics Conference (GSCC) 2021

*Based on Ongoing Joint Work With*

Xi Chen, Alexander Dyachenko, Tomack Gilmore and Alan D. Sokal

## Definition (Total Positivity (TP))

A matrix of real numbers said to be **totally positive (TP)** if all its minors are non-negative

## Definition (Total Positivity (TP))

A matrix of real numbers said to be **totally positive (TP)** if all its minors are non-negative i.e., determinants of all finite square submatrices are non-negative.

## Definition (Total Positivity (TP))

A matrix of real numbers said to be **totally positive (TP)** if all its minors are non-negative i.e., determinants of all finite square submatrices are non-negative.

Array of numbers and not linear operator.

## Definition (Total Positivity (TP))

A matrix of real numbers said to be **totally positive (TP)** if all its minors are non-negative i.e., determinants of all finite square submatrices are non-negative.

Array of numbers and not linear operator.

Need not be a square matrix,

## Definition (Total Positivity (TP))

A matrix of real numbers said to be **totally positive (TP)** if all its minors are non-negative i.e., determinants of all finite square submatrices are non-negative.

Array of numbers and not linear operator.

Need not be a square matrix, or finite!

## Definition (Total Positivity (TP))

A matrix of real numbers said to be **totally positive (TP)** if all its minors are non-negative i.e., determinants of all finite square submatrices are non-negative.

Array of numbers and not linear operator.

Need not be a square matrix, or finite!


Will consider matrix of polynomials soon!

# Historical Note

First defined independently by two different groups in the 30s



(a) M.G. Krein (1907-1989)



(b) I.G. Schoenberg  
(1903-1990)

Source: MacTutor History of Mathematics Archive

We use Schoenberg's terminology.



# Example and basic properties

# Example and basic properties

- Example: Bidiagonal matrices with entries  $\geq 0$

# Example and basic properties

- Example: Bidiagonal matrices with entries  $\geq 0$
- Matrix sum doesn't preserve TP

# Example and basic properties

- Example: Bidiagonal matrices with entries  $\geq 0$
- Matrix sum doesn't preserve TP (Has  $2 \times 2$  example)

# Example and basic properties

- Example: Bidiagonal matrices with entries  $\geq 0$
- Matrix sum doesn't preserve TP (Has  $2 \times 2$  example)
- Matrix product preserves TP

# Example and basic properties

- Example: Bidiagonal matrices with entries  $\geq 0$
- Matrix sum doesn't preserve TP (Has  $2 \times 2$  example)
- Matrix product preserves TP (Cauchy-Binet formula)

# Applications to Combinatorics

- Brenti in 80s started studying TP in combinatorics.



# Applications to Combinatorics

- Brenti in 80s started studying TP in combinatorics.
- Many important combinatorial triangles are TP.

- Brenti in 80s started studying TP in combinatorics.
- Many important combinatorial triangles are TP.

Examples

- Brenti in 80s started studying TP in combinatorics.
- Many important combinatorial triangles are TP.

## Examples

- Binomial triangle. Entries  $\binom{n}{k}$

- Brenti in 80s started studying TP in combinatorics.
- Many important combinatorial triangles are TP.

## Examples

- Binomial triangle. Entries  $\binom{n}{k}$
- Stirling cycle triangle. Entries  $\left[ \begin{smallmatrix} n+1 \\ k \end{smallmatrix} \right]$

- Brenti in 80s started studying TP in combinatorics.
- Many important combinatorial triangles are TP.

## Examples

- Binomial triangle. Entries  $\binom{n}{k}$
- Stirling cycle triangle. Entries  $\left[ \begin{smallmatrix} n+1 \\ k \end{smallmatrix} \right]$
- Stirling subset triangle. Entries  $\left\{ \begin{smallmatrix} n+1 \\ k \end{smallmatrix} \right\}$

- Brenti in 80s started studying TP in combinatorics.
- Many important combinatorial triangles are TP.

## Examples

- Binomial triangle. Entries  $\binom{n}{k}$
- Stirling cycle triangle. Entries  $\left[ \begin{smallmatrix} n+1 \\ k \end{smallmatrix} \right]$
- Stirling subset triangle. Entries  $\left\{ \begin{smallmatrix} n+1 \\ k \end{smallmatrix} \right\}$
- Reversed Stirling cycle triangle. Entries  $\left[ \begin{smallmatrix} n+1 \\ n-k+1 \end{smallmatrix} \right]$

- Brenti in 80s started studying TP in combinatorics.
- Many important combinatorial triangles are TP.

## Examples

- Binomial triangle. Entries  $\binom{n}{k}$
- Stirling cycle triangle. Entries  $\left[ \begin{smallmatrix} n+1 \\ k \end{smallmatrix} \right]$
- Stirling subset triangle. Entries  $\left\{ \begin{smallmatrix} n+1 \\ k \end{smallmatrix} \right\}$
- Reversed Stirling cycle triangle. Entries  $\left[ \begin{smallmatrix} n+1 \\ n-k+1 \end{smallmatrix} \right]$   
(reversal does not preserve TP)

# Reversed Stirling Subset Triangle

Reversed Stirling subset triangle:

$$\left( \left\{ \begin{matrix} n+1 \\ k \end{matrix} \right\}^{\text{rev}} \right)_{n,k \geq 0} = \left( \left\{ \begin{matrix} n+1 \\ n-k+1 \end{matrix} \right\} \right)_{n,k \geq 0} = \begin{bmatrix} 1 & & & & & \\ 1 & 1 & & & & \\ 1 & 3 & 1 & & & \\ 1 & 6 & 7 & 1 & & \\ 1 & 10 & 25 & 15 & 1 & \\ \vdots & \vdots & \vdots & \vdots & \vdots & \ddots \end{bmatrix}$$



# Reversed Stirling Subset Triangle

Reversed Stirling subset triangle:

$$\left( \left\{ \begin{matrix} n+1 \\ k \end{matrix} \right\}^{\text{rev}} \right)_{n,k \geq 0} = \left( \left\{ \begin{matrix} n+1 \\ n-k+1 \end{matrix} \right\} \right)_{n,k \geq 0} = \begin{bmatrix} 1 & & & & & \\ 1 & 1 & & & & \\ 1 & 3 & 1 & & & \\ 1 & 6 & 7 & 1 & & \\ 1 & 10 & 25 & 15 & 1 & \\ \vdots & \vdots & \vdots & \vdots & \vdots & \ddots \end{bmatrix}$$

Conjecture (Us 2019)

*The infinite lower triangular matrix  $\left( \left\{ \begin{matrix} n+1 \\ k \end{matrix} \right\}^{\text{rev}} \right)_{n,k \geq 0} = \left( \left\{ \begin{matrix} n+1 \\ n-k+1 \end{matrix} \right\} \right)_{n,k \geq 0}$  is TP.*

# Eulerian Numbers

Descent of a permutation  $\sigma$ :

Descent of a permutation  $\sigma$ : Position  $i$  such that  $\sigma(i) > \sigma(i+1)$ .

# Eulerian Numbers

Descent of a permutation  $\sigma$ : Position  $i$  such that  $\sigma(i) > \sigma(i+1)$ .

$\text{des}(\sigma)$ : number of descents of  $\sigma$ .

# Eulerian Numbers

Descent of a permutation  $\sigma$ : Position  $i$  such that  $\sigma(i) > \sigma(i+1)$ .

$\text{des}(\sigma)$ : number of descents of  $\sigma$ .

Example: For  $\sigma = 943127685$ ,  $\text{des}(\sigma) = 5$

Descent of a permutation  $\sigma$ : Position  $i$  such that  $\sigma(i) > \sigma(i+1)$ .

$\text{des}(\sigma)$ : number of descents of  $\sigma$ .

Example: For  $\sigma = 943127685$ ,  $\text{des}(\sigma) = 5$

## Definition

The Eulerian number  $\left\langle n \atop k \right\rangle$  is defined to be the cardinality of the set  $\{\sigma \in \mathfrak{S}_n \mid \text{des}(\sigma) = k\}$ .

## Definition

The Eulerian number  $\left\langle n \atop k \right\rangle$  is defined to be the cardinality of the set  $\{\sigma \in \mathfrak{S}_n \mid \text{des}(\sigma) = k\}$ .



## Definition

The Eulerian number  $\langle n \rangle_k$  is defined to be the cardinality of the set  $\{\sigma \in \mathfrak{S}_n \mid \text{des}(\sigma) = k\}$ .

$$\left( \left\langle \begin{matrix} n+1 \\ k \end{matrix} \right\rangle \right)_{n,k \geq 0} = \begin{bmatrix} 1 & & & & & & \\ 1 & 1 & & & & & \\ 1 & 4 & 1 & & & & \\ 1 & 11 & 11 & 1 & & & \\ 1 & 26 & 66 & 26 & 1 & & \\ 1 & 57 & 302 & 302 & 57 & 1 & \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \ddots \end{bmatrix}$$

# Eulerian Triangle Conjecture

$$\left( \left\langle \begin{matrix} n+1 \\ k \end{matrix} \right\rangle \right)_{n,k \geq 0} = \begin{bmatrix} 1 & & & & & & \\ 1 & 1 & & & & & \\ 1 & 4 & 1 & & & & \\ 1 & 11 & 11 & 1 & & & \\ 1 & 26 & 66 & 26 & 1 & & \\ 1 & 57 & 302 & 302 & 57 & 1 & \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \ddots \end{bmatrix}$$

Conjecture (Brenti 1996)

*The infinite lower triangular matrix  $\left( \left\langle \begin{matrix} n+1 \\ k \end{matrix} \right\rangle \right)_{n,k \geq 0}$  is TP.*

# Eulerian Triangle Conjecture

$$\left( \left\langle \begin{matrix} n+1 \\ k \end{matrix} \right\rangle \right)_{n,k \geq 0} = \begin{bmatrix} 1 & & & & & & \\ 1 & 1 & & & & & \\ 1 & 4 & 1 & & & & \\ 1 & 11 & 11 & 1 & & & \\ 1 & 26 & 66 & 26 & 1 & & \\ 1 & 57 & 302 & 302 & 57 & 1 & \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \ddots \end{bmatrix}$$

Conjecture (Brenti 1996)

*The infinite lower triangular matrix  $\left( \left\langle \begin{matrix} n+1 \\ k \end{matrix} \right\rangle \right)_{n,k \geq 0}$  is TP.*

Verified for  $512 \times 512$  using an efficient bidiagonal factorisation algorithm.

# Eulerian Triangle Conjecture

$$\left( \left\langle \begin{matrix} n+1 \\ k \end{matrix} \right\rangle \right)_{n,k \geq 0} = \begin{bmatrix} 1 & & & & & & \\ 1 & 1 & & & & & \\ 1 & 4 & 1 & & & & \\ 1 & 11 & 11 & 1 & & & \\ 1 & 26 & 66 & 26 & 1 & & \\ 1 & 57 & 302 & 302 & 57 & 1 & \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \ddots \end{bmatrix}$$

Conjecture (Brenti 1996)

*The infinite lower triangular matrix  $\left( \left\langle \begin{matrix} n+1 \\ k \end{matrix} \right\rangle \right)_{n,k \geq 0}$  is TP.*

Verified for  $512 \times 512$  using an efficient bidiagonal factorisation algorithm.

Rational numbers with large denominators appear.

# Eulerian Triangle Conjecture

$$\left( \left\langle \begin{matrix} n+1 \\ k \end{matrix} \right\rangle \right)_{n,k \geq 0} = \begin{bmatrix} 1 & & & & & & \\ 1 & 1 & & & & & \\ 1 & 4 & 1 & & & & \\ 1 & 11 & 11 & 1 & & & \\ 1 & 26 & 66 & 26 & 1 & & \\ 1 & 57 & 302 & 302 & 57 & 1 & \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \ddots \end{bmatrix}$$

Conjecture (Brenti 1996)

*The infinite lower triangular matrix  $\left( \left\langle \begin{matrix} n+1 \\ k \end{matrix} \right\rangle \right)_{n,k \geq 0}$  is TP.*

Verified for  $512 \times 512$  using an efficient bidiagonal factorisation algorithm.

Rational numbers with large denominators appear.

General pattern not clear.

# A comparison of the two triangles

Reversed Stirling subset triangle:

$$\left( \left\{ \begin{matrix} n+1 \\ k \end{matrix} \right\}^{\text{rev}} \right)_{n,k \geq 0} = \begin{bmatrix} 1 & & & & & \\ 1 & 1 & & & & \\ 1 & 3 & 1 & & & \\ 1 & 6 & 7 & 1 & & \\ 1 & 10 & 25 & 15 & 1 & \\ \vdots & \vdots & \vdots & \vdots & \vdots & \ddots \end{bmatrix}$$

Eulerian triangle:

$$\left( \left\langle \begin{matrix} n+1 \\ k \end{matrix} \right\rangle \right)_{n,k \geq 0} = \begin{bmatrix} 1 & & & & & \\ 1 & 1 & & & & \\ 1 & 4 & 1 & & & \\ 1 & 11 & 11 & 1 & & \\ 1 & 26 & 66 & 26 & 1 & \\ \vdots & \vdots & \vdots & \vdots & \vdots & \ddots \end{bmatrix}$$

# A comparison of the two triangles

Reversed Stirling subset triangle:

$$\left( \left\{ \begin{matrix} n+1 \\ k \end{matrix} \right\}^{\text{rev}} \right)_{n,k \geq 0} = \begin{bmatrix} 1 & & & & & \\ 1 & 1 & & & & \\ 1 & 3 & 1 & & & \\ 1 & 6 & 7 & 1 & & \\ 1 & 10 & 25 & 15 & 1 & \\ \vdots & \vdots & \vdots & \vdots & \vdots & \ddots \end{bmatrix}$$

$$A_{n,k} = (n - k + 1)A_{n-1,k-1} + 1 \cdot A_{n-1,k}$$

Eulerian triangle:

$$\left( \left\langle \begin{matrix} n+1 \\ k \end{matrix} \right\rangle \right)_{n,k \geq 0} = \begin{bmatrix} 1 & & & & & \\ 1 & 1 & & & & \\ 1 & 4 & 1 & & & \\ 1 & 11 & 11 & 1 & & \\ 1 & 26 & 66 & 26 & 1 & \\ \vdots & \vdots & \vdots & \vdots & \vdots & \ddots \end{bmatrix}$$

$$A_{n,k} = (n - k + 1)A_{n-1,k-1} + (k + 1)A_{n-1,k}$$

# A comparison of the two triangles

Reversed Stirling subset triangle:

$$\left( \left\{ \begin{matrix} n+1 \\ k \end{matrix} \right\}^{\text{rev}} \right)_{n,k \geq 0} = \begin{bmatrix} 1 & & & & & \\ 1 & 1 & & & & \\ 1 & 3 & 1 & & & \\ 1 & 6 & 7 & 1 & & \\ 1 & 10 & 25 & 15 & 1 & \\ \vdots & \vdots & \vdots & \vdots & \vdots & \ddots \end{bmatrix}$$

$$A_{n,k} = (n - k + 1) A_{n-1,k-1} + 1 \cdot A_{n-1,k}$$

Introduce  
variables

Eulerian triangle:

$$\left( \left\langle \begin{matrix} n+1 \\ k \end{matrix} \right\rangle \right)_{n,k \geq 0} = \begin{bmatrix} 1 & & & & & \\ 1 & 1 & & & & \\ 1 & 4 & 1 & & & \\ 1 & 11 & 11 & 1 & & \\ 1 & 26 & 66 & 26 & 1 & \\ \vdots & \vdots & \vdots & \vdots & \vdots & \ddots \end{bmatrix}$$

$$A_{n,k} = (n - k + 1) A_{n-1,k-1} + (k + 1) A_{n-1,k}$$



## Definition (Coefficientwise TP)

A matrix of polynomials with real coefficients is said to be **coefficientwise totally positive (coefficientwise TP)** if all its minors have non-negative coefficients.

## Definition (Coefficientwise TP)

A matrix of polynomials with real coefficients is said to be **coefficientwise totally positive (coefficientwise TP)** if all its minors have non-negative coefficients.

One or several variables.

## Definition (Coefficientwise TP)

A matrix of polynomials with real coefficients is said to be **coefficientwise totally positive (coefficientwise TP)** if all its minors have non-negative coefficients.

One or several variables.

Coefficientwise TP  $\implies$  Pointwise TP.

## Definition (Coefficientwise TP)

A matrix of polynomials with real coefficients is said to be **coefficientwise totally positive (coefficientwise TP)** if all its minors have non-negative coefficients.

One or several variables.

Coefficientwise TP  $\implies$  Pointwise TP.

But much stronger.

# Putting variables in the recurrence

Replace

$$A_{n,k} = (n - k + 1)A_{n-1,k-1} + (k + 1)A_{n-1,k}$$

$$A_{n,k} = (n - k + 1)A_{n-1,k-1} + 1 \cdot A_{n-1,k}$$

Introduce variables  $a, b, c, d, e$ .

$$A_{n,k} = [a(n - k) + b(k - 1) + c]A_{n-1,k-1} + (dk + e)A_{n-1,k}$$

# Putting variables in the recurrence

Replace

$$A_{n,k} = (n - k + 1)A_{n-1,k-1} + (k + 1)A_{n-1,k}$$

$$A_{n,k} = (n - k + 1)A_{n-1,k-1} + 1 \cdot A_{n-1,k}$$

Introduce variables  $a, b, c, d, e$ .

$$A_{n,k} = [a(n - k) + b(k - 1) + c]A_{n-1,k-1} + (dk + e)A_{n-1,k}$$

Is it coefficientwise TP?

(Minors are polynomials with non-negative coefficients?)

# Putting variables in the recurrence

Replace

$$A_{n,k} = (n - k + 1)A_{n-1,k-1} + (k + 1)A_{n-1,k}$$

$$A_{n,k} = (n - k + 1)A_{n-1,k-1} + 1 \cdot A_{n-1,k}$$

Introduce variables  $a, b, c, d, e$ .

$$A_{n,k} = [a(n - k) + b(k - 1) + c]A_{n-1,k-1} + (dk + e)A_{n-1,k}$$

Is it coefficientwise TP?

(Minors are polynomials with non-negative coefficients?)

More general conjecture!

# Putting variables in the recurrence

Replace

$$A_{n,k} = (n - k + 1)A_{n-1,k-1} + (k + 1)A_{n-1,k}$$

$$A_{n,k} = (n - k + 1)A_{n-1,k-1} + 1 \cdot A_{n-1,k}$$

Introduce variables  $a, b, c, d, e$ .

$$A_{n,k} = [a(n - k) + b(k - 1) + c]A_{n-1,k-1} + (dk + e)A_{n-1,k}$$

Is it coefficientwise TP?

(Minors are polynomials with non-negative coefficients?)

More general conjecture!

True till  $11 \times 11$  - a week.



# Putting variables in the recurrence

Replace

$$A_{n,k} = (n - k + 1)A_{n-1,k-1} + (k + 1)A_{n-1,k}$$

$$A_{n,k} = (n - k + 1)A_{n-1,k-1} + 1 \cdot A_{n-1,k}$$

Introduce variables  $a, b, c, d, e$ .

$$A_{n,k} = [a(n - k) + b(k - 1) + c]A_{n-1,k-1} + (dk + e)A_{n-1,k}$$

Is it coefficientwise TP?

(Minors are polynomials with non-negative coefficients?)

More general conjecture!

True till  $11 \times 11$  - a week.

False for  $12 \times 12$  - 99 days CPU time.

# Putting variables in the recurrence

Replace

$$A_{n,k} = (n - k + 1)A_{n-1,k-1} + (k + 1)A_{n-1,k}$$

$$A_{n,k} = (n - k + 1)A_{n-1,k-1} + 1 \cdot A_{n-1,k}$$

Introduce variables  $a, b, c, d, e$ .

$$A_{n,k} = [a(n - k) + b(k - 1) + c]A_{n-1,k-1} + (dk + e)A_{n-1,k}$$

Is it coefficientwise TP?

(Minors are polynomials with non-negative coefficients?)

More general conjecture!

True till  $11 \times 11$  - a week.

False for  $12 \times 12$  - 99 days CPU time. ☹

# Is all hope lost?

Put  $b = 0$  to obtain

$$A_{n,k} = [a(n-k) + c]A_{n-1,k-1} + (dk + e)A_{n-1,k}$$

# Is all hope lost?

Put  $b = 0$  to obtain

$$A_{n,k} = [a(n-k) + c]A_{n-1,k-1} + (dk + e)A_{n-1,k}$$

Empirically true upto  $13 \times 13$ .

# Is all hope lost?

Put  $b = 0$  to obtain

$$A_{n,k} = [a(n-k) + c]A_{n-1,k-1} + (dk + e)A_{n-1,k}$$

Empirically true upto  $13 \times 13$ .

Special cases:

$(a, c, d, e)$	Matrix obtained
----------------	-----------------

# Is all hope lost?

Put  $b = 0$  to obtain

$$A_{n,k} = [a(n-k) + c]A_{n-1,k-1} + (dk + e)A_{n-1,k}$$

Empirically true upto  $13 \times 13$ .

Special cases:

$(a, c, d, e)$	Matrix obtained
$(1, 1, 1, 1)$	clean Eulerian triangle, conjecture
$(1, 0, 1, 1)$	shifted Eulerian triangle, conjecture
$(1, 1, 1, 0)$	shifted Eulerian triangle, conjecture

# Is all hope lost?

Put  $b = 0$  to obtain

$$A_{n,k} = [a(n-k) + c]A_{n-1,k-1} + (dk + e)A_{n-1,k}$$

Empirically true upto  $13 \times 13$ .

Special cases:

$(a, c, d, e)$	Matrix obtained
$(1, 1, 1, 1)$	clean Eulerian triangle, conjecture
$(1, 0, 1, 1)$	shifted Eulerian triangle, conjecture
$(1, 1, 1, 0)$	shifted Eulerian triangle, conjecture
$(0, 1, 0, 1)$	Binomial triangle, TP

# Is all hope lost?

Put  $b = 0$  to obtain

$$A_{n,k} = [a(n-k) + c]A_{n-1,k-1} + (dk + e)A_{n-1,k}$$

Empirically true upto  $13 \times 13$ .

Special cases:

$(a, c, d, e)$	Matrix obtained
$(1, 1, 1, 1)$	clean Eulerian triangle, conjecture
$(1, 0, 1, 1)$	shifted Eulerian triangle, conjecture
$(1, 1, 1, 0)$	shifted Eulerian triangle, conjecture
$(0, 1, 0, 1)$	Binomial triangle, TP
$(0, 1, 1, 1)$	Stirling subset numbers, TP proved by Brenti 1995



# Is all hope lost?


Put  $b = 0$  to obtain

$$A_{n,k} = [a(n-k) + c]A_{n-1,k-1} + (dk + e)A_{n-1,k}$$

Empirically true upto  $13 \times 13$ .

Special cases:

$(a, c, d, e)$	Matrix obtained
$(1, 1, 1, 1)$	clean Eulerian triangle, conjecture
$(1, 0, 1, 1)$	shifted Eulerian triangle, conjecture
$(1, 1, 1, 0)$	shifted Eulerian triangle, conjecture
$(0, 1, 0, 1)$	Binomial triangle, TP
$(0, 1, 1, 1)$	Stirling subset numbers, TP proved by Brenti 1995
$(1, 1, 0, 1)$	Row reversed matrix of Stirling subset numbers, conjecture



$$A_{n,k} = (n - k + 1)A_{n-1,k-1}$$

$$+ (k + 1)A_{n-1,k}$$

$$A_{n,k} = (n - k + 1)A_{n-1,k-1} + 1 \cdot A_{n-1,k}$$



$$A_{n,k} = [a(n - k) + c]A_{n-1,k-1}$$

$$+ (dk + e)A_{n-1,k}$$

# Is all hope lost?

Put  $b = 0$  to obtain

$$A_{n,k} = [a(n-k) + c]A_{n-1,k-1} + (dk + e)A_{n-1,k}$$

This seems to be coefficientwise TP so far.

In fact, we have the following special cases for different substitutions:

$(a, c, d, e)$	Matrix obtained
$(1, 1, 1, 1)$	clean Eulerian triangle, conjecture
$(1, 0, 1, 1)$	Down-shifted Eulerian triangle, conjecture
$(1, 1, 1, 0)$	Diagonally down-shifted Eulerian triangle, conjecture
$(0, 1, 0, 1)$	Binomial triangle, TP
$(0, 1, 1, 1)$	Stirling subset numbers, TP proved by Brenti 1995
$(1, 1, 0, 1)$	<b>Row reversed matrix of Stirling subset numbers, conjecture</b>

# Is all hope lost?

Put  $b = 0$  to obtain

$$A_{n,k} = [a(n-k) + c]A_{n-1,k-1} + (dk + e)A_{n-1,k}$$

This seems to be coefficientwise TP so far.

In fact, we have the following special cases for different substitutions:

$(a, c, d, e)$	Matrix obtained
$(1, 1, 1, 1)$	clean Eulerian triangle, conjecture
$(1, 0, 1, 1)$	Down-shifted Eulerian triangle, conjecture
$(1, 1, 1, 0)$	Diagonally down-shifted Eulerian triangle, conjecture
$(0, 1, 0, 1)$	Binomial triangle, TP
$(0, 1, 1, 1)$	Stirling subset numbers, TP proved by Brenti 1995
$(1, 1, 0, 1)$	<b>Row reversed matrix of Stirling subset numbers, conjecture</b>

## Reversed Stirling subset triangle

$$\left( \left\{ \begin{matrix} n+1 \\ k \end{matrix} \right\}^{\text{rev}} \right)_{n,k \geq 0} = \begin{bmatrix} 1 & & & & & \\ 1 & 1 & & & & \\ 1 & 3 & 1 & & & \\ 1 & 6 & 7 & 1 & & \\ 1 & 10 & 25 & 15 & 1 & \\ \vdots & \vdots & \vdots & \vdots & \vdots & \ddots \end{bmatrix}.$$

ace triangle

$$\begin{bmatrix} 1 & & & & & \\ e & c & & & & \\ e^2 & ae + 2ce & c^2 & & & \\ e^3 & 3ae^2 + 3ce^2 & a^2e + 3ace + 3c^2e & c^3 & & \\ e^4 & 6ae^3 + 4ce^3 & 7a^2e^2 + 12ace^2 + 6c^2e^2 & a^3e + 4a^2ce + 6ac^2e + 4c^3e & c^4 & \\ \vdots & \vdots & \vdots & \vdots & \vdots & \ddots \end{bmatrix}.$$

recurrence

$$A_{n,k} = [\textcolor{green}{a}(n-k) + \textcolor{red}{c}]A_{n-1,k-1} + \textcolor{blue}{e}A_{n-1,k}$$

ace triangle

$$\begin{bmatrix} 1 & & & & & \\ e & c & & & & \\ e^2 & ae + 2ce & c^2 & & & \\ e^3 & 3ae^2 + 3ce^2 & a^2e + 3ace + 3c^2e & c^3 & & \\ e^4 & 6ae^3 + 4ce^3 & 7a^2e^2 + 12ace^2 + 6c^2e^2 & a^3e + 4a^2ce + 6ac^2e + 4c^3e & c^4 & \\ \vdots & \vdots & \vdots & \vdots & \vdots & \ddots \end{bmatrix}.$$


recurrence

$$A_{n,k} = [\textcolor{green}{a}(n-k) + \textcolor{red}{c}]A_{n-1,k-1} + \textcolor{blue}{e}A_{n-1,k}$$

We have three proofs:

- Two Digraph proofs using the Lindström-Gessel-Viennot lemma
  - Bijection between paths in the digraph to set partitions.
  - Exploiting the structure of the digraph.
- Algebraic proof

# An *ace* digraph



—  
1


—  
*a*

—  
*c*

—  
*e*



# An *ace* digraph



We have a weight preserving bijection between paths from  $u_n \rightarrow v_k$  and set partitions of  $\{1, \dots, n+1\}$  into  $n-k+1$  parts.

$$A_{n,k} = (n - k + 1)A_{n-1,k-1} \\ + (k + 1)A_{n-1,k}$$

Eulerian triangle conjecture

$$A_{n,k} = [a(n - k) + c]A_{n-1,k-1} \\ + (dk + e)A_{n-1,k}$$



*acde* conjecture

$$A_{n,k} = [a(n - k) + c]A_{n-1,k-1} \\ + (dk + e)A_{n-1,k} \\ + [f(n - 2) + g]A_{n-2,k-1}$$

*acdefg* conjecture

$$A_{n,k} = [a(n - k) + c]A_{n-1,k-1} \\ + \left( \sum_{i=0}^k d_i \right) A_{n-1,k} \\ + [f(n - 2) + g]A_{n-2,k-1}$$

*acd<sup>d</sup>fg* conjecture



10/27/20

J-fraction for new generating polynomials of TTSa with  $g=f$

$$\delta_n = [c + \binom{n}{2}d] + [c + \binom{n}{2}a]x$$

$$\beta_n = n[cd + ae + f + (n-1)ad]x$$

see conf from only action "Eulerian polynomials ..."

Stokel-Zeng have first J-fraction for permutations

$$\delta_0 = w_0$$

$$\delta_n = [x_2 + (n-1)u_2] + [y_2 + (n-1)v_2] + w_n \text{ for } n \geq 1$$

$$\beta_n = [x_1 + (n-1)u_1][y_1 + (n-1)v_1]$$

so we could take ...

# Thank you

Meme images from internet and digraph drawn by Tomack Gilmore.