

Total Positivity during distressing times

Bishal Deb

University College London

November 17, 2020


Supervisor

Alan D. Sokal

Ugh. Did you spend
all afternoon making
memes again?



Hey, meme creation is
the combinatorics
of culture.



Source: Math with Bad Drawings

- ① Introduction
- ② Proof techniques and some special types of matrices
 - ① LGV lemma
 - ② Hankel matrices
 - ③ Toeplitz matrices
 - ④ Lower triangular matrices
- ③ The Eulerian triangle

- ➊ Introduction
- ➋ Proof techniques and some special types of matrices
 - ➊ LGV lemma
 - ➋ Hankel matrices
 - ➌ Toeplitz matrices
 - ➍ Lower triangular matrices
- ➌ The Eulerian triangle

Horoscope:

Something positive
will happen to you
this month!



Definition (TP and STP)

We say that a matrix (not necessarily finite) is *totally positive (TP)* if all its minors (determinants of finite square submatrices) are non-negative. It is said to be *strictly totally positive (STP)* if instead of non-negativity we want positivity of the minors.

We say that a matrix is totally positive of order r (TP_r) if all its submatrices of size $\leq r$ are non-negative.

Definition (TP and STP)

We say that a matrix (not necessarily finite) is *totally positive (TP)* if all its minors (determinants of finite square submatrices) are non-negative. It is said to be *strictly totally positive (STP)* if instead of non-negativity we want positivity of the minors.

We say that a matrix is totally positive of order r (TP_r) if all its submatrices of size $\leq r$ are non-negative.

In this talk we shall only discuss TP.

Do such matrices exist?

Do such matrices exist?

The answer to the above question is ofcourse true as we have 0 matrices.

$$\begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$$

Do such matrices exist?

Are there any others?

Do such matrices exist?

Are there any others?

Yes, matrices whose entries are all equal to a non-negative constant.

$$\begin{bmatrix} c & c & c \\ c & c & c \end{bmatrix}$$

where $c \geq 0$.

Do such matrices exist?

Any non-constant matrices?

Do such matrices exist?

Any non-constant matrices?

Yes, diagonal matrices with non-negative entries on the diagonal.

$$\begin{bmatrix} d_1 & 0 & 0 & 0 \\ 0 & d_2 & 0 & 0 \\ 0 & 0 & d_3 & 0 \\ 0 & 0 & 0 & d_4 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

where $d_i \geq 0$.

Do such matrices exist?

Anything else?

Do such matrices exist?

Anything else?

Yes, bidiagonal matrices.

$$\begin{bmatrix} d_1 & 0 & 0 & 0 \\ r_1 & d_2 & 0 & 0 \\ 0 & r_2 & d_3 & 0 \\ 0 & 0 & r_3 & d_4 \\ 0 & 0 & 0 & r_4 \end{bmatrix}$$

or

$$\begin{bmatrix} d_1 & r_1 & 0 & 0 \\ 0 & d_2 & r_2 & 0 \\ 0 & 0 & d_3 & r_3 \\ 0 & 0 & 0 & d_4 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

where $d_i \geq 0$ and $r_i \geq 0$.

Do such matrices exist?

Anything else?

Yes, bidiagonal matrices.

$$\begin{bmatrix} d_1 & 0 & 0 & 0 \\ r_1 & d_2 & 0 & 0 \\ 0 & r_2 & d_3 & 0 \\ 0 & 0 & r_3 & d_4 \\ 0 & 0 & 0 & r_4 \end{bmatrix}$$

or

$$\begin{bmatrix} d_1 & r_1 & 0 & 0 \\ 0 & d_2 & r_2 & 0 \\ 0 & 0 & d_3 & r_3 \\ 0 & 0 & 0 & d_4 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

where $d_i \geq 0$ and $r_i \geq 0$.

This is because any square submatrix is either upper triangular or lower triangular and hence the determinant can be computed by looking at the diagonal entries.

Do such matrices exist?

Anything else?

Yes, bidiagonal matrices.

$$\begin{bmatrix} d_1 & 0 & 0 & 0 \\ r_1 & d_2 & 0 & 0 \\ 0 & r_2 & d_3 & 0 \\ 0 & 0 & r_3 & d_4 \\ 0 & 0 & 0 & r_4 \end{bmatrix} \quad \text{or} \quad \begin{bmatrix} d_1 & r_1 & 0 & 0 \\ 0 & d_2 & r_2 & 0 \\ 0 & 0 & d_3 & r_3 \\ 0 & 0 & 0 & d_4 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

where $d_i \geq 0$ and $r_i \geq 0$.

This is because any square submatrix is either upper triangular or lower triangular and hence the determinant can be computed by looking at the diagonal entries.

Can we construct other examples?

Some Operations which Preserve TP

- Taking submatrices,

Some Operations which Preserve TP

- Taking submatrices, adding rows or columns of zeros,

Some Operations which Preserve TP

- Taking submatrices, adding rows or columns of zeros, inserting copies of row or columns next to them.

Some Operations which Preserve TP

- Taking submatrices, adding rows or columns of zeros, inserting copies of row or columns next to them.
- Block matrices of the form

$$\left[\begin{array}{c|c} A & 0 \\ \hline 0 & B \end{array} \right]$$

where both A and B are TP.

Some Operations which Preserve TP

- Taking submatrices, adding rows or columns of zeros, inserting copies of row or columns next to them.
- Block matrices of the form

$$\left[\begin{array}{c|c} A & 0 \\ \hline 0 & B \end{array} \right]$$

where both A and B are TP.

- Matrix Products preserve TP.

Proof: Use Cauchy-Binet formula which we recall here:

Theorem (Cauchy-Binet formula)

Let A be an $m \times n$ matrix and B an $n \times m$ matrix. Then we have that

$$\det(AB) = \sum_{S \in \binom{[n]}{m}} \det(A_{[m],S}) \det(B_{S,[m]})$$

where $[n]$ denotes the set $\{1, \dots, n\}$, and given a set T , $\binom{T}{k}$ is the collection of all k -dimensional subsets of T .

Matrix Sum doesn't generally preserve TP

$$\begin{bmatrix} 1 & 2 \\ 1 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 1 & 1 \end{bmatrix} + \begin{bmatrix} 1 & 2 \\ 0 & 1 \end{bmatrix}$$

where

$$\det \left(\begin{bmatrix} 1 & 2 \\ 1 & 1 \end{bmatrix} \right) = -1$$

Are there any more?

No!!!

Are there any more?

No!!!

It can be shown that all TP matrices are products of bidiagonal and diagonal matrices (Whitney 1952), (Loewner 1955), (Cryer 1972), (Gasca Peña 1990). Fallat (2001) mentions historical remarks about this.

Are there any more?

No!!!

It can be shown that all TP matrices are products of bidiagonal and diagonal matrices (Whitney 1952), (Loewner 1955), (Cryer 1972), (Gasca Peña 1990). Fallat (2001) mentions historical remarks about this.

Infact, this result is algorithmic and we get an algorithm for bidiagonal factorisation of a matrix called Neville elimination!!

Are there any more?

No!!!

It can be shown that all TP matrices are products of bidiagonal and diagonal matrices (Whitney 1952), (Loewner 1955), (Cryer 1972), (Gasca Peña 1990). Fallat (2001) mentions historical remarks about this.

Infact, this result is algorithmic and we get an algorithm for bidiagonal factorisation of a matrix called Neville elimination!!

The algorithm runs in polynomial time and which also provides a polynomial time algorithm to check TP.

What if the entries of the matrix are polynomials?

We can extend the definition of TP to include matrix with polynomial (in one or several variables) entries.

In this case the minors are also polynomials and we have two notions of positivity in this case:

What if the entries of the matrix are polynomials?

We can extend the definition of TP to include matrix with polynomial (in one or several variables) entries.

In this case the minors are also polynomials and we have two notions of positivity in this case:

1. Coefficientwise positivity: all monomials have positive coefficients.

What if the entries of the matrix are polynomials?

We can extend the definition of TP to include matrix with polynomial (in one or several variables) entries.

In this case the minors are also polynomials and we have two notions of positivity in this case:

1. Coefficientwise positivity: all monomials have positive coefficients.
2. Pointwise positivity: all non-negative substitutions give non-negative values.

What if the entries of the matrix are polynomials?

We can extend the definition of TP to include matrix with polynomial (in one or several variables) entries.

In this case the minors are also polynomials and we have two notions of positivity in this case:

1. Coefficientwise positivity: all monomials have positive coefficients.
2. Pointwise positivity: all non-negative substitutions give non-negative values.

Coefficientwise Positivity \implies Pointwise Positivity

Converse is not true. Example: $x^2 - 2x + 1$.

- 1 Introduction
- 2 Proof techniques and some special types of matrices
 - 1 LGV lemma
 - 2 Hankel matrices
 - 3 Toeplitz matrices
 - 4 Lower triangular matrices
- 3 The Eulerian triangle

The essence of mathematics is proving theorems - and so, that is what mathematicians do: They prove theorems. But to tell the truth, what they really want to prove, once in their lifetime, is a Lemma, like the one by Fatou in analysis, the Lemma of Gauss in number theory, or the Burnside-Frobenius Lemma in combinatorics.

...

- Chapter 25, Lattice Paths and Determinants
Proofs from the Book

The essence of mathematics is proving theorems - and so, that is what mathematicians do: They prove theorems. But to tell the truth, what they really want to prove, once in their lifetime, is a Lemma, like the one by Fatou in analysis, the Lemma of Gauss in number theory, or the Burnside-Frobenius Lemma in combinatorics.

... In this chapter we look at one such marvelous piece of mathematical reasoning, a counting lemma that first appeared in a paper by Bernt Lindström in 1972. Largely overlooked at the time, the result became an instant classic in 1985, when Ira Gessel and Gerard Viennot rediscovered it and demonstrated in a wonderful paper how the lemma could be successfully applied to a diversity of difficult combinatorial enumeration problems.

- Chapter 25, Lattice Paths and Determinants
Proofs from the Book

Let $G = (V, E)$ be an acyclic digraph, which is possibly infinite but each vertex has finitely many neighbours,

Let $G = (V, E)$ be an acyclic digraph, which is possibly infinite but each vertex has finitely many neighbours, such that each of its edges e have weights $w_e \in R$ where R is a commutative ring.

Let $G = (V, E)$ be an acyclic digraph, which is possibly infinite but each vertex has finitely many neighbours, such that each of its edges e have weights $w_e \in R$ where R is a commutative ring.

The weight of a directed path \mathcal{P} , $w(\mathcal{P})$ is the product of the weight of the edges traversed by the path and the weight of a collection of paths is the product of the weight of the individual paths.

Let $G = (V, E)$ be an acyclic digraph, which is possibly infinite but each vertex has finitely many neighbours, such that each of its edges e have weights $w_e \in R$ where R is a commutative ring.

The weight of a directed path \mathcal{P} , $w(\mathcal{P})$ is the product of the weight of the edges traversed by the path and the weight of a collection of paths is the product of the weight of the individual paths.

Given $u, v \in V$, we define

$$P(u \rightarrow v) = \sum_{\mathcal{P}: u \rightarrow v} w(\mathcal{P}).$$

LGV lemma

Given $u, v \in V$, we define

$$P(u \rightarrow v) = \sum_{\mathcal{P}: u \rightarrow v} w(\mathcal{P}).$$

LGV lemma

Given $u, v \in V$, we define

$$P(u \rightarrow v) = \sum_{\mathcal{P}: u \rightarrow v} w(\mathcal{P}).$$

Let $\mathcal{U} = \{u_1, u_2, \dots, u_n\}$ and $\mathcal{V} = \{v_1, v_2, \dots, v_n\}$ be two sets of vertices which we call sources and sinks, respectively.

LGV lemma

Given $u, v \in V$, we define

$$P(u \rightarrow v) = \sum_{\mathcal{P}: u \rightarrow v} w(\mathcal{P}).$$

Let $\mathcal{U} = \{u_1, u_2, \dots, u_n\}$ and $\mathcal{V} = \{v_1, v_2, \dots, v_n\}$ be two sets of vertices which we call sources and sinks, respectively.

Let M be the matrix whose ij^{th} entry is $P(u_i \rightarrow v_j)$.

LGV lemma

Given $u, v \in V$, we define

$$P(u \rightarrow v) = \sum_{\mathcal{P}: u \rightarrow v} w(\mathcal{P}).$$

Let $\mathcal{U} = \{u_1, u_2, \dots, u_n\}$ and $\mathcal{V} = \{v_1, v_2, \dots, v_n\}$ be two sets of vertices which we call sources and sinks, respectively.

Let M be the matrix whose ij^{th} entry is $P(u_i \rightarrow v_j)$.

Then

$$\det(M) = \sum_{\sigma \in S_n} \text{sign}(\sigma) P(u_1 \rightarrow v_{\sigma(1)}) \cdots P(u_n \rightarrow v_{\sigma(n)})$$

LGV lemma

Given $u, v \in V$, we define

$$P(u \rightarrow v) = \sum_{\mathcal{P}: u \rightarrow v} w(\mathcal{P}).$$

Let $\mathcal{U} = \{u_1, u_2, \dots, u_n\}$ and $\mathcal{V} = \{v_1, v_2, \dots, v_n\}$ be two sets of vertices which we call sources and sinks, respectively.

Let M be the matrix whose ij^{th} entry is $P(u_i \rightarrow v_j)$.

Then

$$\det(M) = \sum_{\substack{\mathcal{P}_1: u_1 \rightarrow v_{\sigma(1)} \\ \vdots \\ \mathcal{P}_n: u_n \rightarrow v_{\sigma(n)}}} \text{sign}(\sigma) w(\mathcal{P}_1) \dots w(\mathcal{P}_n)$$

LGV lemma

Given $u, v \in V$, we define

$$P(u \rightarrow v) = \sum_{\mathcal{P}: u \rightarrow v} w(\mathcal{P}).$$

Let $\mathcal{U} = \{u_1, u_2, \dots, u_n\}$ and $\mathcal{V} = \{v_1, v_2, \dots, v_n\}$ be two sets of vertices which we call sources and sinks, respectively.

Let M be the matrix whose ij^{th} entry is $P(u_i \rightarrow v_j)$.

Then

$$\det(M) = \sum_{\substack{\mathcal{P}_1: u_1 \rightarrow v_{\sigma(1)} \\ \vdots \\ \mathcal{P}_n: u_n \rightarrow v_{\sigma(n)} \\ \mathcal{P}_1, \dots, \mathcal{P}_n \text{ vertex disjoint}}} \text{sign}(\sigma) w(\mathcal{P}_1) \dots w(\mathcal{P}_n)$$

Given $u, v \in V$, we define

$$P(u \rightarrow v) = \sum_{\mathcal{P}: u \rightarrow v} w(\mathcal{P}).$$

Let $\mathcal{U} = \{u_1, u_2, \dots, u_n\}$ and $\mathcal{V} = \{v_1, v_2, \dots, v_n\}$ be two sets of vertices which we call sources and sinks, respectively.

Let M be the matrix whose ij^{th} entry is $P(u_i \rightarrow v_j)$.

Then

$$\det(M) = \sum_{\substack{\mathcal{P}_1: u_1 \rightarrow v_{\sigma(1)} \\ \vdots \\ \mathcal{P}_n: u_n \rightarrow v_{\sigma(n)} \\ \mathcal{P}_1, \dots, \mathcal{P}_n \text{ vertex disjoint}}} \text{sign}(\sigma) w(\mathcal{P}_1) \dots w(\mathcal{P}_n)$$

In most digraphs that we work with, the only non-intersecting paths occur when the permutation σ is the identity permutation.

Given $u, v \in V$, we define

$$P(u \rightarrow v) = \sum_{\mathcal{P}: u \rightarrow v} w(\mathcal{P}).$$

Let $\mathcal{U} = \{u_1, u_2, \dots, u_n\}$ and $\mathcal{V} = \{v_1, v_2, \dots, v_n\}$ be two sets of vertices which we call sources and sinks, respectively.


Let M be the matrix whose ij^{th} entry is $P(u_i \rightarrow v_j)$.

Then

$$\det(M) = \sum_{\substack{\mathcal{P}_1: u_1 \rightarrow v_{\sigma(1)} \\ \vdots \\ \mathcal{P}_n: u_n \rightarrow v_{\sigma(n)} \\ \mathcal{P}_1, \dots, \mathcal{P}_n \text{ vertex disjoint}}} \text{sign}(\sigma) w(\mathcal{P}_1) \dots w(\mathcal{P}_n)$$


In most digraphs that we work with, the only non-intersecting paths occur when the permutation σ is the identity permutation. Furthermore, if R is the \mathbb{R} , $\mathbb{R}[x]$, $\mathbb{R}[x, y, z]$, etc. where each edge has non-negative weight, the determinant computed is non-negative.

Example: Lower Bidiagonal matrix



$$\begin{bmatrix} d_1 & 0 & 0 & 0 & 0 \\ r_1 & d_2 & 0 & 0 & 0 \\ 0 & r_2 & d_3 & 0 & 0 \\ 0 & 0 & r_3 & d_4 & 0 \\ 0 & 0 & 0 & r_4 & d_5 \end{bmatrix}$$

Example: Upper Bidiagonal matrix



$$\begin{bmatrix} d_1 & r_1 & 0 & 0 & 0 & 0 \\ 0 & d_2 & r_2 & 0 & 0 & 0 \\ 0 & 0 & d_3 & r_3 & 0 & 0 \\ 0 & 0 & 0 & d_4 & r_4 & 0 \\ 0 & 0 & 0 & 0 & d_5 & r_5 \\ 0 & 0 & 0 & 0 & 0 & d_6 \end{bmatrix}$$

Bidiagonal Matrix

Neville elimination gives us that a real matrix is TP if and only if it has an LGV digraph!

Bidiagonal Matrix

Neville elimination gives us that a real matrix is TP if and only if it has an LGV digraph!

In general, Existence of an LGV digraph \implies TP.

Bidiagonal Matrix

Neville elimination gives us that a real matrix is TP if and only if it has an LGV digraph!

In general, Existence of an LGV digraph \implies TP.

Converse not true in general.


Bidiagonal Matrix

Neville elimination gives us that a real matrix is TP if and only if it has an LGV digraph!

In general, Existence of an LGV digraph \implies TP.

Converse not true in general. Not even true for integers as Neville factorisation involves division.


Example:



All edges have weights 1 and are directed from left to right.

$$P(u_n \rightarrow v_k) = ??$$


Example: Binomial Triangle



All edges have weights 1 and are directed from left to right.

$$P(u_n \rightarrow v_k) = \binom{n}{k}$$

Example: Binomial Triangle



All edges have weights 1 and are directed from left to right.

$$\begin{bmatrix} 1 & 0 & 0 & 0 & 0 \\ 1 & 1 & 0 & 0 & 0 \\ 1 & 2 & 1 & 0 & 0 \\ 1 & 3 & 3 & 1 & 0 \\ 1 & 4 & 6 & 4 & 1 \end{bmatrix}$$

- 1 Introduction
- 2 Proof techniques and some special types of matrices
 - 1 LGV lemma
 - 2 Hankel matrices
 - 3 Toeplitz matrices
 - 4 Lower triangular matrices
- 3 The Eulerian triangle

Hankel Matrix

Given a sequence a_0, a_1, \dots the infinite matrix $H_\infty(\mathbf{a})$ whose ij^{th} entry is a_{i+j} is called the Hankel matrix of $(a_n)_{n \geq 0}$.

$$\begin{array}{cccccc} a_0 & a_1 & a_2 & a_3 & a_4 & \dots \\ a_1 & a_2 & a_3 & a_4 & a_5 & \dots \\ a_2 & a_3 & a_4 & a_5 & a_6 & \dots \\ a_3 & a_4 & a_5 & a_6 & a_7 & \dots \\ a_4 & a_5 & a_6 & a_7 & a_8 & \dots \\ \vdots & \vdots & \vdots & \vdots & \vdots & \end{array}$$

Hankel Matrix

Given a sequence a_0, a_1, \dots the infinite matrix $H_\infty(\mathbf{a})$ whose ij^{th} entry is a_{i+j} is called the Hankel matrix of $(a_n)_{n \geq 0}$.

$$\begin{array}{cccccc} a_0 & a_1 & a_2 & a_3 & a_4 & \dots \\ a_1 & a_2 & a_3 & a_4 & a_5 & \dots \\ a_2 & a_3 & a_4 & a_5 & a_6 & \dots \\ a_3 & a_4 & a_5 & a_6 & a_7 & \dots \\ a_4 & a_5 & a_6 & a_7 & a_8 & \dots \\ \vdots & \vdots & \vdots & \vdots & \vdots & \end{array}$$

We say that a sequence $(a_n)_{n \geq 0}$ is Hankel-totally positive (Hankel-TP in short) if its Hankel matrix is TP.

Fundamental Fact about Hankel-TP

Theorem (Stieltjes, Gantmacher-Krein)

For a sequence $(a_n)_{n \geq 0}$ of real numbers, the following are equivalent:

- 1 $(a_n)_{n \geq 0}$ is Hankel-TP.
- 2 There exists a positive measure μ on $[0, \infty)$ such that $a_n = \int_0^\infty x^n d\mu$ for all $n \geq 0$. ($(a_n)_{n \geq 0}$ is a Stieltjes moment sequence.)
- 3 There exists numbers $\alpha_0, \alpha_1, \dots \geq 0$ such that

$$\sum_{n=0}^{\infty} a_n t^n = \frac{\alpha_0}{1 - \frac{\alpha_1 t}{1 - \frac{\alpha_2 t^2}{1 - \ddots}}}$$

or in other words the generating function of $(a_n)_{n \geq 0}$ has a Stieltjes continued fraction expansion.

Fundamental Fact about Hankel-TP

Theorem (Stieltjes, Gantmacher-Krein)

For a sequence $(a_n)_{n \geq 0}$ of real numbers, the following are equivalent:

- 1 $(a_n)_{n \geq 0}$ is Hankel-TP.
- 2 There exists a positive measure μ on $[0, \infty)$ such that $a_n = \int_0^\infty x^n d\mu$ for all $n \geq 0$. ($(a_n)_{n \geq 0}$ is a Stieltjes moment sequence.)
- 3 There exists numbers $\alpha_0, \alpha_1, \dots \geq 0$ such that

$$\sum_{n=0}^{\infty} a_n t^n = \frac{\alpha_0}{1 - \frac{\alpha_1 t}{1 - \frac{\alpha_2 t^2}{1 - \ddots}}}$$

or in other words the generating function of $(a_n)_{n \geq 0}$ has a Stieltjes continued fraction expansion.

This doesn't hold entirely in the general case.

Combinatorial interpretations of Stieltjes Moment Sequences

Flajolet (1980) gave a combinatorial interpretation of Stieltjes continued fractions (and also Jacobi continued fractions) by showing that Stieltjes continued fractions are generating series of Dyck paths (Motzkin paths resp.) with height dependent weights.

Combinatorial interpretations of Stieltjes Moment Sequences

Flajolet (1980) gave a combinatorial interpretation of Stieltjes continued fractions (and also Jacobi continued fractions) by showing that Stieltjes continued fractions are generating series of Dyck paths (Motzkin paths resp.) with height dependent weights.

Using this interpretation he gave Stieltjes continued fractions for several sequences such as Catalan numbers, Bell and Stirling numbers, tangent and secant numbers, etc. and of orthogonal polynomials such as Tchebycheff, Charlier, Hermite, Laguerre and Meixner polynomials.

Combinatorial interpretations of Stieltjes Moment Sequences

Flajolet (1980) gave a combinatorial interpretation of Stieltjes continued fractions (and also Jacobi continued fractions) by showing that Stieltjes continued fractions are generating series of Dyck paths (Motzkin paths resp.) with height dependent weights.

Using this interpretation he gave Stieltjes continued fractions for several sequences such as Catalan numbers, Bell and Stirling numbers, tangent and secant numbers, etc. and of orthogonal polynomials such as Tchebycheff, Charlier, Hermite, Laguerre and Meixner polynomials.

Sokal and his collaborators have used the interpretation of Flajolet and generalisations along with LGV lemma to prove several sufficient conditions for coefficientwise Hankel-TP.

- 1 Introduction
- 2 Proof techniques and some special types of matrices
 - 1 LGV lemma
 - 2 Hankel matrices
 - 3 Toeplitz matrices
 - 4 Lower triangular matrices
- 3 The Eulerian triangle

Toeplitz Matrix

Given a sequence a_0, a_1, \dots the infinite matrix $T_\infty(\mathbf{a})$ whose ij^{th} entry is a_{i-j} for $i \leq j$ and 0 otherwise is called the Toeplitz matrix of $(a_n)_{n \geq 0}$.

$$\begin{array}{cccccc} a_0 & 0 & 0 & 0 & 0 & \dots \\ a_1 & a_0 & 0 & 0 & 0 & \dots \\ a_2 & a_1 & a_0 & 0 & 0 & \dots \\ a_3 & a_2 & a_1 & a_0 & 0 & \dots \\ a_4 & a_3 & a_2 & a_1 & a_0 & \dots \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \end{array}$$

Toeplitz Matrix

Given a sequence a_0, a_1, \dots the infinite matrix $T_\infty(\mathbf{a})$ whose ij^{th} entry is a_{i-j} for $i \leq j$ and 0 otherwise is called the Toeplitz matrix of $(a_n)_{n \geq 0}$.

$$\begin{array}{cccccc} a_0 & 0 & 0 & 0 & 0 & \dots \\ a_1 & a_0 & 0 & 0 & 0 & \dots \\ a_2 & a_1 & a_0 & 0 & 0 & \dots \\ a_3 & a_2 & a_1 & a_0 & 0 & \dots \\ a_4 & a_3 & a_2 & a_1 & a_0 & \dots \\ \vdots & \vdots & \vdots & \vdots & \vdots & \end{array}$$

We say that a sequence $(a_n)_{n \geq 0}$ is Toeplitz-totally positive (Toeplitz-TP in short) or a Polya frequency sequence (PF sequence) if its Toeplitz matrix is TP.

Fundamental Fact about Toeplitz-TP

Theorem (Aissen-Schoenberg-Whitney, Edrei 1952)

For a sequence $(a_n)_{n \geq 0}$ of real numbers with $a_0 = 1$, the following are equivalent:

- ① *$(a_n)_{n \geq 0}$ is Toeplitz-TP.*
- ② *There exists $\alpha_i \geq 0$, $\beta_j \geq 0$ and $\gamma \geq 0$ such that*

$$\sum_{n=0}^{\infty} a_n t^n = e^{\gamma t} \frac{\prod_i (1 + \alpha_i t)}{\prod_j (1 - \beta_j t)}.$$

Fundamental Fact about Toeplitz-TP

Theorem (Aissen-Schoenberg-Whitney, Edrei 1952)

For a sequence $(a_n)_{n \geq 0}$ of real numbers with $a_0 = 1$, the following are equivalent:

- ① *$(a_n)_{n \geq 0}$ is Toeplitz-TP.*
- ② *There exists $\alpha_i \geq 0$, $\beta_j \geq 0$ and $\gamma \geq 0$ such that*

$$\sum_{n=0}^{\infty} a_n t^n = e^{\gamma t} \frac{\prod_i (1 + \alpha_i t)}{\prod_j (1 - \beta_j t)}.$$

$(2) \implies (1)$ is easy and even holds coefficientwise.

Fundamental Fact about Toeplitz-TP

Theorem (Aissen-Schoenberg-Whitney, Edrei 1952)

For a sequence $(a_n)_{n \geq 0}$ of real numbers with $a_0 = 1$, the following are equivalent:

- ① *$(a_n)_{n \geq 0}$ is Toeplitz-TP.*
- ② *There exists $\alpha_i \geq 0$, $\beta_j \geq 0$ and $\gamma \geq 0$ such that*

$$\sum_{n=0}^{\infty} a_n t^n = e^{\gamma t} \frac{\prod_i (1 + \alpha_i t)}{\prod_j (1 - \beta_j t)}.$$

(2) \implies (1) is easy and even holds coefficientwise. (1) \implies (2) is hard and requires Nevanlinna theory of meromorphic functions.

Fundamental Fact about Toeplitz-TP

Theorem (Aissen-Schoenberg-Whitney, Edrei 1952)

For a sequence $(a_n)_{n \geq 0}$ of real numbers with $a_0 = 1$, the following are equivalent:

- ① *$(a_n)_{n \geq 0}$ is Toeplitz-TP.*
- ② *There exists $\alpha_i \geq 0$, $\beta_j \geq 0$ and $\gamma \geq 0$ such that*

$$\sum_{n=0}^{\infty} a_n t^n = e^{\gamma t} \frac{\prod_i (1 + \alpha_i t)}{\prod_j (1 - \beta_j t)}.$$

$(2) \implies (1)$ is easy and even holds coefficientwise. $(1) \implies (2)$ is hard and requires Nevanlinna theory of meromorphic functions.

For a finite sequence we need to show that the generating polynomial is negative real rooted.

Theorem (Katkova 2006)

Let $\xi(z) = \frac{1}{2}z(z-1)\pi^{-z/2}\Gamma(z/2)\zeta(z)$ be the Reimann- ξ function and let $\xi_1(z) = \xi(\sqrt{z} + 1/2)$.

Connection of Toeplitz-TP to Reimann Hypothesis

Theorem (Katkova 2006)

Let $\xi(z) = \frac{1}{2}z(z-1)\pi^{-z/2}\Gamma(z/2)\zeta(z)$ be the Reimann- ξ function and let $\xi_1(z) = \xi(\sqrt{z} + 1/2)$. Then ξ_1 is Toeplitz-TP if and only if the Reimann Hypothesis is true.

Log Concavity and Log Convexity

Consider the sequence $(a_n)_{n \geq 0}$.

Log Concavity and Log Convexity

Consider the sequence $(a_n)_{n \geq 0}$.

Toeplitz- TP_2 implies log-concavity i.e., $a_n^2 - a_{n-1}a_{n+1} \geq 0$.

Hankel- TP_2 implies log-convexity i.e., $a_n^2 - a_{n-1}a_{n+1} \leq 0$.

Lower Triangular Matrices and three types of TP

Consider an infinite lower triangular matrix A with entries a_{ij} where the indexing of the rows and columns begins from 0.

$$\begin{array}{cccccc} a_{00} & 0 & 0 & 0 & 0 & \dots \\ a_{10} & a_{11} & 0 & 0 & 0 & \dots \\ a_{20} & a_{21} & a_{22} & 0 & 0 & \dots \\ a_{30} & a_{31} & a_{32} & a_{33} & 0 & \dots \\ a_{40} & a_{41} & a_{42} & a_{43} & a_{44} & \dots \\ \vdots & \vdots & \vdots & \vdots & \vdots & \end{array}$$

- 1 Introduction
- 2 Proof techniques and some special types of matrices
 - 1 LGV lemma
 - 2 Hankel matrices
 - 3 Toeplitz matrices
 - 4 Lower triangular matrices
- 3 The Eulerian triangle

Lower Triangular Matrices and three types of TP

Let $A_n(x)$ denote the row generating polynomial of the n^{th} row i.e.

$$A_n(x) = \sum_{i=0}^n a_{ni} x^i$$

Lower Triangular Matrices and three types of TP

Let $A_n(x)$ denote the row generating polynomial of the n^{th} row i.e.

$$A_n(x) = \sum_{i=0}^n a_{ni} x^i$$

We can ask three different questions:

Lower Triangular Matrices and three types of TP

Let $A_n(x)$ denote the row generating polynomial of the n^{th} row i.e.

$$A_n(x) = \sum_{i=0}^n a_{ni} x^i$$

We can ask three different questions:

- 1 Is A TP?

Lower Triangular Matrices and three types of TP

Let $A_n(x)$ denote the row generating polynomial of the n^{th} row i.e.

$$A_n(x) = \sum_{i=0}^n a_{ni} x^i$$

We can ask three different questions:

- 1 Is A TP?
- 2 Are $A_n(x)$ negative real rooted or is the finite sequence a_{n0}, \dots, a_{nn} Toeplitz-TP?

Lower Triangular Matrices and three types of TP

Let $A_n(x)$ denote the row generating polynomial of the n^{th} row i.e.

$$A_n(x) = \sum_{i=0}^n a_{ni} x^i$$

We can ask three different questions:

- 1 Is A TP?
- 2 Are $A_n(x)$ negative real rooted or is the finite sequence a_{n0}, \dots, a_{nn} Toeplitz-TP?
- 3 Is the sequence of polynomials $(A_n(x))_{n \geq 0}$ Hankel-TP?

Lower Triangular Matrices and three types of TP

Let $A_n(x)$ denote the row generating polynomial of the n^{th} row i.e.

$$A_n(x) = \sum_{i=0}^n a_{ni} x^i$$

We can ask three different questions:

- 1 Is A TP?
- 2 Are $A_n(x)$ negative real rooted or is the finite sequence a_{n0}, \dots, a_{nn} Toeplitz-TP?
- 3 Is the sequence of polynomials $(A_n(x))_{n \geq 0}$ Hankel-TP?

In several cases where the answer to one of these questions is true, the others also seem to be true. It is not clear which way the implications hold.

Easy example: Binomial Triangle

For A the triangle of binomial numbers,

$$A = \begin{bmatrix} 1 & 0 & 0 & 0 & 0 & \dots \\ 1 & 1 & 0 & 0 & 0 & \dots \\ 1 & 2 & 1 & 0 & 0 & \dots \\ 1 & 3 & 3 & 1 & 0 & \dots \\ 1 & 4 & 6 & 4 & 1 & \dots \\ \vdots & \vdots & \vdots & \vdots & \vdots & \end{bmatrix}$$

Easy example: Binomial Triangle

For A the triangle of binomial numbers,

$$A = \begin{bmatrix} 1 & 0 & 0 & 0 & 0 & \dots \\ 1 & 1 & 0 & 0 & 0 & \dots \\ 1 & 2 & 1 & 0 & 0 & \dots \\ 1 & 3 & 3 & 1 & 0 & \dots \\ 1 & 4 & 6 & 4 & 1 & \dots \\ \vdots & \vdots & \vdots & \vdots & \vdots & \end{bmatrix}$$

- 1 A is TP by LGV lemma.

Easy example: Binomial Triangle

For A the triangle of binomial numbers,

$$A = \begin{bmatrix} 1 & 0 & 0 & 0 & 0 & \dots \\ 1 & 1 & 0 & 0 & 0 & \dots \\ 1 & 2 & 1 & 0 & 0 & \dots \\ 1 & 3 & 3 & 1 & 0 & \dots \\ 1 & 4 & 6 & 4 & 1 & \dots \\ \vdots & \vdots & \vdots & \vdots & \vdots & \ddots \end{bmatrix}$$

- ① A is TP by LGV lemma.
- ② The row generating polynomials $A_n(x) = (1+x)^n$ are clearly negative real rooted and hence the sequence $\binom{n}{0}, \dots, \binom{n}{n}$ is Toeplitz-TP.

Easy example: Binomial Triangle

For A the triangle of binomial numbers,

$$A = \begin{bmatrix} 1 & 0 & 0 & 0 & 0 & \dots \\ 1 & 1 & 0 & 0 & 0 & \dots \\ 1 & 2 & 1 & 0 & 0 & \dots \\ 1 & 3 & 3 & 1 & 0 & \dots \\ 1 & 4 & 6 & 4 & 1 & \dots \\ \vdots & \vdots & \vdots & \vdots & \vdots & \ddots \end{bmatrix}$$

- ① A is TP by LGV lemma.
- ② The row generating polynomials $A_n(x) = (1+x)^n$ are clearly negative real rooted and hence the sequence $\binom{n}{0}, \dots, \binom{n}{n}$ is Toeplitz-TP.
- ③ The sequence $(A_n(x))$ is Hankel-TP as we have the easy continued fraction expansion

$$\sum_{n=0}^{\infty} (1+x)^n t^n = \frac{1}{1 - (1+x)t}.$$

- 1 Introduction
- 2 Proof techniques and some special types of matrices
 - 1 LGV lemma
 - 2 Hankel matrices
 - 3 Toeplitz matrices
 - 4 Lower triangular matrices
- 3 The Eulerian triangle**

Ongoing joint work with X. Chen, A. Dyachenko, T. Gilmore, A.D. Sokal.



Ongoing joint work with X. Chen, A. Dyachenko, T. Gilmore, A.D. Sokal.

Special case of the Eulerian triangle

We can define the Eulerian numbers:

Definition

The *descent* of a permutation of $\sigma \in \mathfrak{S}_n$ is an $i \in [n-1]$ such that $\sigma(i) > \sigma(i+1)$. Let $\text{des}(\sigma)$ denote the number of descents of a permutation σ .

Example: For $\sigma = 943127685$ the $\text{des}(\sigma) = \{1, 2, 3, 6, 8\}$

Special case of the Eulerian triangle

We can define the Eulerian numbers:

Definition

The *descent* of a permutation of $\sigma \in \mathfrak{S}_n$ is an $i \in [n-1]$ such that $\sigma(i) > \sigma(i+1)$. Let $\text{des}(\sigma)$ denote the number of descents of a permutation σ .

Example: For $\sigma = 943127685$ the $\text{des}(\sigma) = \{1, 2, 3, 6, 8\}$

Definition

The Eulerian number $A_{n,k}$ is defined to be the cardinality of the set $\{\sigma \in \mathfrak{S}_{n+1} \mid \text{des}(\sigma) = k\}$.

Special case of the Eulerian triangle

We can define the Eulerian numbers:

Definition

The *descent* of a permutation of $\sigma \in \mathfrak{S}_n$ is an $i \in [n-1]$ such that $\sigma(i) > \sigma(i+1)$. Let $\text{des}(\sigma)$ denote the number of descents of a permutation σ .

Example: For $\sigma = 943127685$ the $\text{des}(\sigma) = \{1, 2, 3, 6, 8\}$

Definition

The Eulerian number $A_{n,k}$ is defined to be the cardinality of the set $\{\sigma \in \mathfrak{S}_{n+1} \mid \text{des}(\sigma) = k\}$.

These numbers satisfy the recurrence

$$A_{n,k} = (n-k+1)A_{n-1,k-1} + (k+1)A_{n-1,k}.$$

Eulerian Triangle Conjecture

Conjecture (Brenti 1996)

The infinite lower triangular matrix whose $(n, k)^{\text{th}}$ entry is the Eulerian number $A_{n,k}$ is totally positive.

Eulerian Triangle Conjecture

Conjecture (Brenti 1996)

The infinite lower triangular matrix whose $(n, k)^{\text{th}}$ entry is the Eulerian number $A_{n,k}$ is totally positive.

Using Neville elimination we have verified the conjecture for the upper left 500×500 submatrix. But the general pattern is not clear.

Putting variables in the recurrence

We can define the matrix just from the recurrence

$$A_{n,k} = (n - k + 1)A_{n-1,k-1} + (k + 1)A_{n-1,k}.$$

Putting variables in the recurrence

We can define the matrix just from the recurrence

$$A_{n,k} = (n - k + 1)A_{n-1,k-1} + (k + 1)A_{n-1,k}.$$

We can instead look at the matrix generated by the recurrence

$$A_{n,k} = [a(n - k) + b(k - 1) + c]A_{n-1,k-1} + (dk + e)A_{n-1,k}$$

and ask if it is coefficientwise TP in all five variables a, b, c, d, e .

Setting $a = 1, b = 0, c = 1, d = 1, e = 1$ gives the Eulerian triangle and hence, coefficientwise TP of the $abcde$ matrix can be used to prove that the Eulerian triangle is TP.

Putting variables in the recurrence

We can define the matrix just from the recurrence

$$A_{n,k} = (n - k + 1)A_{n-1,k-1} + (k + 1)A_{n-1,k}.$$

We can instead look at the matrix generated by the recurrence

$$A_{n,k} = [a(n - k) + b(k - 1) + c]A_{n-1,k-1} + (dk + e)A_{n-1,k}$$

and ask if it is coefficientwise TP in all five variables a, b, c, d, e .

Setting $a = 1, b = 0, c = 1, d = 1, e = 1$ gives the Eulerian triangle and hence, coefficientwise TP of the $abcde$ matrix can be used to prove that the Eulerian triangle is TP.

We checked it upto 11×11 which took about a week to compute.

Putting variables in the recurrence

We can define the matrix just from the recurrence

$$A_{n,k} = (n - k + 1)A_{n-1,k-1} + (k + 1)A_{n-1,k}.$$

We can instead look at the matrix generated by the recurrence


$$A_{n,k} = [a(n - k) + b(k - 1) + c]A_{n-1,k-1} + (dk + e)A_{n-1,k}$$

and ask if it is coefficientwise TP in all five variables a, b, c, d, e .

Setting $a = 1, b = 0, c = 1, d = 1, e = 1$ gives the Eulerian triangle and hence, coefficientwise TP of the $abcde$ matrix can be used to prove that the Eulerian triangle is TP.

We checked it upto 11×11 which took about a week to compute.

Unfortunately the 12×12 matrix turned out false after about 2 months of real time.



Is all hope lost?

We can look at the matrix with $b = 0$ and which is generated by the recurrence

$$A_{n,k} = [a(n - k) + c]A_{n-1,k-1} + (dk + e)A_{n-1,k}$$

and ask if this is coefficientwise TP in all variables a, c, d, e .

Is all hope lost?

We can look at the matrix with $b = 0$ and which is generated by the recurrence

$$A_{n,k} = [a(n - k) + c]A_{n-1,k-1} + (dk + e)A_{n-1,k}$$

and ask if this is coefficientwise TP in all variables a, c, d, e . This seems to be true so far.

Is all hope lost?

We can look at the matrix with $b = 0$ and which is generated by the recurrence

$$A_{n,k} = [a(n-k) + c]A_{n-1,k-1} + (dk + e)A_{n-1,k}$$


and ask if this is coefficientwise TP in all variables a, c, d, e . This seems to be true so far.

Infact, we have the following special cases for different substitutions:

Values of a, c, d, e	Matrix obtained
$a = 1, c = 1, d = 1, e = 1$	Eulerian triangle
$a = 1, c = 0, d = 1, e = 1$	Down-shifted Eulerian triangle
$a = 1, c = 1, d = 1, e = 0$	Diagonally down-shifted Eulerian triangle
$a = 0, c = 1, d = 1, e = 1$	k -dependent recurrence which gives the matrix of Stirling subset numbers. Brenti 1995.
$a = 1, c = 1, d = 0, e = 1$	Row reversed matrix of Stirling subset numbers

The coefficientwise results for cases of $\{a = 0, d = 0\}$ (weighted Binomial triangle), $a = 0$ (weighted Stirling subset triangle) were already known. We managed to resolve the $c = 0$ (weighted reverse Stirling subset) case and gave two different proofs for the result.

The first proof involves a bijective argument involving the LGV lemma. The second proof consists of a linear algebraic method involving matrix products which we call the quasi-production matrix method.



$$T(n, k) = (n - k + 1)T(n - 1, k - 1)$$

$$+ (k + 1)T(n - 1, k)$$



$$T(n, k) = [a(n - k) + c]T(n - 1, k - 1)$$

$$+ (dk + e)T(n - 1, k)$$

$$T(n, k) = (n - k + 1)T(n - 1, k - 1) \\ + (k + 1)T(n - 1, k)$$

Eulerian triangle conjecture

$$T(n, k) = [a(n - k) + c]T(n - 1, k - 1) \\ + (dk + e)T(n - 1, k)$$



acde conjecture

$$T(n, k) = [a(n - k) + c]T(n - 1, k - 1) \\ + (dk + e)T(n - 1, k) \\ + [f(n - 2) + g]T(n - 2, k - 1)$$

acdefg conjecture

$$T(n, k) = [a(n - k) + c]T(n - 1, k - 1) \\ + \left(\sum_{i=0}^k d_i \right) T(n - 1, k) \\ + [f(n - 2) + g]T(n - 2, k - 1)$$

acd^dfg conjecture



- ① Introduction - Definitions and generating examples
- ② Proof techniques and some special types of matrices
 - ① LGV lemma - description and illustration using some simple examples
 - ② Hankel matrices - definition and fundamental theorem relating to Stieltjes moment sequences and Stieltjes Continued fractions
 - ③ Toeplitz matrices - definition and fundamental theorem relating to negative real zeros and positive real poles
 - ④ Lower triangular matrices - three different types of TP
- ③ The Eulerian triangle - the conjecture, partial progress and a lot of other stronger conjectures