

# Lecture slides

Philip Murray

# Lecture 1

- ▶ Introduction to nonlinear difference equations
- ▶ The Malthusian model
- ▶ The Ricker model

Why difference equations?

## A general model

Consider the first order difference equation

$$N_{t+1} = N_t f(N_t) = H(N_t), \quad (1)$$

where  $f(N_t)$  is a function that defines the per capita growth rate.  
The function  $H(N_t)$  describes the total (net) growth rate.

# The Malthusian model

The population size at time  $t + 1$  is

$$N_{t+1} = N_t + bN_t - dN_t = rN_t,$$

Exercise: solve the Malthusian model and classify qualitative behaviours

# Nonlinear models

- ▶ Beverton-Holt

$$N_{t+1} = \frac{rN_t}{1 + \frac{N_t}{K}},$$

- ▶ Hassell model

$$N_{t+1} = \frac{rN_t}{(1 + \frac{N_t}{K})^b},$$

- ▶ Ricker model

$$N_{t+1} = N_t e^{r(1 - \frac{N_t}{K})}.$$

# Numerical simulation of the Ricker model




Figure 1: A plot of numerical solutions of the Ricker model. (a)  $r=0.5$ .



# Summary

- ▶ Motivated use of difference equation models
- ▶ Introduced general model for one population
- ▶ Solved the Malthusian model
- ▶ Introduced nonlinear models

## Lecture 2 - General techniques for solving nonlinear difference equations

$$N_{t+1} = N_t f(N_t) = H(N_t), \quad (2)$$

- ▶ Computational solutions
- ▶ Fixed points
- ▶ Linear stability of fixed points
- ▶ Cobweb diagrams
- ▶ Bifurcation diagrams
- ▶ Identify how model solutions depend on model parameters

## Fixed points

Suppose the solution at the next iteration is equal to that at a given iteration, i.e. there exists some  $N^*$  such that

$$N^* = N_{t+1} = N_t$$



Fixed point definition

$$N^* = H(N^*), N^* \geq 0$$

Biological relevance: non-negative solutions

Linear stability analysis - how do small perturbations about  $N^*$  behave?

## Linear stability analysis (ctd)

💡 Linear stability is determined by the derivative of  $H$  evaluated at  $N^*$

$$|H'(N^*)| < 1 \implies \text{linear stability of } N^*.$$

## Exercise

Identify the fixed points of the Malthusian model

$$N_{t+1} = rN_t$$

and identify their linear stability.

# Cobweb diagrams

## Definition

A cobweb diagram is a technique for computing graphical solutions of a difference equation.

Use previous analyses to identify different qualitative cases (one cobweb diagram for each fixed point).

For each case:

- ▶ Sketch a graph of  $H$  to evaluate iterative solutions
- ▶ Compute an iterative solution

# Example




Figure 2: Generating a cobweb plot.



Bifurcation diagrams - Plot fixed points against a parameter and annotate their stability

## Exercise

Draw cobweb diagrams for the Malthusian model.

$$N_{t+1} = rN_t$$

and identify their linear stability.

## Lecture 3 - Preparation for tutorial 3

- Curve sketching nonlinear functions in qualitatively distinct cases

Example:

Sketch a graph of

$$f(x) = xe^{-r(1-\frac{x}{K})}, \quad r, K \in \mathfrak{R}^+, \quad x \in \mathfrak{R}, x \geq 0$$

### Approach

Identify properties of  $H$  to distinguish qualitatively distinct cases

# Roots

# Turning points

Limit as  $x \rightarrow \infty$

Limiting behaviour as  $x \rightarrow 0$

# Tutorial sheet 1



## Lecture 4

Consider the model

$$N_{t+1} = \frac{\gamma N_t}{1 + N_t^2}, \quad \gamma \in \mathfrak{R}^+.$$

# Fixed points

# Linear stability

# Cobweb diagrams

# Bifurcations

## Symbolic computations

The FPs are:

[{N: 0}, {N: -sqrt(gamma - 1)}, {N: sqrt(gamma - 1)}]

The derivative of H is:

$\text{gamma} \cdot (1 - N^2) / (N^2 + 1)^2$

The derivative evaluated at FP 1 is:

gamma

The derivative evaluated at FP 2 is:

$(2 - \text{gamma}) / \text{gamma}$