

thornado-Hydro (xCFC)

Samuel J. Dunham

October 21, 2024




toolkit for high-order neutrino-radiation hydrodynamics

- DG
- SSPRK/IMEX
- GR (xCFC)
- Hydro^a (Valencia)
- Neutrino transport^b (M1)
- Interfaces to tabulated EoS/Opacities (weaklib: <https://github.com/starkiller-astro/weaklib>)
- Fluid self-gravity via Poseidon: <https://github.com/jrober50/Poseidon>
- GPUs via OpenACC or OpenMP pragmas
- MPI parallelism and AMR via AMReX: <https://github.com/AMReX-Codes/amrex>

^aEndeve et al. (2019); Dunham et al. (2020); Pochik et al. (2021)

^bLaiu et al. (2021)

3+1 Decomposition



$$ds^2 = g_{\mu\nu} dx^\mu dx^\nu = -\alpha^2 dt^2 + \gamma_{ij} (dx^i + \beta^i dt) (dx^j + \beta^j dt)$$

Conformally-Flat Condition

Developed by Wilson et al. (1996),
extended by Cordero-Carrión et al.
(2009)

Special case: Schwarzschild spacetime
in isotropic coordinates ($G = c = 1$)

$$\gamma_{ij}(x) = \psi^4(x) \bar{\gamma}_{ij}(x^i)$$

$$K = 0, \quad \partial_t K = 0$$

(Always and everywhere)

$$\alpha = \left(1 + \frac{1}{2}\Phi\right) \left(1 - \frac{1}{2}\Phi\right)^{-1}$$

$$\psi = 1 - \frac{1}{2}\Phi$$


$$\beta^i = 0,$$

- Exact in spherical symmetry!
- Hyperbolic \rightarrow Elliptic equations
- Good for long-time simulations


with

$$\Phi(r) := -\frac{M}{r}$$


Discontinuous Galerkin (DG) Method




Discontinuous Galerkin (DG) Method




Discontinuous Galerkin (DG) Method



Discontinuous Galerkin (DG) Method




Discontinuous Galerkin (DG) Method




Discontinuous Galerkin (DG) Method

$$u_h(x, t) := \sum_{i=1}^{k+1} u_i(t) \ell_i(x)$$




- For hydro-only, strong-stability-preserving Runge–Kutta (SSPRK) methods (e.g., see Gottlieb et al. (2001))
 - Convex combinations of forward-Euler timesteps
- For hydro+neutrinos, ImEx methods (e.g., see Pareschi and Russo (2005))

Convergence Rates for Sine Wave Advection (1D)




Slope Limiter (MinMod)




Slope Limiter (MinMod)

$$k = 9$$




Slope Limiter (MinMod)




Slope Limiter (MinMod)

$$\bar{u}(t) := \frac{1}{V(t)} \int_K u_h(x, t) \sqrt{\gamma}(x, t) dx, \quad V(t) := \int_K \sqrt{\gamma}(x, t) dx$$




Slope Limiter (MinMod)

$$u_h(x, t) = \sum_{i=1}^{k+1} u_i(t) \ell_i(x) = \sum_{n=0}^k C_n(t) P_n(x)$$




Slope Limiter (MinMod)

-- : $\bar{u}_K - \bar{u}_{K-1}$ -- : $\bar{u}_{K+1} - \bar{u}_K$



Slope Limiter (MinMod)


$$\sum_{n=1}^k C_n P_n = u_h \rightarrow \tilde{u}_h := \Lambda_{\text{SL}}(u_h) := C_0 P_0 + \tilde{C}_1 P_1$$



Slope Limiter (MinMod)


$$k = 9$$

$$\tilde{k} = 1$$




Troubled Cell Indicator

Proposed in Fu and Shu (2017)




Bound-Enforcing Limiter (Λ_{BEL})

- Proposed in Qin et al. (2016)
- Define \mathcal{G} as the (convex) set of physically-admissible states; i.e., positivity density and energy density, subluminal fluid velocity (ideal EoS only)



$$u_h = u_1 \ell_1 + u_2 \ell_2$$



$$\xrightarrow{\Lambda_{\text{BEL}}}$$



$$\tilde{u}_h := \Lambda_{\text{BEL}}(u_h) = \tilde{u}_1 \ell_1 + \tilde{u}_2 \ell_2$$


Standing Accretion Shock Instability

Used thornado to investigate the role of GR on the SASI¹



¹Dunham et al. (2020, 2024)

Mesh Refinement with AMReX



Conservative Interpolation (1D)

- Schaal et al. (2015)


Given a solution in the nodal points of a coarse element K , $\{U_j\}_{j=1}^N$, define the solution in the nodal points of the j th ($j \in \{1, 2\}$) fine element $k^{(j)}$, $\{u_i^{(j)}\}_{i=1}^N$ via the minimization of an L^2 -projection:

$$\forall i \in \{1, \dots, N\} : \min_{\tilde{u}_i^{(j)}} \int_{k^{(j)}} \left(\tilde{U}_h - \tilde{u}_h^{(j)} \right)^2 dx,$$




where $\tilde{U} := \sqrt{\gamma} U$. For the geometry fields, we use the same procedure but do not pre-multiply with $\sqrt{\gamma}$.

Show what we store in the flux register


Kelvin–Helmholtz Instability




2D Blast Wave (Del Zanna and Bucciantini, 2002)



Adiabatic Collapse (Collapse Phase)



Adiabatic Collapse (Post-Bounce Phase)





Bibliography

- Eirik Endeve, Jesse Buffaloe, Samuel J. Dunham, Nick Roberts, Kristopher Andrew, Brandon Barker, David Pochik, Juliana Pulsinelli, and Anthony Mezzacappa. thornado-hydro: towards discontinuous Galerkin methods for supernova hydrodynamics. In *Journal of Physics Conference Series*, volume 1225 of *Journal of Physics Conference Series*, page 012014, May 2019. doi: 10.1088/1742-6596/1225/1/012014.
- Samuel J. Dunham, E. Endeve, A. Mezzacappa, J. Buffaloe, and K. Holley-Bockelmann. A discontinuous Galerkin method for general relativistic hydrodynamics in thornado. In *Journal of Physics Conference Series*, volume 1623 of *Journal of Physics Conference Series*, page 012012, September 2020. doi: 10.1088/1742-6596/1623/1/012012.
- David Pochik, Brandon L. Barker, Eirik Endeve, Jesse Buffaloe, Samuel J. Dunham, Nick Roberts, and Anthony Mezzacappa. thornado-hydro: A Discontinuous Galerkin Method for Supernova Hydrodynamics with Nuclear Equations of State. *ApJS*, 253(1):21, March 2021. doi: 10.3847/1538-4365/abd700.
- M. Paul Laiu, Eirik Endeve, Ran Chu, J. Austin Harris, and O. E. Bronson Messer. A dg-imex method for two-moment neutrino transport: Nonlinear solvers for neutrino-matter coupling*. *The Astrophysical Journal Supplement Series*, 253(2):52, apr 2021. doi: 10.3847/1538-4365/abe2a8. URL <https://dx.doi.org/10.3847/1538-4365/abe2a8>.
- J. R. Wilson, G. J. Mathews, and P. Marronetti. Relativistic numerical model for close neutron-star binaries. *Phys. Rev. D*, 54(2):1317–1331, July 1996. doi: 10.1103/PhysRevD.54.1317.
- Isabel Cordero-Carrión, Pablo Cerdá-Durán, Harald Dimmelmeier, José Luis Jaramillo, Jérôme Novak, and Ericourgoulhon. Improved constrained scheme for the Einstein equations: An approach to the uniqueness issue. *Phys. Rev. D*, 79(2):024017, January 2009. doi: 10.1103/PhysRevD.79.024017.
- Sigal Gottlieb, Chi-Wang Shu, and Eitan Tadmor. Strong Stability-Preserving High-Order Time Discretization Methods. *SIAM Review*, 43(1):89–112, January 2001. doi: 10.1137/S003614450036757X.
- L. Pareschi and G. Russo. *Journal of Scientific Computing*, 25:129–155, 2005. doi: 10.1007/BF02728986.
- Guosheng Fu and Chi-Wang Shu. A new troubled-cell indicator for discontinuous Galerkin methods for hyperbolic conservation laws. *Journal of Computational Physics*, 347:305–327, October 2017. doi: 10.1016/j.jcp.2017.06.046.
- Tong Qin, Chi-Wang Shu, and Yang Yang. Bound-preserving discontinuous Galerkin methods for relativistic hydrodynamics. *Journal of Computational Physics*, 315:323–347, June 2016. doi: 10.1016/j.jcp.2016.02.079.
- Samuel J. Dunham, Eirik Endeve, Anthony Mezzacappa, John M. Blondin, Jesse Buffaloe, and Kelly Holley-Bockelmann. A Parametric Study of the SASI Comparing General Relativistic and Nonrelativistic Treatments. *ApJ*, 964(1):38, March 2024. doi: 10.3847/1538-4357/ad206c.
- Kevin Schaal, Andreas Bauer, Praveen Chandrasekar, Rüdiger Pakmor, Christian Klingenberg, and Volker Springel. Astrophysical hydrodynamics with a high-order discontinuous Galerkin scheme and adaptive mesh refinement. *MNRAS*, 453(4):4278–4300, November 2015. doi: 10.1093/mnras/stv1859.
- L. Del Zanna and N. Bucciantini. An efficient shock-capturing central-type scheme for multidimensional relativistic flows. I. Hydrodynamics. *A&A*, 390:1177–1186, August 2002. doi: 10.1051/0004-6361:20020776.

Summary

Can run multi-D pure hydro problems in GR with AMR



Can run 1D hydro+self-gravity problems in GR with AMR

Working on coupling GR transport to existing hydro+gravity modules