




20.1 Rate covariance matrix for the whole tree using all taxa (45 species), transforming inapplicable states to zeroes. Covariances scaled to correlations. All characters estimated simultaneously under Brownian Motion.




20.2 Rate covariance matrix for the whole tree using only taxa without inapplicable states (24 species). Covariances scaled to correlations. All characters estimated simultaneously under Brownian Motion.




20.3 Rate covariance matrix for the whole tree using only taxa with diet data (22 species), transforming inapplicable states to zeroes. Covariances scaled to correlations. All characters estimated simultaneously under Brownian Motion.



20.4 Best models (lowest AIC) supported in a pairwise character rate covariance analysis comparing correlated Brownian Motion models across the five selective regimes. Selective regimes were mapped onto the tree (22 species with diet data) using an ancestral state reconstruction of the feeding guilds. Blank cells represent computationally singular contrasts.



20.5 Pairwise estimated rate covariance matrices across the five selective regimes, using only taxa with diet data (22 species). Inapplicable states transformed to zeroes. Covariances scaled to correlations. Selective regimes were mapped onto the tree (22 species with diet data) using an ancestral state reconstruction of the feeding guilds. Question marks represent computationally singular contrasts.



20.6 Scaled differences between the regime-specific covariance matrices in Appendix 20.5 and the whole tree covariance matrix.

