




# fearmem\_report

## recognition memory

### non-responses

Starting with rigorous outlier detection. We had previously identified some subjects to exclude, but that was over a year ago so I wanted to start from scratch.

First, I started by looking at the % of missing data for each subject - trials they didn't actually respond.



subject	missing
20	0.179167
110	0.091667
120	0.229167


Above are the subjects who had >5% missing data.

Based on this info, I am excluding subjects 20 and 120. These subjects are not present in any further data or graphs

## poor memory

Next, I examined the corrected recognition for each group individually for outliers

Using boxplots, there were no outliers in the healthy group




Below are all points  $<.05$  corrected recognition. The only thing that stands out to me is subject 18, as they have poor memory for CS+ acquisition, should be the best memory for the entire experiment. Based on this, subject 18 is also excluded.

group	subject	condition	encode_phase	cr
healthy	18	CS+	acquisition	0.041667
healthy	3	CS-	acquisition	0.000000
healthy	13	CS+	extinction	0.041667
healthy	23	CS+	extinction	0.000000
healthy	16	CS-	extinction	0.020833

## PTSS group

Looking at our ptsd group I did not find any more suspect data



There were participants with low CR, but nothing to warrant exclusion.

group	subject	condition	encode_phase	cr
ptsd	101	CS-	baseline	0.000000
ptsd	102	CS+	extinction	0.020833
ptsd	116	CS+	extinction	0.020833
ptsd	116	CS-	extinction	0.041667

This leaves us with **N=22** in the healthy group, and **N=23** in the PTSS group.

## Normality of the data

Before running statistics, I checked the assumption of normality for corrected recognition in both groups separately, using a Shapiro-Wilk test.

group	W	pval	normal
healthy	0.961719	0.000910	False
ptsd	0.960665	0.000533	False

Both groups were determined to be non-normal, so I will be employing non-parametric statistics to analyze the recognition memory data.


## Analysis

I started with a full mixed ANOVA, using permutation tests to obtain p-values with 10,000 iterations, with a minimum possible p-value of 0.0001

Group was entered as a between-subjects factor, and Condition (CS+/-) and Encoding Phase were entered as within-subjects factors.

As previously reported, there were significant main effects of Condition ( $F(1,43) = 28.95$ ,  $P_{perm} = 0.0001$ ) Encoding Phase ( $F(2,86) = 19.28$ ,  $P_{perm} = 0.0001$ ) and a Condition X Encoding Phase interaction ( $F(2,86) = 9.15$ ,  $P_{perm} = 0.0005$ )

Next, I investigated the emotional enhancement ( $CS+ > CS-$ ). I've graphed it here as a difference score just for ease of seeing the enhancement effect - publication graphics will be full data.



For follow up comparisons, I used wilcoxon signed-rank tests, which is a non-parametric version of a paired t-test

**As we have a very specific directional hypothesis, I used 1-sided p-values.**

RBC = non-parametric effect size (rank-biserial correlation)

CLES = common language effect size

Retroactive effect: Baseline  $CS+ > CS-$

	W-val	tail	p-val	RBC	CLES
Healthy	183.0	greater	0.009903	0.584416	0.584711
PTSD	194.0	greater	0.045647	0.405797	0.60397

Emotional enhancement: Acquisition CS+ > CS-

	W-val	tail	p-val	RBC	CLES
Healthy	209.0	greater	0.000613	0.809524	0.747934
PTSD	229.0	greater	0.000464	0.810277	0.689981

Proactive effect: Extinction CS+ > CS-

	W-val	tail	p-val	RBC	CLES
Healthy	179.5	greater	0.013642	0.554113	0.606405
PTSD	196.5	greater	0.0388	0.423913	0.608696

**Thus, when using appropriate statistics (i.e. non-parametric and motivated by *a priori* hypothesis), we find that there is increased memory for CS+ relative to CS- for all phases and in both groups.**

## source memory

Carrying over our exclusions from the recognition memory, we have **N=17** in the healthy group and **N=17** in the PTSD group. We technically have the data from the 3 subs we excluded, since the source memory is self-paced they did answer everything. It could be interesting to see what they guessed, but for now just keeping them out.

## normality


Again before analysis I checked the overall normality of the source memory data.

	W	pval	normal
healthy	0.965078	9.845692e-07	False
ptsd	0.964084	7.086799e-07	False

Again, we are dealing with non-normal data.

## analysis

Here is source memory for items that were encoded during Baseline, broken down by CS type and Response Phase (the phase they said during the source memory test)




Running a full mixed ANOVA as before, we find there is a significant main effect of Response Phase ( $F(2,64) = 3.39$ ,  $P_{perm} = 0.0422$ ) and a Condition by Response Phase interaction ( $F(2,64) = 10.29$ ,  $P_{perm} = 0.0001$ )

I again using 1-sided Wilcoxon tests to determine if the proportion of CS+ items mis-sourced to acquisition was higher than CS- in each group.

	W-val	tail	p-val	RBC	CLES
Healthy	117.0	greater	0.006044	0.720588	0.688581
PTSD	114.0	greater	0.009209	0.676471	0.65917

In both groups CS+B were misplaced as being from acquisition more compared to CS-B

BUT - How does memory factor in? Here is the same graph, but now **Proportion** refers to the proportion of remembered items, which varies per subject.



Again main effect of Response Phase, and significant interaction of Condition X Response Phase

Follow up comparisons CS+B vs. CS-B responded to as from acquisition




	W-val	tail	p-val	RBC	CLES
Healthy	121.0	greater	0.003317	0.779412	0.66436
PTSD	111.0	greater	0.053754	0.45098	0.653979

So the effect is still significant for Healthy, but NOT for PTSS. Although it is trending.

We know that each subject has different hit rates for each condition and phase, so a more rigorous analysis should incorporate mixed effect modeling. But I'm really not sure how to approach it with this data, as we have binary memory status (hit vs. miss) and then three alternate choice data. I've spent a while thinking about this but there is not an "obvious" model to me. Please let me know what you think

Next we can do the same things but for extinction, looking at the proactive effect




Main effect of response Phase, Condition by Response Phase interaction

Again the critical comparisons here are CS+E vs CS-E that people endorse as being from acquisition

	W-val	tail	p-val	RBC	CLES
healthy	119.0	greater	0.004481	0.75	0.780277
PTSS	61.0	greater	0.045778	0.564103	0.704152

Significant in both groups, but only just barely in PTSS

Lets look at just the items that were remembered:



Main effect of response phase only

Follow up tests:


	W-val	tail	p-val	RBC	CLES
Healthy	116.0	greater	0.032395	0.51634	0.67474
PTSS	76.5	greater	0.339533	0.125	0.544983

Effect is significant for healthy, but definitely NOT for PTSS. First major group difference. Although important to note that this is definitely being driven by PTSS endorsing more CS-E as being from acquisition. I didn't run any tests, but I'd bet good money there is no difference in the CS+E being endorsed as acquisition between the two groups in the above graph.

# Recognition X Source memory

Correlations seeing if differential (CS+ > CS-) source memory predicts differential recognition memory. I'm not sure if restricting analysis to just remembered items is really appropriate here or not, so I'm showing it both ways.


Starting with Baseline. X-axis is the difference in CS+B - CS-B endorsed as being from acquisition, against differential corrected recognition for all of baseline on the Y-axis



healthy:  $r = 0.41$ ,  $p = 0.09$   
ptss:  $r = 0.31$ ,  $p = 0.22$

Trending in Healthy, NS in PTSS

Here is the same thing, using the difference in proportion of remembered items:




```
healthy: r = 0.25, p = 0.32
ptsd:    r = 0.45, p = 0.069
```

So here, NS in healthy, trending in PTSS

What this tells me is that there is probably something interesting going on with recognition memory here, but we need a more direct test to tease it out.

Lastly for now, here is the same gambit for extinction.


X-axis is the difference in CS+E - CS-E endorsed as being from acquisition, against differential corrected recognition for all of extinction on the Y-axis



healthy:  $r = 0.46$ ,  $p = 0.095$   
 ptss:  $r = 0.24$ ,  $p = 0.35$

So trending in Healthy, NS in PTSS

Finally, same as above but using only remembered items for the source memory:



```
healthy: r = 0.18, p = 0.48  
ptss:   r = 0.15, p = 0.55
```

Neither group is significant.

So in sum, I think we need to run a multinomial logistic regression in order to better relate performance on the 3-AFC source memory test to recognition memory performance.